

## Problem assignment 1.

Algebraic Geometry and Commutative Algebra

Joseph Bernstein

February 21, 2011.

**A remark on problems in different areas.** In my assignments I will try to single out problems that are not directly related to algebraic geometry. For example sign (CA) signifies a problem (or a definition) from commutative algebra, (LA) stands for linear algebra, (Top) for topology.

**A remark on different kinds of problems.** In all my home assignments I will use the following system.

The problems without marking are just exercises. You have to convince yourself that you can do them but it is not necessary to write them down (if you have difficulties with one of these problems ask me or Sasha).

The problems marked by [P] you should hand in for grading.

The sign (\*) marks more difficult problems.

The sign ( $\nabla$ ) marks more challenging and more interesting problems which are related to some interesting subjects. They are not always directly needed in the course, but I definitely advise you to think about these problems.

**Remark.** In this assignment you can freely use the Nullstellensatz. Please specify every time you use it (for example put the sign (NS)). Check for yourself that you really need it for your argument to work.

**1.** (i) Let  $A$  be a finitely generated  $k$ -algebra with no nilpotents. Consider the set  $X = \text{Mor}_{k\text{-alg}}(A, k)$ .

Show that  $A$  embeds into the algebra  $\mathcal{F}(X)$  of  $k$ -valued functions on  $X$  and that the pair  $(X, A)$  is an affine algebraic variety.

(ii) If  $X, Y$  are affine algebraic varieties, show that

$$\text{Mor}(X, Y) = \text{Mor}_{k\text{-alg}}(\mathcal{P}(Y), \mathcal{P}(X)).$$

**Definition.** (CA) Let  $A$  be a ring (commutative, with 1). Given an element  $f \in A$  we define the **localization** of  $A$  with respect to  $f$  to be the ring  $A_f = A[t]/(ft - 1)A[t]$ .

**2.** (CA) (i) Show that we can define the ring  $A_f$  as a set of fractions  $\frac{a}{f^k}$  modulo the following equivalence relation:  $\frac{a}{f^k}$  is equivalent to  $\frac{b}{f^l}$  iff for some  $N$  we have  $f^N f^l a = f^N f^k b$

(ii) Describe the kernel of the canonical morphism  $i : A \rightarrow A_f$ .

In what cases the localized ring  $A_f$  is trivial, i.e. consists of one element 0 ?

(iii) Show that if  $A$  has no nilpotents (resp. no zero divisors), then  $A_f$  has no nilpotents (resp. no zero divisors).

[P] **3.** Let  $(X, \mathcal{P}(X))$  be an affine algebraic variety. Pick an element  $f$  in  $\mathcal{P}(X)$ . Show that the algebra  $\mathcal{P}(X)_f$  is canonically isomorphic to the algebra of polynomial functions on the basic open subset  $X_f$ .

[P] 4. Let  $X$  be an affine algebraic variety and  $B$  a subset of  $X$ . Suppose we know that  $B$  is a basic open subset, i.e. it has a form  $B = X_f$  for some function  $f \in \mathcal{P}(X)$ .

Show how to describe the subalgebra  $\mathcal{P}(B) \subset \mathcal{F}(B)$  in terms of the set  $B$ , without knowing the function  $f$  that defines it.

5. Consider the subvariety  $Y \subset \mathbf{A}^2$  defined by one equation  $x^2 - y^3 = 0$ .

Describe a morphism  $\nu : \mathbf{A}^1 \rightarrow Y$  which is bijective. Show that it is a homeomorphism, but not an isomorphism of algebraic varieties.

[P] 6. Let  $K$  be some field. Consider a system of polynomial equations  $\Xi = (P_\alpha)$ , where  $P_\alpha$  are polynomials in variables  $x_1, \dots, x_n$ .

Suppose we know that there exists a field extension  $\Omega$  such that the system of equations  $\Xi$  has some solution in  $\Omega$ , i.e. there exists a collection of elements  $x_i \in \Omega$  that satisfy all the equations in  $\Xi$ .

Show that the system  $\Xi$  has a solution in the field  $k = \bar{K}$  - algebraic closure of  $K$ .

7. Let  $X$  be an affine algebraic variety. Denote by  $\mathcal{B}$  the family of basic open subsets of  $X$ .

Show that  $\mathcal{B}$  forms a base of a topology and that the topology defined by  $\mathcal{B}$  is the Zariski topology on  $X$ .

**Definition.** A topological space  $X$  is called **quasicompact** if any open covering  $\{U_\alpha\}$  of  $X$  has finite subcovering.

[P] 8. Show that any affine algebraic variety is quasicompact in Zariski topology.