

Blowing up the power of a singular cardinal of uncountable cofinality.

Moti Gitik*

February 13, 2018

We will present here a new method for blowing up the power of a singular cardinal which differs from those used in [1] and in [2] to deal with cofinality ω . The advantage of the present technique is that it generalizes to singular cardinals of uncountable cofinality, which was open.

Let us start with countable cofinality.

1 Blowing up the power of a singular cardinal of cofinality ω .

Let $\langle \kappa_n \mid n < \omega \rangle$ be an increasing sequence of cardinals and $\langle E_n \mid n < \omega \rangle$ such that for every $n < \omega$

1. $E(n)$ is a $(\kappa_n, \kappa_\omega^{++})$ -extender,
i.e. $j_{E(n)} : V \rightarrow M_{E(n)} \simeq \text{Ult}(V, E(n))$, $\text{crit}(j_{E(n)}) = \kappa_n$, $j_{E(n)}(\kappa_n) > \kappa_\omega^{++}$,
 $M_{E(n)} \supseteq V_{\kappa_\omega+2, \kappa_n} M_{E(n)} \subseteq M_{E(n)}$;
2. $E(n) \triangleleft E(n+1)$,
where $\kappa_\omega = \bigcup_{n < \omega} \kappa_n$.

Denote by $\mathcal{P}(n)$ the one element extender based Prikry forcing with $E(n)$. We would like to combine $\mathcal{P}(n)$'s together. It would be a kind of Magidor product, but will involve restrictions and reflections. Namely, if for some $n < \omega$ a non-direct extension is made in $\mathcal{P}(n)$, then we will restrict each $E(m)$, $m < n$ to the corresponding member of the Prikry

*The work was partially supported by Israel Science Foundation Grant No. 58/14. We are grateful to Carmi Merimovich for reading a draft of the paper and specially for the inspiration we got from his works on extender based forcings.

sequence for κ_n and reflect the information the condition contains about coordinates $m < n$ below κ_n .

Let us start with a simpler situation where instead of ω extenders we have only two.

1.1 A single extender.

Let us describe a variation of one element extender based Prikry forcing that will be used here. It will be very close to those of C. Merimovich [7]. A difference will be that sequences inside conditions will be either empty or of the length one only.

Let E be a (κ, λ) -extender. Define the corresponding forcing \mathcal{P}_E .

Let $d \subseteq \lambda \setminus \kappa$ of cardinality at most κ . Define a κ -ultrafilter $E(d)$ on $[d \times \kappa]^{<\kappa}$ as follows:

$$X \in E(d) \Leftrightarrow \{\langle j_E(\alpha), \alpha \rangle \mid \alpha \in d\} \in j_E(X).$$

Actually, $E(d)$ concentrates on a smaller set called $OB(d)$ in [7].

The advantage of using $E(d)$ is that once A is typical set of $E(d)$ -measure one and $a \in A$, then a is of the form $\langle \langle \alpha_\xi, \beta_\xi \rangle \mid \xi < \rho \rangle$, where

1. $\rho < \kappa$,
2. $\text{dom}(a) = \{\alpha_\xi \mid \xi < \rho\} \subseteq d$,
3. $\beta_\xi < \kappa$, for every $\xi < \rho$.

So, already a measure one set provides an explicit connection between elements of Prikry sequences and the measures to which they belong.

We assume further that always $\langle \alpha_\xi \mid \xi < \rho \rangle$ and $\langle \beta_\xi \mid \xi < \rho \rangle$ are strictly increasing sequences of ordinals.

Definition 1.1 Let \mathcal{P}_E^* be the set of all functions f such that

1. $\text{dom}(f) \subseteq \lambda \setminus \kappa$ of cardinality at most κ ,
2. $\kappa \in \text{dom}(f)$,
3. for every $\alpha \in \text{dom}(f)$, $f(\alpha)$ is either empty or a one element sequence which consists of an element of κ .

Definition 1.2 Let $f, g \in \mathcal{P}_E^*$. Set $f \geq^* g$ iff $f \subseteq g$.

Definition 1.3 (One element extension) Let $f \in \mathcal{P}_E^*$ and $\vec{v} \in [\text{dom}(f) \times \kappa]^{<\kappa}$. Define $g = f_{\langle \vec{v} \rangle} \in \mathcal{P}_E^*$ as follows:

1. $\text{dom}(g) = \text{dom}(f)$,
2. for every $\alpha \in \text{dom}(g)$,
$$g(\alpha) = \begin{cases} \langle \vec{v}(\alpha) \rangle, & \text{if } \alpha \in \text{dom}(\vec{v}) \text{ and } f(\alpha) \text{ is empty sequence;} \\ \langle \vec{v}(\alpha) \rangle, & \text{if } \alpha \in \text{dom}(\vec{v}), f(\alpha) \text{ is not empty and } \vec{v}(\alpha) > f(\alpha); \\ f(\alpha), & \text{otherwise.} \end{cases}$$

The difference from the original definition by Merimovich in [7], is that we do not keep $f(\alpha)$ if $\vec{v}(\alpha) > f(\alpha)$, but rather replace $f(\alpha)$ by $\vec{v}(\alpha)$.

Define now the pure part \mathcal{P}_E^0 of the main forcing \mathcal{P}_E .

Definition 1.4 A pure condition $p \in \mathcal{P}_E^0$ is of the form $\langle f, A \rangle$, where

1. $f \in \mathcal{P}_E^*$,
2. $f(\kappa)$ is the empty sequence,
3. $A \in E(\text{dom}(f))$.

Define the order on \mathcal{P}_E^0 as follows:

Definition 1.5 Let $p = \langle f, A \rangle, q = \langle g, B \rangle \in \mathcal{P}_E^0$. Set $p \geq^* q$ iff

1. $f \geq^* g$ in \mathcal{P}_E^* ,
2. $A \upharpoonright \text{dom}(g) \subseteq B$.

The forcing \mathcal{P}_E will be the union of \mathcal{P}_E^0 with

$$\{f \in \mathcal{P}_E^* \mid f(\kappa) \neq \langle \rangle\}.$$

The direct order extension will be just the union of \leq^* orders of both parts. Let us define the forcing order \leq on \mathcal{P} . We do this by defining one element extensions of members of \mathcal{P}_E^0 .

Definition 1.6 Let $p = \langle f, A \rangle$ be in \mathcal{P}_E^0 and $\vec{v} \in A$. Define $p \hat{\sim} \vec{v} \in \mathcal{P}_E^*$ to be $f_{\langle \vec{v} \rangle}$.

Definition 1.7 Let $p = \langle f, A \rangle$ be in \mathcal{P}_E^0 and g be in \mathcal{P}_E^* . Set $p \leq g$ iff there is $\vec{v} \in A$ such that $f_{\langle \vec{v} \rangle} \leq^* g$.

The next lemma follows from the definitions:

Lemma 1.8 *The forcing $\langle \mathcal{P}_E, \leq \rangle$ is equivalent to the Cohen forcing for adding λ -many Cohen subsets to κ^+ .*

However, more can be deduced:

Lemma 1.9 *$\langle \mathcal{P}, \leq, \leq^* \rangle$ is a Prikry type forcing notion.*

Proof. Let us sketch the basic argument following Merimovich presentation [7].

Let $p = \langle f^p, A^p \rangle \in \mathcal{P}_E^0$ and σ be a statement of the forcing language.

We would like to find a direct extension of p which decides σ . Suppose that there is no such extension.

Proceed as in 3.12 of [6]. Construct by induction an increasing chain of elementary submodels $\langle N_\xi \mid \xi < \kappa \rangle$ of H_χ , for χ large enough, and a sequence $\langle f_\xi \mid \xi < \kappa \rangle$ of members of \mathcal{P}_E^* , such that

1. $p, \mathcal{P}_E, \sigma \in N_0$,
2. $N_0 \supseteq \kappa$,
3. for every $\xi < \kappa$,
 - (a) $|N_\xi| = \kappa$,
 - (b) ${}^{\kappa >} N_\xi \subseteq N_\xi$,
 - (c) $\langle f_\zeta \mid \zeta < \xi \rangle \in N_\xi$,
 - (d) $f_\xi \in \bigcap \{D' \in N_\xi \mid D' \text{ is a dense open subset of } \mathcal{P}_E^* \text{ above } f^p\}$,
 - (e) $f^p \leq^* f_0$,
 - (f) $f_\xi \geq^* f_\zeta$, for every $\zeta < \xi$.

Set $N = \bigcup_{\xi < \kappa} N_\xi$ and $f^* = \bigcup \{f_\xi \mid \xi < \kappa\}$.¹ to construct Let $A \subseteq [\text{dom}(f^*) \times \kappa]^{<\kappa}$ be such that

- $A \upharpoonright \text{dom}(f^p) \subseteq A^p$,
- $A \in E(\text{dom}(f^*))$.

¹Carmi Merimovich pointed out that there is no need here in elementary chain of models and it is possible to define N directly. This observation applies also to our further constructions.

Note that $A \subseteq N$, since $\text{dom}(f^*) \subseteq N$, and so, $[\text{dom}(f^*) \times \kappa]^{<\kappa} \subseteq N$.

Let $\vec{\nu} \in A$.

Define $D_{\vec{\nu}}$ to be the set of all $f \in \mathcal{P}_E^*$, $f \geq f^p$ such that

$$f_{\vec{\nu}} \parallel \sigma.$$

Then $D_{\vec{\nu}}$ is a dense open subset of \mathcal{P}_E^* above f^p .

It is definable with parameters in N , hence $D_{\vec{\nu}} \in N$.

Then, $f^* \in D_{\vec{\nu}}$.

Shrink now A to $A^* \in E(\text{dom}(f^*))$, if necessary, such that for every $\vec{\nu}, \vec{\nu}'$ inside we will have

$$f_{\vec{\nu}}^* \Vdash \sigma \text{ iff } f_{\vec{\nu}'}^* \Vdash \sigma.$$

Suppose that for every $\vec{\nu} \in A^*$, $f_{\vec{\nu}}^* \Vdash \sigma$.

Now, we claim that already $\langle f^*, A^* \rangle \Vdash \sigma$.

Suppose otherwise. Then there is $g \geq \langle f^*, A^* \rangle$ which forces $\neg\sigma$. Then for some $\vec{\nu} \in A^*$, $g \geq f_{\vec{\nu}}^*$, by Definition 1.7. But $f^* \in D_{\vec{\nu}}$, hence already $f_{\vec{\nu}}^* \Vdash \neg\sigma$, which is impossible by the choice of A^* .

Contradiction.

□

1.2 Two extenders.

We deal now with $E(0) \triangleleft E(1)$.

Let $\mathcal{P}_{E(i)}^*$, $\mathcal{P}_{E(i)}$, $\mathcal{P}_{E(i)}^0$, $i < 2$ be as defined in the previous subsection.

Define first components of the main forcing $\mathcal{P}_{\langle E(0), E(1) \rangle}$.

Definition 1.10 The set of pure conditions $\mathcal{P}_{\langle E(0), E(1) \rangle}^{(0,0),(1,0)}$ consists of all pairs $\langle p(0), p(1) \rangle$ such that

1. $p(0) = \langle f^0, A^0 \rangle \in \mathcal{P}_{E(0)}^0$,
2. $p(1) = \langle f^1, A^1 \rangle \in \mathcal{P}_{E(1)}^0$,
3. $\text{dom}(f^0) \setminus \kappa_1 \subseteq \text{dom}(f^1)$,
4. for every $\alpha \in \text{dom}(f^0) \setminus \kappa_1$, if $f^1(\alpha)$ is not the empty sequence, then for every $\vec{\nu} \in A^1$, $\alpha \in \text{dom}(\vec{\nu})$ and $\vec{\nu}(\alpha) > f^1(\alpha)$.

The intuition behind this condition is that the current value $f^1(\alpha)$ may interfere with

values of one element Prikry sequences over κ_0 . Namely, with the α -th Prikry sequence over κ_0 . Now, if $\vec{v}(\alpha) > f^1(\alpha)$, then $f^1_{\vec{v}}(\alpha) = \vec{v}(\alpha)$, by Definition 1.3, and so, the value $f^1(\alpha)$ just disappears.

5. For every $\gamma \in \text{dom}(f^0) \cap \kappa_1, \vec{v} \in A^1$ and $\alpha \in \text{dom}(\vec{v}), \vec{v}(\alpha) > \gamma$.

Note that $|\text{dom}(f^0)| \leq \kappa_0$, so it is easy to arrange this.

6. For every $\vec{v} \in A^1$, the measures $E(0)(\text{dom}(f^0))$ and

$E(0)((\text{dom}(f^0) \cap \kappa_1) \cup \{\vec{v}(\alpha) \mid \alpha \in \text{dom}(f^0) \setminus \kappa_1\})$ are basically the same in the following sense:

$$X \in E(0)(\text{dom}(f^0)) \text{ iff } X^{ref} \in E(0)((\text{dom}(f^0) \cap \kappa_1) \cup \{\vec{v}(\alpha) \mid \alpha \in \text{dom}(f^0) \setminus \kappa_1\}),$$

where

$$X^{ref} = \{(\alpha, \beta) \in X \mid \alpha < \kappa_1\} \cup \{(\vec{v}(\alpha), \beta) \mid (\alpha, \beta) \in X, \alpha \geq \kappa_1\}.$$

Note that this property is true in the ultrapower by $E(1)$, so it holds on a set of measure one, as well.

Turn now to non-pure extensions.

First consider the situation with non-pure part over κ_0 .

Definition 1.11 The set of conditions $\mathcal{P}_{\langle E(0), E(1) \rangle}^{(1,0)}$ consists of all pairs $\langle f^0, p(1) \rangle$ such that

1. $f^0 \in \mathcal{P}_{E(0)}^*$,
2. $p(1) = \langle f^1, A^1 \rangle \in \mathcal{P}_{E(1)}$,
3. $\text{dom}(f^0) \setminus \kappa_1 \subseteq \text{dom}(f^1)$,
4. for every $\alpha \in \text{dom}(f^0) \setminus \kappa_1$, if $f^1(\alpha)$ is not the empty sequence, then for every $\vec{v} \in A^1$, $\alpha \in \text{dom}(\vec{v})$ and $\vec{v}(\alpha) > f^1(\alpha)$,
5. for every $\gamma \in \text{dom}(f^0) \cap \kappa_1, \vec{v} \in A^1$ and $\alpha \in \text{dom}(\vec{v}), \vec{v}(\alpha) > \gamma$.

Define conditions with a pure part over κ_0 and a non-pure over κ_1 .

Assume for simplicity that there is $h_\lambda : \kappa_1 \rightarrow \kappa_1$ such that $j_{E(1)}(h_\lambda)(\kappa_1) = \lambda$.

Definition 1.12 The set of conditions $\mathcal{P}_{\langle E(0), E(1) \rangle}^{(0,0)}$ consists of all pairs $\langle p(0), f^1 \rangle$ such that

1. $f^1 \in \mathcal{P}_{E(1)}^*$,

2. $f^1(\kappa_1)$ is non-empty,
3. $p(0) \in \mathcal{P}_{E(0) \upharpoonright h_\lambda(f^1(\kappa_1))}$. The meaning is that if the value of the Prikry sequence for the normal measure of $E(1)$ is decided, then we cut $E(0)$ to the reflection of λ below κ_1 , i.e. to $h_\lambda(f^1(\kappa_1))$.

Define now a completely non-pure part of the forcing.

Definition 1.13 The set of conditions $\mathcal{P}_{\langle E(0), E(1) \rangle}^*$ consists of all pairs $\langle f^0, f^1 \rangle$ such that

1. $f^1 \in \mathcal{P}_{E(1)}^*$,
2. $f^1(\kappa_1)$ is non-empty,
3. $f^0 \in \mathcal{P}_{E(0)}^*$,
4. $f^0(\kappa_0)$ is non-empty,
5. $\text{dom}(f^0) \subseteq h_\lambda(f^1(\kappa_1))$.

The meaning is that if the value of the Prikry sequence for the normal measure of $E(1)$ is decided, then we add only $h_\lambda(f^1(\kappa_1))$ Cohen subsets to κ_0^+ .

Now let us put everything together.

Definition 1.14 $\mathcal{P}_{\langle E(0), E(1) \rangle} = \mathcal{P}_{\langle E(0), E(1) \rangle}^{(0,0),(1,0)} \cup \mathcal{P}_{\langle E(0), E(1) \rangle}^{(1,0)} \cup \mathcal{P}_{\langle E(0), E(1) \rangle}^{(0,0)} \cup \mathcal{P}_{\langle E(0), E(1) \rangle}^*$.

Define the orders \leq, \leq^* over $\mathcal{P}_{\langle E(0), E(1) \rangle}$.

\leq^* is just the union of the orders at each of the components.

Let us give now the main definition.

Definition 1.15 Let $p, q \in \mathcal{P}_{\langle E(0), E(1) \rangle}$. If p, q are in the same component, then set $p \geq q$ iff $p \geq^* q$. Suppose that they are in different components.

Split into cases.

1. Suppose that $q \in \mathcal{P}_{\langle E(0), E(1) \rangle}^{(0,0),(1,0)}$, i.e. in the pure part of $\mathcal{P}_{\langle E(0), E(1) \rangle}$, $p \in \mathcal{P}_{\langle E(0), E(1) \rangle}^{(1,0)}$, i.e. only the part of p over κ_1 is a pure condition.

Let then $q = \langle \langle g^0, B^0 \rangle, \langle g^1, B^1 \rangle \rangle, p = \langle f^0, \langle f^1, A^1 \rangle \rangle$.

Set $p \geq q$ iff $f^0 \geq \langle g^0, B^0 \rangle$ in $\mathcal{P}_{E(0)}$ and $\langle f^1, A^1 \rangle \geq^* \langle g^1, B^1 \rangle$ in $\mathcal{P}_{E(1)}$.

2. Suppose that $q \in \mathcal{P}_{\langle E(0), E(1) \rangle}^{(0,0)}$, i.e. in the part over κ_0 is pure and those over κ_1 is not pure, $p \in \mathcal{P}_{\langle E(0), E(1) \rangle}^*$, i.e. p is a completely non-pure condition.
 Let then $q = \langle \langle g^0, B^0 \rangle, g^1 \rangle$ and $p = \langle f^0, f^1 \rangle$.
 Set $p \geq q$ iff $f^0 \geq \langle g^0, B^0 \rangle$ in $\mathcal{P}_{E(0)}$ and $f^1 \geq g^1$ in $\mathcal{P}_{E(1)}$.

3. (Principal case 1.)

Suppose that $q \in \mathcal{P}_{\langle E(0), E(1) \rangle}^{(1,0)}$, i.e. in the part over κ_1 is pure and those over κ_0 is not pure, $p \in \mathcal{P}_{\langle E(0), E(1) \rangle}^*$, i.e. p is a completely non-pure condition.
 Let then $q = \langle g^0, \langle g^1, B^1 \rangle \rangle$ and $p = \langle f^0, f^1 \rangle$.
 Set $p \geq q$ iff $f^1 \geq \langle g^1, B^1 \rangle$ in $\mathcal{P}_{E(1)}$ and $f^0 \geq (g^0)^{ref}$ in $\mathcal{P}_{E(0) | h_\lambda(f^1(\kappa_1))}$, where $(g^0)^{ref}$ the reflection of g^0 below κ_1 is defined as follows:

- (a) $\text{dom}((g^0)^{ref}) = (\text{dom}(g^0) \cap \kappa_1) \cup \{f^1(\alpha) \mid \alpha \in \text{dom}(g^0) \setminus \kappa_1\}$,
- (b) for every $\alpha \in \text{dom}(g^0) \cap \kappa_1 = \text{dom}(g^0) \cap \text{dom}((g^0)^{ref})$, $(g^0)^{ref}(\alpha) = g^0(\alpha)$,
- (c) for every $\alpha \in \text{dom}(g^0) \setminus \kappa_1$, $(g^0)^{ref}(f^1(\alpha)) = g^0(\alpha)$.

It is crucial here that $f^1 \upharpoonright (\text{dom}(g^0) \setminus \kappa_1)$ is one to one and the values there are above $\text{rng}(g^0) \cap \kappa_1$.

This follows by conditions (4),(5) of Definitions 1.10,1.11.

4. (Principal case 2.)

Suppose that $q \in \mathcal{P}_{\langle E(0), E(1) \rangle}^{(0,0),(1,0)}$, i.e. both parts are pure, $p \in \mathcal{P}_{\langle E(0), E(1) \rangle}^{(0,0)}$, i.e. only the part over κ_0 is pure.
 Let then $q = \langle \langle g^0, B^0 \rangle, \langle g^1, B^1 \rangle \rangle$ and $p = \langle \langle f^0, A^0 \rangle, f^1 \rangle$.
 Set $p \geq q$ iff $f^1 \geq \langle g^1, B^1 \rangle$ in $\mathcal{P}_{E(1)}$ and $\langle f^0, A^0 \rangle \geq (\langle g^0, B^0 \rangle)^{ref}$ in $\mathcal{P}_{E(0) | h_\lambda(f^1(\kappa_1))}$, where $(\langle g^0, B^0 \rangle)^{ref}$ the reflection of $\langle g^0, B^0 \rangle$ below κ_1 is defined as follows:

- (a) $\text{dom}((\langle g^0, B^0 \rangle)^{ref}) = (\text{dom}(g^0) \cap \kappa_1) \cup \{f^1(\alpha) \mid \alpha \in \text{dom}(g^0) \setminus \kappa_1\}$,
- (b) for every $\alpha \in \text{dom}(g^0) \cap \kappa_1 = \text{dom}(g^0) \cap \text{dom}((\langle g^0, B^0 \rangle)^{ref})$, $(\langle g^0, B^0 \rangle)^{ref}(\alpha) = g^0(\alpha)$,
- (c) for every $\alpha \in \text{dom}(g^0) \setminus \kappa_1$, $(\langle g^0, B^0 \rangle)^{ref}(f^1(\alpha)) = g^0(\alpha)$.

Again, it is crucial here that $f^1 \upharpoonright (\text{dom}(g^0) \setminus \kappa_1)$ is one to one and the values there are above $\text{rng}(g^0) \cap \kappa_1$, and this follows by conditions (4),(5) of Definitions 1.10,1.11.

One more crucial observation here is that the measure $(E(0))(\text{dom}(g^0)$, to which B^0 belongs, reflects to basically the same measure,

It follows by (6) of Definitions 1.10.

- (d) $A^0 \upharpoonright \text{dom}((g^0)^{ref}) \subseteq (B^0)^{ref}$, where $(B^0)^{ref} = \{\vec{v}^{ref} \mid \vec{v} \in B^0\}$ and if $\vec{v} = \langle \langle \alpha_\xi, \beta_\xi \rangle \mid \xi < \rho \rangle$, then $\vec{v}^{ref} = \langle \langle \alpha_\xi, \beta_\xi \rangle \mid \xi < \rho, \alpha < \kappa_1 \rangle \hat{\ } \langle \langle f^1(\alpha_\xi), \beta_\xi \rangle \mid \xi < \rho, \alpha \geq \kappa_1 \rangle$.

Denote further in this subsection $\mathcal{P}_{\langle E(0), E(1) \rangle}$ by just \mathcal{P} .

The next lemma follows from the definitions:

Lemma 1.16 *The forcing $\langle \mathcal{P}, \leq \rangle$ is equivalent to $\text{Cohen}(\kappa_0^+, \eta) \times \text{Cohen}(\kappa_1^+, \lambda)$, for some $\eta < \kappa_1$ which depends on choice of a non-pure condition for $\mathcal{P}_{E(1)}$.*

However, as usual, more can be deduced:

Lemma 1.17 *$\langle \mathcal{P}, \leq, \leq^* \rangle$ is a Prikry type forcing notion.*

Proof. The proof is similar to those of Lemma 1.9 (and in turn to those of Merimovich [7]).

Suppose otherwise.

Let $p \in \mathcal{P}$ be a pure condition and σ a statement of the forcing language which is undecided by pure extensions of p . Then p is of the form $\langle \langle f^{p^0}, A^{p^0} \rangle, \langle f^{p^1}, A^{p^1} \rangle \rangle$.

Proceed as in 3.12 of [6]. Construct by induction an increasing chain of elementary submodels $\langle N_\xi^1 \mid \xi < \kappa_1 \rangle$ of H_χ , for χ large enough, and a sequence $\langle f_\xi^1 \mid \xi < \kappa_1 \rangle$ of members of $\mathcal{P}_{E(1)}^*$, such that

1. $p, \mathcal{P}, \sigma \in N_0^1$,
2. $N_0^1 \supseteq \kappa_1$,
3. for every $\xi < \kappa_1$,
 - (a) $|N_\xi^1| = \kappa_1$,
 - (b) $\kappa_1 > N_\xi^1 \subseteq N_\xi^1$,
 - (c) $\langle f_\zeta^1 \mid \zeta < \xi \rangle \in N_\xi^1$,
 - (d) $f_\xi^1 \in \bigcap \{D' \in N_\xi^1 \mid D' \text{ is a dense open subset of } \mathcal{P}_{E(1)}^* \text{ above } f^{1p}\}$,
 - (e) $f^{1p} \leq^* f_0^1$,
 - (f) $f_\xi^1 \geq^* f_\zeta^1$, for every $\zeta < \xi$.

Set $N^1 = \bigcup_{\xi < \kappa_1} N_\xi^1$ and $f^{1*} = \bigcup \{f_\xi^1 \mid \xi < \kappa_1\}$. Let $A \subseteq [\text{dom}(f^{1*}) \times \kappa_1]^{< \kappa_1}$ be such that

- $A \upharpoonright \text{dom}(f^{p^1}) \subseteq A^{p^1}$,

- $A \in (E(1))(\text{dom}(f^{1*}))$.

Note that $A \subseteq N^1$, since $\text{dom}(f^{1*}) \subseteq N^1$, and so, $[\text{dom}(f^{1*}) \times \kappa_1]^{<\kappa_1} \subseteq N^1$.

Let $\vec{\nu} \in A$. Consider $\lambda_1^{\vec{\nu}} := h_\lambda(\vec{\nu}(\kappa_1))$, i.e. the cardinal below κ_1 that now corresponds to λ . Suppose for simplicity that $\text{dom}(f^{p_0}) \subseteq \lambda_1^{\vec{\nu}}$, otherwise just reflect the part above κ_1 below as in Definition 1.15.

Consider $\mathcal{P}_{E(0) \upharpoonright \lambda_1^{\vec{\nu}}}$. Clearly, it is contained and belongs to N^1 .

Let $\langle t_\xi \mid \xi < \lambda_1^{\vec{\nu}} \rangle$ be an enumeration of this forcing notion in N^1 .

Let $f \in \mathcal{P}_{E(1)}^*$, $f \geq^* f^{p_1}$.

Proceed by induction on $\xi < \lambda_1^{\vec{\nu}}$ and define an \leq^* -increasing sequence $\langle f_\xi \mid \xi < \lambda_1^{\vec{\nu}} \rangle$ of direct extensions of f such that, for every $\xi < \lambda_1^{\vec{\nu}}$, either

$$(1) \langle t_\xi, (f_\xi)_{\vec{\nu}} \rangle \parallel \sigma,$$

or

$$(2) \text{ for every } g \geq^* (f_\xi)_{\vec{\nu}}, \langle t_\xi, g \rangle \not\parallel \sigma.$$

Let $\bar{f} = \bigcup_{\xi < \lambda_1^{\vec{\nu}}} f_\xi$.

Then, for every $t \in \mathcal{P}_{E(0) \upharpoonright \lambda_1^{\vec{\nu}}}$ either

$$(1) \langle t, \bar{f}_{\vec{\nu}} \rangle \parallel \sigma,$$

or

$$(2) \text{ for every } g \geq^* \bar{f}_{\vec{\nu}}, \langle t, g \rangle \not\parallel \sigma.$$

Consider now the following statement of the forcing language of $\mathcal{P}_{E(0) \upharpoonright \lambda_1^{\vec{\nu}}}$:

$$\varphi \equiv \exists t \in \mathcal{G}(\langle t, \bar{f}_{\vec{\nu}} \rangle \parallel \sigma).$$

By the Prikry condition of the forcing $\mathcal{P}_{E(0) \upharpoonright \lambda_1^{\vec{\nu}}}$ (Lemma 1.9), there is $t^* \geq^* \langle f^{p_0}, A^{p_0} \rangle$ which decides φ .

Claim 1 $t^* \Vdash \varphi$.

Proof. Suppose otherwise. Then $t^* \Vdash \neg \varphi$. This means that whenever $t \in \mathcal{P}_{E(0) \upharpoonright \lambda_1^{\vec{\nu}}}$ and $t \geq t^*$, $\langle t, \bar{f}_{\vec{\nu}} \rangle \not\parallel \sigma$.

Pick now some $\langle t, g \rangle \in \mathcal{P}_{E(0), E(1)}$, $\langle t, g \rangle \geq \langle t^*, \bar{f}_{\vec{\nu}} \rangle$ which decides σ .

Then, for some $\xi < \lambda_1^{\vec{\nu}}$, $t = t_\xi$, and then, $\langle t, (f_\xi)_{\vec{\nu}} \rangle \parallel \sigma$. So, $\langle t, \bar{f}_{\vec{\nu}} \rangle \parallel \sigma$.

Contradiction.

□ of the claim.

Now use again the Prikry condition of the forcing $\mathcal{P}_{E(0)|\lambda_1^{\vec{v}}}$ to decide the following statement

$$\psi \equiv \exists t \in \mathcal{G}(\langle t, \bar{f}_{\vec{v}} \rangle \Vdash \sigma).$$

Let $t(\vec{v}, f) \geq^* t^*$ be a condition which decides ψ .

Assume that $t(\vec{v}, f) \Vdash \psi$. Then $\langle t(\vec{v}, f), \bar{f}_{\vec{v}} \rangle \Vdash \sigma$.

Define $D_{\vec{v}}$ to be the set of all $f \in \mathcal{P}_{E(1)}^*$, $f \geq^* f^{p1}$ such that

$$\langle t(\vec{v}, f), f_{\vec{v}} \rangle \Vdash \sigma.$$

The next claim follows now:

Claim 2 $D_{\vec{v}}$ is a dense open subset of $\mathcal{P}_{E(1)}^*$ above f^{p1} .

$D_{\vec{v}}$ is definable with parameters in N , hence $D_{\vec{v}} \in N$.

Then, $f^{1*} \in D_{\vec{v}}$, for every $\vec{v} \in A$.

So, $\langle t(\vec{v}, f^{1*}), f_{\vec{v}}^{1*} \rangle \Vdash \sigma$, for every $\vec{v} \in A$. Shrink A , if necessary, to a set

$A^{1*} \in (E(1))(\text{dom}(f^{1*}))$, such that for any two $\vec{v}, \vec{v}' \in A^{1*}$ the decision is the same, say σ is forced.

Consider now $\langle f^{1*}, A^{1*} \rangle$. It is a pure condition in $\mathcal{P}_{E(1)}$. Use the function $\vec{v} \mapsto t(\vec{v}, f^{1*})$ in order to get a pure condition in $\mathcal{P}_{E(0)}$, just use the one which this function represents in the ultrapower by $(E(1))(\text{dom}(f^{1*}))$.

Let us explain how do we naturally combine the result into a condition in $\mathcal{P}_{E(0), E(1)}$.

Let $t(\vec{v}, f^{1*}) = \langle f^{0\vec{v}}, A^{0\vec{v}} \rangle$, for every $\vec{v} \in A^{1*}$. Consider $f^{0\vec{v}}$. It is a set of at most κ_0 many pairs (α, β) , where $\alpha < \lambda_1^{\vec{v}} < \kappa_1$ and β is either the empty sequence or an ordinal $< \kappa_0$.

Shrinking A^{1*} if necessary, we can assume that there are x and $\kappa_0^* < \kappa_0^+$ such that for every $\vec{v}, \vec{v}' \in A^{1*}$ the following hold:

1. $\text{dom}(f^{0\vec{v}}) \cap \vec{v}(\kappa_1) = x$,
2. $\text{dom}(f^{0\vec{v}}) \setminus \vec{v}(\kappa_1) = \{\gamma_\tau^{\vec{v}} \mid \tau < \kappa_0^*\}$ is an increasing enumeration,
3. for every $\alpha \in x$, $f^{0\vec{v}}(\alpha) = f^{0\vec{v}'}(\alpha)$,
4. for every $\tau < \kappa_0^*$, $f^{0\vec{v}}(\gamma_\tau^{\vec{v}}) = f^{0\vec{v}'}(\gamma_\tau^{\vec{v}'})$

Consider, for every $\tau < \kappa_0^*$ a function s_τ on A^{1^*} defined by setting $s_\tau(\vec{\nu}) = \gamma_\tau^{\vec{\nu}}$.

Let

$$\gamma_\tau = j_{E(1)}(s_\tau)(\langle (j_{E(1)}(\alpha), \alpha) \mid \alpha \in \text{dom}(f^{1^*}) \rangle).$$

Extend now f^{1^*} to $f^{1^{**}}$ by adding all $\gamma_\tau, \tau < \kappa_0^*$ to its domain and setting $f^{1^{**}}(\gamma_\tau)$ to be the empty sequence whenever $\gamma_\tau \notin \text{dom}(f^{1^*})$.

Define $A^{1^{**}} \in E(1)(\text{dom}(f^{1^{**}}))$ as follows.

Set $\vec{\nu} \in A^{1^{**}}$ iff

1. $\vec{\nu} \upharpoonright \text{dom}(f^{1^*}) \in A^{1^*}$,
2. $\text{dom}(\vec{\nu}) \supseteq \{\gamma_\tau \mid \tau < \kappa_0^*\}$,
3. if $\gamma_\tau \in \text{dom}(f^{1^*})$ and $f^{1^*}(\gamma_\tau)$ is not the empty sequence, then $\vec{\nu}(\gamma_\tau) > f^{1^*}(\gamma_\tau)$,
4. $\vec{\nu}(\gamma_\tau) = s_\tau(\vec{\nu} \upharpoonright \text{dom}(f^{1^*}))$.

For every $\vec{\nu} \in A^{1^{**}}$, set $\langle f^{0\vec{\nu}}, A^{0\vec{\nu}} \rangle = \langle f^{0\vec{\nu} \upharpoonright \text{dom}(f^{1^*})}, A^{0\vec{\nu} \upharpoonright \text{dom}(f^{1^*})} \rangle$.

Consider the function $\vec{\nu} \mapsto \langle f^{0\vec{\nu}}, A^{0\vec{\nu}} \rangle, \vec{\nu} \in A^{1^{**}}$. Let $\langle f^{0^*}, A^{0^*} \rangle$ be represented by it in the ultrapower with $E(1)$.

It follows that $\langle \langle f^{0^*}, A^{0^*} \rangle, \langle f^{1^{**}}, A^{1^{**}} \rangle \rangle$ is a pure condition in $\mathcal{P}_{E(0), E(1)}$ which extends p .

The next claim completes the argument:

Claim 3 $\langle \langle f^{0^*}, A^{0^*} \rangle, \langle f^{1^{**}}, A^{1^{**}} \rangle \rangle \Vdash \sigma$.

Proof. Suppose otherwise. Then there is $\langle f, g \rangle \geq \langle \langle f^{0^*}, A^{0^*} \rangle, \langle f^{1^{**}}, A^{1^{**}} \rangle \rangle$ a non-pure in both coordinates condition which forces $\neg\sigma$. There is $\vec{\nu} \in A^{1^{**}} \upharpoonright \text{dom}(f^{1^*})$ such that $g \geq^* f_{\vec{\nu}}^{1^*}$. But then $f \geq t(\vec{\nu}, f^{1^*})$, and so, $\langle f, f_{\vec{\nu}}^{1^*} \rangle \Vdash \sigma$. Contradiction.

□ of the claim.

□

1.3 ω -many extenders.

So, we deal now with a sequence $\langle E(n) \mid n < \omega \rangle$ where each $E(n)$ is a (κ_n, λ) -extender and $\langle \kappa_n \mid n < \omega \rangle$ is an increasing sequence.

Let $\mathcal{P}_{E(i)}^*, \mathcal{P}_{E(i)}, i < \omega$ be as defined before.

Define first components of the main forcing $\mathcal{P}_{\langle E(n) \mid n < \omega \rangle}$.

Definition 1.18 The set of pure conditions $\mathcal{P}_{\langle E(n) \mid n < \omega \rangle}$ consists of all sequences $\langle p(n) \mid n < \omega \rangle$ such that for every $n < \omega$, the following hold:

1. $p(n) = \langle f^n, A^n \rangle \in \mathcal{P}_{E(n)}$,
2. $\text{dom}(f^n) \setminus \kappa_{n+1} \subseteq \text{dom}(f^{n+1})$,
3. for every $m \leq n$, for every $\alpha \in \text{dom}(f^m) \setminus \kappa_{n+1}$, if $f^{n+1}(\alpha)$ is not the empty sequence, then for every $\vec{\nu} \in A^{n+1}$, $\alpha \in \text{dom}(\vec{\nu})$ and $\vec{\nu}(\alpha) > f^{n+1}(\alpha)$.

The idea behind is as in the case of two extenders.

4. For every $\vec{\nu} \in A^{n+1}$ and $m \leq n$, the measures $E(m)(\text{dom}(f^m))$ and $E(m)((\text{dom}(f^m) \cap \kappa_{n+1}) \cup \{\vec{\nu}(\alpha) \mid \alpha \in \text{dom}(f^m) \setminus \kappa_{n+1}\})$ are basically the same in the following sense:

$$X \in E(m)(\text{dom}(f^m)) \text{ iff}$$

$$X^{ref} \in E(m)((\text{dom}(f^m) \cap \kappa_{n+1}) \cup \{\vec{\nu}(\alpha) \mid \alpha \in \text{dom}(f^m) \setminus \kappa_{n+1}\}),$$

where

$$X^{ref} = \{(\alpha, \beta) \in X \mid \alpha < \kappa_{n+1}\} \cup \{(\vec{\nu}(\alpha), \beta) \mid (\alpha, \beta) \in X, \alpha \geq \kappa_{n+1}\}.$$

Note that this property is true in the ultrapower by $E(n+1)$, so it holds on a set of measure one, as well.

Turn now to non-pure extensions. As usual, in Magidor type of iterations, non-pure extensions are allowed only at finitely many coordinates.

Start with a non-pure extension at a single coordinate and then proceed by induction.

We assume that for each $m < \omega$ there is a function $h_\lambda^m : \kappa_m \rightarrow \kappa_m$ such that $j_{E(m)}(h_\lambda^m)(\kappa_m) = \lambda$.

Definition 1.19 Let $m < \omega$. Define the set $\mathcal{P}_{\langle E(n) \mid n < \omega \rangle}^{\{m\}}$ of conditions with only non-pure part over the coordinate m . $\mathcal{P}_{\langle E(n) \mid n < \omega \rangle}^{(m)}$ consists of all sequences $\langle p(n) \mid n < \omega \rangle$ such that for every $n < \omega$, the following hold:

1. $\langle p(n) \mid n < \omega, n \neq m \rangle$ is a pure condition in $\mathcal{P}_{\langle E(n) \mid n < \omega, n \neq m \rangle}$,
2. $p(m) = f^m \in \mathcal{P}_{E(m)}^*$,
3. $\text{dom}(f^m) \setminus \kappa_n \subseteq \text{dom}(f^n)$, for every $n, m < n < \omega$,
4. for every $n, m < n < \omega$, for every $\alpha \in \text{dom}(f^m) \setminus \kappa_n$, if $f^n(\alpha)$ is not the empty sequence, then for every $\vec{\nu} \in A^n$, $\alpha \in \text{dom}(\vec{\nu})$ and $\vec{\nu}(\alpha) > f^n(\alpha)$,

5. for every $n, m < n < \omega$, for every $\gamma \in \text{dom}(f^m) \cap \kappa_n, \vec{v} \in A^n$ and $\alpha \in \text{dom}(\vec{v}), \vec{v}(\alpha) > \gamma$.
6. If $m > 0$, then the sequence $\langle p(n) \mid n < m \rangle$ will be a condition in the pure part of $\mathcal{P}_{\langle E(n) \mid h_\lambda(f^m(\kappa_m)) \mid n < m \rangle}$. The meaning is that if the value of the Prikry sequence for the normal measure of $E(m)$ is decided, then we cut all extenders $E(n), n < m$ to the reflection of λ below κ_m , i.e. to $h_\lambda^m(f^m(\kappa_m))$.

Let $m_1 < \dots < m_k < \omega, 1 \leq k < \omega$ and suppose that $\mathcal{P}_{\langle E(n) \mid n < \omega \rangle}^{\{m_1, \dots, m_k\}}$ the set of conditions with non-pure extensions over coordinates (m_1, \dots, m_k) only, is defined.

Let $m < \omega, m \notin \{m_1, \dots, m_k\}$.

Define non-pure extensions at the set of coordinates $\{m_1, \dots, m_k\} \cup \{m\}$.

Definition 1.20 Let $m < \omega$. Define the set $\mathcal{P}_{\langle E(n) \mid n < \omega \rangle}^{\{m_1, \dots, m_k\} \cup \{m\}}$ of conditions with only non-pure part over the coordinate m_1, \dots, m_k and m . $\mathcal{P}_{\langle E(n) \mid n < \omega \rangle}^{\{m_1, \dots, m_k\} \cup \{m\}}$ consists of all sequences $\langle p(n) \mid n < \omega \rangle$ such that for every $n < \omega$, the following hold:

1. $\langle p(n) \mid n < \omega, n \neq m \rangle$ is a condition in $\mathcal{P}_{\langle E(n) \mid n < \omega, n \neq m \rangle}^{\{m_1, \dots, m_k\}}$,
2. $p(m) = f^m \in \mathcal{P}_{E(m)}^*$.
3. If for every $i, 1 \leq i \leq k, m_i < m$, then following hold:
 - (a) $\text{dom}(f^m) \setminus \kappa_n \subseteq \text{dom}(f^n)$, for every $n, m < n < \omega$,
 - (b) for every $n, m < n < \omega$, for every $\alpha \in \text{dom}(f^m) \setminus \kappa_n$, if $f^n(\alpha)$ is not the empty sequence, then for every $\vec{v} \in A^n, \alpha \in \text{dom}(\vec{v})$ and $\vec{v}(\alpha) > f^n(\alpha)$,
 - (c) for every $n, m < n < \omega$, for every $\gamma \in \text{dom}(f^m) \cap \kappa_n, \vec{v} \in A^n$ and $\alpha \in \text{dom}(\vec{v}), \vec{v}(\alpha) > \gamma$.
 - (d) If $m > 0$, then the sequence $\langle p(n) \mid n < m \rangle$ will be a condition in $\mathcal{P}_{\langle E(n) \mid h_\lambda(f^m(\kappa_m)) \mid n < m \rangle}^{\{m_1, \dots, m_k\}}$.
The meaning is that if the value of the Prikry sequence for the normal measure of $E(m)$ is decided, then we cut all extenders $E(n), n < m$ to the reflection of λ below κ_m , i.e. to $h_\lambda^m(f^m(\kappa_m))$.

4. If there is $i, 1 \leq i \leq k, m_i > m$, then let i^* be the least such i . We require the following:

$$(a) \langle p(n) \mid n < m_{i^*} \rangle \in \mathcal{P}_{\langle E(n) \mid h_\lambda(f^{m_{i^*}}(\kappa_{m_{i^*}})) \mid n < m_{i^*} \rangle}^{\{m_1, \dots, m_{i^*-1}, m\}}$$

Finally set

$$\mathcal{P}_{\langle E(n)|n<\omega \rangle} = \bigcup \{ \mathcal{P}_{\langle E(n)|n<\omega \rangle}^{\{m_1, \dots, m_k\}} \mid k < \omega, m_1 < \dots < m_k < \omega \}.$$

Define the direct extension order \leq^* over $\mathcal{P}_{\langle E(n)|n<\omega \rangle}$ to be the union of such order over every $\mathcal{P}_{\langle E(n)|n<\omega \rangle}^{\{m_1, \dots, m_k\}}$, for every $k < \omega, m_1 < \dots < m_k < \omega$.

Turn now to the definition of the forcing order \leq over $\mathcal{P}_{\langle E(n)|n<\omega \rangle}$.

Let $m < \omega, m \notin \{m_1, \dots, m_k\}$. Define a one element extension at coordinate m of a condition in $\mathcal{P}_{\langle E(n)|n<\omega \rangle}^{\{m_1, \dots, m_k\}}$.

Definition 1.21 Let $p \in \mathcal{P}_{\langle E(n)|n<\omega \rangle}^{\{m_1, \dots, m_k\} \cup \{m\}}$ and $q \in \mathcal{P}_{\langle E(n)|n<\omega \rangle}^{\{m_1, \dots, m_k\}}$. Set $p \geq q$ iff the following hold:

1. Suppose that $m = 0$.

Then $p(0) = f^0 \in \mathcal{P}_{E(0)}^*$ and $q(0) = \langle g^0, B^0 \rangle$ is a pure condition in $\mathcal{P}_{E(0)}$.

Set $p \geq q$ iff $f^0 \geq \langle g^0, B^0 \rangle$ in $\mathcal{P}_{E(0)}$ and $\langle p(n) \mid 0 < n < \omega \rangle \geq^* \langle q(n) \mid 0 < n < \omega \rangle$ in $\mathcal{P}_{\langle E(n)|0 < n < \omega \rangle}$.

2. Suppose that $m > 0$.

Then $p(m) = f^m \in \mathcal{P}_{E(m)}^*$ and $q(m) = \langle g^m, B^m \rangle$ is a pure condition in $\mathcal{P}_{E(m)}$.

Set $p \geq q$ iff

(a) $f^m \geq \langle g^m, B^m \rangle$ in $\mathcal{P}_{E(m)}$ and $\langle p(n) \mid m < n < \omega \rangle \geq^* \langle q(n) \mid m < n < \omega \rangle$ in $\mathcal{P}_{\langle E(n)|m < n < \omega \rangle}$.

And

(b) $\langle p(n) \mid n < m \rangle \geq^* \langle q(n) \mid n < m \rangle^{ref}$ in $\mathcal{P}_{\langle E(n)|n < m \rangle}$, where $\langle q(n) \mid n < m \rangle^{ref}$ - the reflection of $\langle q(n) \mid n < m \rangle$ below κ_m is defined as follows, where $q(n) = \langle g^n, B^n \rangle$, if $n \notin \{m_1, \dots, m_k\}$ and $q(n) = \langle g^n \rangle$ otherwise.

i. Suppose first that $n \in \{m_1, \dots, m_k\}$.

Then

A. $\text{dom}((g^n)^{ref}) = (\text{dom}(g^n) \cap \kappa_m) \cup \{f^m(\alpha) \mid \alpha \in \text{dom}(g^n) \setminus \kappa_m\}$,

B. for every $\alpha \in \text{dom}(g^n) \cap \kappa_m = \text{dom}(g^n) \cap \text{dom}((g^n)^{ref})$, $(g^n)^{ref}(\alpha) = g^n(\alpha)$,

C. for every $\alpha \in \text{dom}(g^n) \setminus \kappa_m$, $(g^n)^{ref}(f^m(\alpha)) = g^n(\alpha)$.

It is crucial here that $f^m \upharpoonright (\text{dom}(g^n) \setminus \kappa_m)$ is one to one and the values there are above $\text{rng}(g^n) \cap \kappa_m$.

This follows by conditions (4),(5) of Definitions 1.10,1.11.

ii. Suppose now that $n \notin \{m_1, \dots, m_k\}$.

Then

- A. $\text{dom}((g^n)^{ref}) = (\text{dom}(g^n) \cap \kappa_m) \cup \{f^m(\alpha) \mid \alpha \in \text{dom}(g^n) \setminus \kappa_m\}$,
- B. for every $\alpha \in \text{dom}(g^n) \cap \kappa_m = \text{dom}(g^n) \cap \text{dom}((g^n)^{ref})$, $(g^n)^{ref}(\alpha) = g^n(\alpha)$,
- C. for every $\alpha \in \text{dom}(g^n) \setminus \kappa_m$, $(g^n)^{ref}(f^m(\alpha)) = g^n(\alpha)$.

Again, it is crucial here that $f^m \upharpoonright (\text{dom}(g^n) \setminus \kappa_m)$ is one to one and the values there are above $\text{rng}(g^n) \cap \kappa_m$, and this follows by conditions (3),(4) of Definition 1.18 and (4),(5) of Definition 1.19.

One more crucial observation here is that the measure $(E(n))(\text{dom}(g^n))$, to which B^n belongs, reflects to basically the same measure,

It follows by (4) of Definition 1.18.

- D. $A^n \upharpoonright \text{dom}((g^n)^{ref}) \subseteq \{(f^m(\alpha), \beta) \mid (\alpha, \beta) \in B^n\}$.

Denote further in this subsection $\mathcal{P}_{\langle E(n) \mid n < \omega \rangle}$ by just \mathcal{P} .

The next lemma follows from the definitions:

Lemma 1.22 *For every $m < \omega$, the forcing $\langle \mathcal{P}_{\langle E(n) \mid n < m \rangle}, \leq \rangle$ is equivalent to product of Cohen forcings $\text{Cohen}(\kappa_n^+, \eta_n)$'s, for some $\eta_n < \kappa_{n+1}$'s which depend on choice of a non-pure condition for $\mathcal{P}_{E(n+1)}$.*

Lemma 1.23 *For every $m < \omega$, the forcing $\langle \mathcal{P}_{\langle E(n) \mid m \leq n < \omega \rangle}, \leq^* \rangle$ is κ_m -closed.*

Lemma 1.24 *The forcing $\langle \mathcal{P}, \leq \rangle$ satisfies κ_ω^{++} -c.c.*

Proof. Use the standard Δ -system argument.

□

Lemma 1.25 *$\langle \mathcal{P}, \leq, \leq^* \rangle$ is a Prikry type forcing notion.*

Proof. The proof is similar to those of Lemmas 1.9, 1.17 (and in turn to those of Merimovich [7]).

Assume that for every $m < \omega$, $\langle \mathcal{P}_{\langle E(n) \mid n < m \rangle}, \leq, \leq^* \rangle$ is a Prikry type forcing notion.

Suppose that $\langle \mathcal{P}, \leq, \leq^* \rangle$ does not have the Prikry property.

Let $p \in \mathcal{P}$ be a pure condition and σ a statement of the forcing language which is undecided by pure extensions of p . Then p is of the form $\langle \langle f^m, A^m \rangle \mid n < \omega \rangle$.

Proceed by induction on $m < \omega$ and define an \leq^* -increasing sequence $\langle p_m \mid m < \omega \rangle$ of direct extensions of p .

Assume that for every $n < m$, p_n is defined. Define p_m . If $m = 0$, then let p_{-1} be p . At stage m we deal with the coordinate m of the condition.

Construct by induction an increasing chain of elementary submodels $\langle N_\xi^m \mid \xi < \kappa_m \rangle$ of H_χ , for χ large enough, and a sequence $\langle f_\xi \mid \xi < \kappa \rangle$ of members of $\mathcal{P}_{E(m)}^*$, such that

1. $p, p_{m-1}, \mathcal{P}, \sigma \in N_0^m$,
2. $N_0^m \supseteq \kappa_m$,
3. for every $\xi < \kappa_m$,
 - (a) $|N_\xi^m| = \kappa_m$,
 - (b) $\kappa_m > N_\xi^m \subseteq N_\xi^m$,
 - (c) $\langle f_\zeta^m \mid \zeta < \xi \rangle \in N_\xi^m$,
 - (d) $f_\xi^m \in \bigcap \{D' \in N_\xi^m \mid D' \text{ is a dense open subset of } \mathcal{P}_{E(m)}^* \text{ above } f^{mp}\}$,
 - (e) $f^{mp} \leq^* f_0^m$,
 - (f) $f_\xi^m \geq^* f_\zeta^m$, for every $\zeta < \xi$.

Set $N^m = \bigcup_{\xi < \kappa_1} N_\xi^m$ and $f^{m*} = \bigcup \{f_\xi^m \mid \xi < \kappa\}$. Let $A \subseteq [\text{dom}(f^{m*}) \times \kappa_m]^{<\kappa_m}$ be such that

- $A \upharpoonright \text{dom}(f^{pm}) \subseteq A^{pm}$,
- $A \in (E(m))(\text{dom}(f^{m*}))$.

Note that $A \subseteq N^m$, since $\text{dom}(f^{m*}) \subseteq N^m$, and so, $[\text{dom}(f^{m*}) \times \kappa_m]^{<\kappa_m} \subseteq N^m$.

Let $\vec{\nu} \in A$. Consider $\lambda_m^{\vec{\nu}} := h_\lambda^m(\vec{\nu}(\kappa_m))$, i.e. the cardinal below κ_m that now corresponds to λ . Suppose for simplicity that $\text{dom}(f^{pn}) \subseteq \lambda_m^{\vec{\nu}}$, for every $n < m$, otherwise just reflect the part above κ_m below as in Definition 1.21.

Consider $\mathcal{P}_{\langle E(n) \upharpoonright \lambda_m^{\vec{\nu}} \mid n < m \rangle}$. Clearly, it is contained and belongs to N^m .

Let $\langle t_\xi \mid \xi < \lambda_m^{\vec{\nu}} \rangle$ be an enumeration of this forcing notion in N^m .

Let $f \in \mathcal{P}_{E(m)}^*$, $f \geq^* f^{pm}$.

Proceed by induction on $\xi < \lambda_m^{\vec{\nu}}$. Define an \leq^* -increasing sequence $\langle f_\xi \mid \xi < \lambda_m^{\vec{\nu}} \rangle$ of direct extensions of f and an \leq^* -increasing sequence $\langle p_\xi^{>m} \mid \xi < \lambda_m^{\vec{\nu}} \rangle$ of direct extensions of $\langle p_m(n) \mid m < n < \omega \rangle$

such that, for every $\xi < \lambda_m^{\vec{\nu}}$, either

(1) $\langle t_\xi, (f_\xi)_{\vec{v}}, p_\xi^{>m} \rangle \parallel \sigma$,

or

(2) for every $q \geq^* \langle (f_\xi)_{\vec{v}}, p_\xi^{>m} \rangle$, $\langle t_\xi, q \rangle \not\parallel \sigma$.

Let $\bar{f} = \bigcup_{\xi < \lambda_1^{\vec{v}}} f_\xi$ and $\bar{p}^{>m} = \bigcup_{\xi < \lambda_1^{\vec{v}}} p_\xi^{>m}$.

Then, for every $t \in \mathcal{P}_{E(0) \upharpoonright \lambda_1^{\vec{v}}}$ either

(1) $\langle t, \bar{f}_{\vec{v}}, \bar{p}^{>m} \rangle \parallel \sigma$,

or

(2) for every $q \geq^* \langle \bar{f}_{\vec{v}}, \bar{p}^{>m} \rangle$, $\langle t, q \rangle \not\parallel \sigma$.

Consider now the following statement of the forcing language of $\mathcal{P}_{\langle E(n) \upharpoonright \lambda_m^{\vec{v}} \mid n < m \rangle}$:

$$\varphi \equiv \exists t \in \mathcal{G}(\langle t, \bar{f}_{\vec{v}}, \bar{p}^{>m} \rangle \parallel \sigma).$$

By the Prikry condition of the forcing $\mathcal{P}_{\langle E(n) \upharpoonright \lambda_m^{\vec{v}} \mid n < m \rangle}$, there is $t^* \geq^* \langle p_{m-1}(n) \mid n < m \rangle$ which decides φ .

If $t^* \Vdash \neg \varphi$, then set $t(\vec{v}, f) = t^*$.

If $t^* \Vdash \varphi$. Use again the Prikry condition of the forcing $\mathcal{P}_{\langle E(n) \upharpoonright \lambda_m^{\vec{v}} \mid n < m \rangle}$ to decide the following statement

$$\psi \equiv \exists t \in \mathcal{G}(\langle t, \bar{f}_{\vec{v}}, \bar{p}^{>m} \rangle \Vdash \sigma).$$

Let $t(\vec{v}, f) \geq^* t^*$ be a condition which decides ψ .

Claim 4 Let $t \geq t(\vec{v}, f)$ in $\mathcal{P}_{\langle E(n) \upharpoonright \lambda_m^{\vec{v}} \mid n < m \rangle}$, $\langle g, q \rangle \geq^* \langle \bar{f}_{\vec{v}}, \bar{p}^{>m} \rangle$ in $\mathcal{P}_{\langle E(n) \mid m \leq n < \omega \rangle}$.

Suppose that $\langle t, g, q \rangle \Vdash \sigma$ (or $\langle t, g, q \rangle \Vdash \neg \sigma$),

then already $\langle t(\vec{v}, f), \bar{f}_{\vec{v}}, \bar{p}^{>m} \rangle \Vdash \sigma$ (or $\langle t(\vec{v}, f), \bar{f}_{\vec{v}}, \bar{p}^{>m} \rangle \Vdash \neg \sigma$).

Proof. Let $t \geq t(\vec{v}, f)$ in $\mathcal{P}_{\langle E(n) \upharpoonright \lambda_m^{\vec{v}} \mid n < m \rangle}$, $\langle g, q \rangle \geq^* \langle \bar{f}_{\vec{v}}, \bar{p}^{>m} \rangle$ in $\mathcal{P}_{\langle E(n) \mid m \leq n < \omega \rangle}$.

Suppose that $\langle t, g, q \rangle \Vdash \sigma$.

Then, for some $\xi < \lambda_1^{\vec{v}}$, $t = t_\xi$, and then, $\langle t, (f_\xi)_{\vec{v}}, p_\xi^{>m} \rangle \parallel \sigma$. So, $\langle t, \bar{f}_{\vec{v}}, \bar{p}^{>m} \rangle \parallel \sigma$.

Then $t^* \Vdash \varphi$. Hence, $\langle t(\vec{v}, f), \bar{f}_{\vec{v}}, \bar{p}^{>m} \rangle \Vdash \sigma$.

□ of the claim.

Define $D_{\vec{v}}$ to be the set of all $f \in \mathcal{P}_{E(m)}^*$, $f \geq^* f^{p_{m-1}^m}$ so that there is $p_f^{>m} \in \mathcal{P}_{\langle E(n) \mid m < n < \omega \rangle}$, $p_f^{>m} \geq^* p_{m-1}^{>m}$, such that either

(1) $\langle t(\vec{v}, f), f_{\vec{v}}, p_f^{>m} \rangle \parallel \sigma$

or

(2) for every $t \geq t(\vec{v}, f)$ in $\mathcal{P}_{\langle E(n) \mid \lambda_m^{\vec{v}} \mid n < m \rangle}$, for every $\langle g, q \rangle \geq^* \langle f_{\vec{v}}, p_f^{>m} \rangle$ in $\mathcal{P}_{\langle E(n) \mid m \leq n < \omega \rangle}$, $\langle t, g, q \rangle \not\parallel \sigma$.

The next claim follows now from the previous one:

Claim 5 $D_{\vec{v}}$ is a dense open subset of $\mathcal{P}_{E(m)}^*$ above $f^{p_{m-1}m}$.

$D_{\vec{v}}$ is definable with parameters in N^m , hence $D_{\vec{v}} \in N^m$.

Then, $f^{m*} \in D_{\vec{v}}$, for every $\vec{v} \in A$.

So, for every $\vec{v} \in A$ we have either

(3) $\langle t(\vec{v}, f^{m*}), f_{\vec{v}}^{m*}, p_{f^{m*}}^{>m} \rangle \parallel \sigma$

or

(4) for every $t \geq t(\vec{v}, f^{m*})$ in $\mathcal{P}_{\langle E(n) \mid \lambda_m^{\vec{v}} \mid n < m \rangle}$, for every $\langle g, q \rangle \geq^* \langle f_{\vec{v}}^{m*}, p_{f^{m*}}^{>m} \rangle$ in $\mathcal{P}_{\langle E(n) \mid m \leq n < \omega \rangle}$, $\langle t, g, q \rangle \not\parallel \sigma$.

Shrink A , if necessary, to a set $A^{m*} \in (E(m))(\text{dom}(f^{m*}))$, such that for any two $\vec{v}, \vec{v}' \in A^{m*}$ the decision is the same.

Consider now $\langle f^{m*}, A^{m*} \rangle$ it is a pure condition in $\mathcal{P}_{E(m)}$. Use the function $\vec{v} \mapsto t(\vec{v}, f^{m*})$ in order to get a pure condition in $\mathcal{P}_{\langle E(n) \mid n < m \rangle}$, just use the one this function represents in the ultrapower by $(E(m))(\text{dom}(f^{m*}))$. Denote it by $\langle \langle f^{n*}, A^{n*} \rangle \mid n < m \rangle$.

Let us explain how do we naturally combine the result into a condition in $\mathcal{P}_{\langle E(n) \mid n < \omega \rangle}$.

Let $t(\vec{v}, f^{m*}) = \langle \langle f^{n\vec{v}}, A^{n\vec{v}} \rangle \mid n < m \rangle$, for every $\vec{v} \in A^{m*}$. Consider $f^{n\vec{v}}, n < m$. It is a set of at most κ_n many pairs (α, β) , where $\alpha < \lambda_m^{\vec{v}} < \kappa_m$ and β is either the empty sequence or an ordinal $< \kappa_n$.

Shrinking A^{m*} if necessary, we can assume that there are $\langle x_n \mid n < m \rangle$ and $\kappa_n^* < \kappa_n^+, n < m$ such that for every $\vec{v}, \vec{v}' \in A^{m*}$, for every $n < m$, the following hold:

1. $\text{dom}(f^{n\vec{v}}) \cap \vec{v}(\kappa_m) = x_n$,
2. $\text{dom}(f^{n\vec{v}}) \setminus \vec{v}(\kappa_m) = \{\gamma_{\tau n}^{\vec{v}} \mid \tau < \kappa_n^*\}$ is an increasing enumeration,
3. for every $\alpha \in x_n$, $f^{n\vec{v}}(\alpha) = f^{n\vec{v}'}(\alpha)$,
4. for every $\tau < \kappa_n^*$, $f^{n\vec{v}}(\gamma_{\tau n}^{\vec{v}}) = f^{n\vec{v}'}(\gamma_{\tau n}^{\vec{v}'})$

Consider, for every $n < m$ and $\tau < \kappa_n^*$ a function $s_{\tau n}$ on A^{m*} defined by setting $s_{\tau n}(\vec{\nu}) = \gamma_{\tau n}^{\vec{\nu}}$.
Let

$$\gamma_{\tau n} = j_{E(m)}(s_{\tau n})(\langle (j_{E(m)}(\alpha), \alpha) \mid \alpha \in \text{dom}(f^{m*}) \rangle).$$

Extend now f^{m*} to f^{m**} by adding all $\gamma_{\tau n}, \tau < \kappa_n^*, n < m$ to its domain and setting $f^{m**}(\gamma_{\tau n})$ to be the empty sequence whenever $\gamma_{\tau n} \notin \text{dom}(f^{m*})$.

Define $A^{m**} \in E(m)(\text{dom}(f^{m**}))$ as follows.

Set $\vec{\nu} \in A^{m**}$ iff

1. $\vec{\nu} \upharpoonright \text{dom}(f^{m*}) \in A^{m*}$,
2. $\text{dom}(\vec{\nu}) \supseteq \{\gamma_{\tau n} \mid \tau < \kappa_n^*, n < m\}$,
3. if $\gamma_{\tau n} \in \text{dom}(f^{m*})$ and $f^{m*}(\gamma_{\tau n})$ is not the empty sequence, then $\vec{\nu}(\gamma_{\tau n}) > f^{m*}(\gamma_{\tau n})$, for every $n < m$,
4. $\vec{\nu}(\gamma_{\tau n}) = s_{\tau n}(\vec{\nu} \upharpoonright \text{dom}(f^{m*}))$, for every $n < m$.

For every $\vec{\nu} \in A^{m**}, n < m$, set $\langle f^{n\vec{\nu}}, A^{n\vec{\nu}} \rangle = \langle f^{n\vec{\nu} \upharpoonright \text{dom}(f^{m*})}, A^{n\vec{\nu} \upharpoonright \text{dom}(f^{m*})} \rangle$.

Consider the function $\vec{\nu} \mapsto \langle \langle f^{n\vec{\nu}}, A^{n\vec{\nu}} \rangle \mid n < m \rangle, \vec{\nu} \in A^{m**}$. Let $\langle \langle f^{n*}, A^{n*} \rangle \mid n < m \rangle$ be represented by it in the ultrapower with $E(m)$.

It follows that $\langle \langle \langle f^{n*}, A^{n*} \rangle \mid n < m \rangle, \langle f^{m**}, A^{m**} \rangle \rangle$ is a pure condition in $\mathcal{P}_{\langle E(n) \mid n \leq m \rangle}$ which extends $p_{m-1} \upharpoonright \mathcal{P}_{\langle E(n) \mid n \leq m \rangle}$.

Extend purely $p_{f^{m*}}^{>m}$ in the obvious fashion to a condition $p_{f^{m**}}^{>m}$ in $\mathcal{P}_{\langle E(n) \mid m < n < \omega \rangle}$ such that $\langle \langle \langle f^{n*}, A^{n*} \rangle \mid n < m \rangle, \langle f^{m**}, A^{m**} \rangle, p_{f^{m**}}^{>m} \rangle$ is a pure condition in $\mathcal{P}_{\langle E(n) \mid n < \omega \rangle}$. Then it extends p_{m-1} .

Set p_m to be $\langle \langle \langle f^{n*}, A^{n*} \rangle \mid n < m \rangle, \langle f^{m**}, A^{m**} \rangle, p_{f^{m**}}^{>m} \rangle$.

This completes the recursive construction of $\langle p_m \mid m < \omega \rangle$. Let $p_* \geq p_m$, for every $m < \omega$.

The next claim completes the argument:

Claim 6 $p_* \parallel \sigma$.

Proof. Suppose otherwise. Pick then $q \geq p_*$ to be a condition which decides σ and such that its last coordinate at which a non-direct extension was made is as small as possible.

Let $q \Vdash \sigma$ and this coordinate is some $m < \omega$.

Then there is $\vec{\nu} \in A^{p_*(m)}$ such that $q(m) \geq^* f^{p_*(m)}_{\vec{\nu}}$ in $\mathcal{P}_{E(m)}^*$. In addition, $q^{>m} \geq^* p_*^{>m}$ in $\mathcal{P}_{\langle E(n) \mid m < n < \omega \rangle}$, by the choice of m .

But, then the condition (4) above cannot hold. Hence (3) is true, which means, that

$$\langle t(\vec{v}, f^{m*}), f_{\vec{v}}^{m*}, p_{f^{m*}}^{>m} \rangle \Vdash \sigma.$$

Then the same holds for every $\vec{v}' \in A^{p^*}(m)$. So, already $p_* \Vdash \sigma$.

Contradiction.

□ of the claim.

□

It follows now that the forcing $\langle \mathcal{P}, \leq \rangle$ preserves all the cardinals but κ_ω^+ . Using the arguments of the previous lemma it is possible to show that κ_ω^+ is preserved as well.

Let G be a generic subset of $\langle \mathcal{P}, \leq \rangle$.

Lemma 1.26 κ_ω remains a strong limit cardinal in $V[G]$.

Proof. Given $p \in \mathcal{P}$ and $m < \omega$. Suppose that $p(m)$ is non-pure. Then $p(m)(\kappa_m)$ is defined, and hence also the reflection $h_\lambda^m(p(m)(\kappa_m))$ of λ below κ_m . By the definition of the forcing, then the part $\mathcal{P}_{\langle E(n) \mid n < m \rangle}$ above p will act as $\mathcal{P}_{\langle E(n) \mid h_\lambda^m(p(m)(\kappa_m)) \mid n < m \rangle}$. In particular, $2^{\kappa_n} \leq h_\lambda^m(p(m)(\kappa_m)) < \kappa_m$. The upper part of the forcing, i.e. $\mathcal{P}_{\langle E(n) \mid m \leq n < \omega \rangle}$, does not add new bounded subsets to κ_m .

So we are done.

□

Lemma 1.27 $(\kappa_\omega^+)^V$ remains a cardinal in $V[G]$.

Let us state first the following:

Lemma 1.28 Let $p \in \mathcal{P}$ and ζ be a $\langle \mathcal{P}, \leq \rangle$ -name of an ordinal or just $p \Vdash_{\langle \mathcal{P}, \leq \rangle} \zeta$ is an ordinal.

Then there are $p^* \geq^* p$ and $n_1 < \dots < n_k$, for some $k < \omega$, such that

1. for every $i, 1 \leq i \leq k$, $p^*(n_i) = \langle f_{n_i}^{p^*}, A_{n_i}^{p^*} \rangle$,
2. for every $\vec{v}_1 \in A_{n_1}^{p^*}, \dots, \vec{v}_k \in A_{n_k}^{p^*}$,
 $p^* \frown \vec{v}_1 \dots \frown \vec{v}_k$ decides ζ .

The proof of this lemma repeats the proof of the Prikry condition of the forcing.

Proof of 1.27. Suppose otherwise. Then there is $\mu < \kappa_\omega$ such that, in $V[G]$, $\text{cof}((\kappa_\omega^+)^V) = \mu$. Back in V , let $\langle \zeta_\tau \mid \tau < \mu \rangle$ be a name of a witnessing sequence.

Pick $\bar{n} < \omega$ with $\kappa_{\bar{n}} > \mu$. Let $p \in \mathcal{P}$ be such that $p(\bar{n}) \in \mathcal{P}_{E(\bar{n})}^*$, i.e. its \bar{n} -th coordinate is non-pure. Then above p the part $\mathcal{P}_{E(n)|n < \bar{n}}$ reflects down to $\mathcal{P}_{\langle E(n) \upharpoonright h_{\bar{n}}^\lambda(p(\bar{n})) \upharpoonright n < \bar{n} \rangle}$, and so has cardinality below $\kappa_{\bar{n}}$.

Construct a sequence $\langle p_\tau \mid \tau < \mu \rangle$ of \leq^* -extensions of p such that, for every $\tau < \mu$,

1. p_τ satisfies the conclusion of Lemma 1.28 for ζ_τ ,
2. $\langle p_\tau(n) \mid \bar{n} \leq n < \omega \rangle \leq^* \langle p_{\tau'}(n) \mid \bar{n} \leq n < \omega \rangle$ in the forcing $\mathcal{P}_{\langle E(n) \upharpoonright \bar{n} \leq n < \omega \rangle}$, for every $\tau < \tau' < \mu$.

Let $s \geq^* \langle p_\tau(n) \mid \bar{n} \leq n < \omega \rangle$ in the forcing $\mathcal{P}_{\langle E(n) \upharpoonright \bar{n} \leq n < \omega \rangle}$, for every $\tau < \mu$. Set $r = p \upharpoonright \bar{n} \frown s$. Then, for every $\tau < \mu$, there is $\xi_\tau < \kappa_\omega^+$ such that

$$r \Vdash_{\langle \mathcal{P}, \leq \rangle} \zeta_\tau < \xi_\tau,$$

since by the choice of p_τ , the number of possibilities for ζ_τ has cardinality $< \kappa_\omega$.

Set $\xi = \bigcup_{\tau < \mu} \xi_\tau < \kappa_\omega^+$.

$$r \Vdash_{\langle \mathcal{P}, \leq \rangle} \langle \zeta_\tau \mid \tau < \mu \rangle \text{ is bounded by } \xi.$$

Contradiction.

□

Given $p \in \mathcal{P}$. Denote by $\text{np}(p)$ the set of all coordinates n of p such that $p(n) \in \mathcal{P}_{E(n)}^*$, i.e. a non-pure extension was made at the coordinate n .

Let $\beta \in [\kappa_\omega, \lambda)$ we define in $V[G]$ a function $t_\beta : \omega \rightarrow \kappa_\omega$ as follows.

For every $n < \omega$, find $p \in G$ such that $n \in \text{np}(p)$ and if $n_1 < \dots < n_k$ is the increasing enumeration of $\text{np}(p) \setminus n$ (i.e. $n = n_1$), then the following hold:

1. $\beta \in \text{dom}(p(n_k))$.
Set $\beta_k = \beta$.
2. For every $i, 1 \leq i \leq k - 1$, $\beta_i \in \text{dom}(p(n_i))$,
where $\beta_i = p(n_{i+1})(\beta_{i+1})$.

Set $t_\beta(n) = p(n)(\beta_1)$.

Lemma 1.29 *In $V[G]$, if $\beta, \gamma \in [\kappa_\omega, \lambda)$ and $\beta < \gamma$, then there is $n^* < \omega$ such that for every $n, n^* \leq n < \omega$, $t_\beta(n) < t_\gamma(n)$.*

Proof. Work in V . Let $p \in \mathcal{P}$ be any condition and $\beta, \gamma \in [\kappa_\omega, \lambda), \beta < \gamma$.

Let n^* be a coordinate above $\text{np}(p)$. Then $p(n) = \langle f_n^p, A_n^p \rangle$, for every $n, n^* \leq n < \omega$.

Extend p to p^* by adding β, γ to all $\text{dom}(f_n^p)$ with $n^* \leq n < \omega$.

Now, by the definition of the order on \mathcal{P} , for every $n, n^* \leq n < \omega$ and every $q \geq p^*$ such that q defines $t_\beta(n)$ and $t_\gamma(n)$, we will have $t_\beta(n) < t_\gamma(n)$.

So,

$$p^* \Vdash (\forall n)(n^* \leq n < \omega \rightarrow \underset{\sim}{t}_\beta(n) < \underset{\sim}{t}_\gamma(n)).$$

□

It is possible to say a bit more. Namely, let in $V[G]$, for every $n < \omega$, λ_n be the reflection of λ below κ_n , i.e. for some $p \in G$ with $p(n) = f_n^p$, $\lambda_n = h_\lambda^n(f_n^p(\kappa_n))$. Then the following holds:

Lemma 1.30 *The sequence $\langle t_\beta \mid \beta \in [\kappa_\omega, \lambda) \rangle$ is a scale in $\langle \prod_{n < \omega} \lambda_n, <_{J^{bd}} \rangle$.*

2 Arbitrary cofinality.

Let η be any ordinal. We generalize the construction of the previous section to sequences of extenders of the length η . The generalization is straightforward. Let us repeat just the main points.

So, we deal now with a sequence $\langle E(\alpha) \mid \alpha < \eta \rangle$, where each $E(\alpha)$ is a (κ_α, λ) -extender and $\langle \kappa_\alpha \mid n < \omega \rangle$ is an increasing sequence with $\eta < \kappa_0$.

Let $\mathcal{P}_{E(i)}^*, \mathcal{P}_{E(i)}, i < \eta$ be as defined before.

Define components of the main forcing $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}$.

Definition 2.1 The set of pure conditions $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}$ consists of all sequences $\langle p(\alpha) \mid \alpha < \eta \rangle$ such that for every $\alpha < \eta$, the following hold:

1. $p(\alpha) = \langle f^\alpha, A^\alpha \rangle \in \mathcal{P}_{E(\alpha)}$,
2. $\text{dom}(f^\alpha) \setminus \kappa_{\alpha+1} \subseteq \text{dom}(f^{\alpha+1})$,
3. for every $\beta \leq \alpha$, for every $\xi \in \text{dom}(f^\beta) \setminus \kappa_{\alpha+1}$, if $f^{\alpha+1}(\xi)$ is not the empty sequence, then for every $\vec{v} \in A^{\alpha+1}$, $\xi \in \text{dom}(\vec{v})$ and $\vec{v}(\xi) > f^{\alpha+1}(\xi)$.

The idea behind is as in the case of two extenders.

4. For every $\vec{v} \in A^{\alpha+1}$ and $\beta \leq \alpha$, the measures $E(\beta)(\text{dom}(f^\beta))$ and $E(\beta)((\text{dom}(f^\beta) \cap \kappa_{\alpha+1}) \cup \{\vec{v}(\xi) \mid \xi \in \text{dom}(f^\beta) \setminus \kappa_{\alpha+1}\})$ are basically the same in the following sense:

$$X \in E(\beta)(\text{dom}(f^\beta)) \text{ iff}$$

$$X^{ref} \in E(\beta)((\text{dom}(f^\beta) \cap \kappa_{\alpha+1}) \cup \{\vec{v}(\xi) \mid \xi \in \text{dom}(f^\beta) \setminus \kappa_{\alpha+1}\}),$$

where

$$X^{ref} = \{(\xi, \beta) \in X \mid \xi < \kappa_{\alpha+1}\} \cup \{(\vec{v}(\xi), \beta) \mid (\xi, \beta) \in X, \xi \geq \kappa_{\alpha+1}\}.$$

Note that this property is true in the ultrapower by $E(\alpha + 1)$, so it holds on a set of measure one, as well.

Turn now to non-pure extensions. As usual, in Magidor type of iterations, non-pure extensions are allowed only at finitely many coordinates.

Start with a non-pure extension at a single coordinate and then proceed by induction.

We assume that for each $\alpha < \eta$ there is a function $h_\lambda^\alpha : \kappa_\alpha \rightarrow \kappa_\alpha$ such that $j_{E(\alpha)}(h_\lambda^\alpha)(\kappa_\alpha) = \lambda$.

Definition 2.2 Let $\beta < \eta$. Define the set $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}^{\{\beta\}}$ of conditions with only non-pure part over the coordinate β . $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}^{(\beta)}$ consists of all sequences $\langle p(\alpha) \mid \alpha < \eta \rangle$ such that for every $\alpha < \eta$, the following hold:

1. $\langle p(\alpha) \mid \alpha < \eta, \alpha \neq \beta \rangle$ is a pure condition in $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta, \alpha \neq \beta \rangle}$,
2. $p(\beta) = f^\beta \in \mathcal{P}_{E(\beta)}^*$,
3. $\text{dom}(f^\beta) \setminus \kappa_\alpha \subseteq \text{dom}(f^\alpha)$, for every $\alpha, \beta < \alpha < \eta$,
4. for every $\alpha, \beta < \alpha < \eta$, for every $\xi \in \text{dom}(f^\beta) \setminus \kappa_\alpha$, if $f^\alpha(\xi)$ is not the empty sequence, then for every $\vec{v} \in A^\alpha$, $\xi \in \text{dom}(\vec{v})$ and $\vec{v}(\xi) > f^\alpha(\xi)$,
5. for every $\alpha, \beta < \alpha < \eta$, for every $\gamma \in \text{dom}(f^\beta) \cap \kappa_\alpha$, $\vec{v} \in A^\alpha$ and $\xi \in \text{dom}(\vec{v})$, $\vec{v}(\xi) > \gamma$.
6. If $\beta > 0$, then the sequence $\langle p(\alpha) \mid \alpha < \beta \rangle$ will be a condition in the pure part of $\mathcal{P}_{\langle E(\alpha) \mid h_\lambda(f^\beta(\kappa_\beta)) \mid \alpha < \beta \rangle}$. The meaning is that if the value of the Prikry sequence for the normal measure of $E(\beta)$ is decided, then we cut all extenders $E(\alpha), \alpha < \beta$ to the reflection of λ below κ_β , i.e. to $h_\lambda^\beta(f^\beta(\kappa_\beta))$.

Let $\beta_1 < \dots < \beta_k < \eta, 1 \leq k < \omega$ and suppose that $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}^{\{\beta_1, \dots, \beta_k\}}$ the set of conditions with non-pure extensions over coordinates $(\beta_1, \dots, \beta_k)$ only, is defined.

Let $\beta < \eta, \beta \notin \{\beta_1, \dots, \beta_k\}$.

Define non-pure extensions at the set of coordinates $\{\beta_1, \dots, \beta_k\} \cup \{\beta\}$.

Definition 2.3 Let $\beta < \eta$. Define the set $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}^{\{\beta_1, \dots, \beta_k\} \cup \{\beta\}}$ of conditions with only non-pure part over the coordinate β_1, \dots, β_k and β . $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}^{\{\beta_1, \dots, \beta_k\} \cup \{\beta\}}$ consists of all sequences $\langle p(\alpha) \mid \alpha < \eta \rangle$ such that for every $\alpha < \eta$, the following hold:

1. $\langle p(\alpha) \mid \alpha < \eta, \alpha \neq \beta \rangle$ is a condition in $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta, \alpha \neq \beta \rangle}^{\{\beta_1, \dots, \beta_k\}}$,
2. $p(\beta) = f^\beta \in \mathcal{P}_{E(\beta)}^*$.
3. If for every $i, 1 \leq i \leq k, \beta_i < \beta$, then following hold:
 - (a) $\text{dom}(f^\beta) \setminus \kappa_\alpha \subseteq \text{dom}(f^\alpha)$, for every $\alpha, \beta < \alpha < \eta$,
 - (b) for every $\alpha, \beta < \alpha < \eta$, for every $\xi \in \text{dom}(f^\beta) \setminus \kappa_\alpha$, if $f^\alpha(\xi)$ is not the empty sequence, then for every $\vec{v} \in A^\alpha$, $\xi \in \text{dom}(\vec{v})$ and $\vec{v}(\xi) > f^\alpha(\xi)$,

(c) for every $\alpha, \beta < \alpha < \eta$, for every $\gamma \in \text{dom}(f^\beta) \cap \kappa_\alpha, \vec{v} \in A^\alpha$ and $\xi \in \text{dom}(\vec{v})$, $\vec{v}(\xi) > \gamma$.

(d) If $\beta > 0$, then the sequence $\langle p(\alpha) \mid \alpha < \beta \rangle$ will be a condition

in $\mathcal{P}_{\langle E(\alpha) \upharpoonright h_\lambda(f^\beta(\kappa_\beta)) \mid \alpha < \beta \rangle}^{\{\beta_1, \dots, \beta_k\}}$.

The meaning is that if the value of the Prikry sequence for the normal measure of $E(\beta)$ is decided, then we cut all extenders $E(\alpha), \alpha < \beta$ to the reflection of λ below κ_β , i.e. to $h_\lambda^\beta(f^\beta(\kappa_\beta))$.

4. If there is $i, 1 \leq i \leq k, \beta_i > \beta$, then let i^* be the least such i . We require the following:

(a) $\langle p(\alpha) \mid \alpha < \beta_{i^*} \rangle \in \mathcal{P}_{\langle E(\alpha) \upharpoonright h_\lambda(f^{\beta_{i^*}}(\kappa_{\beta_{i^*}})) \mid \alpha < \beta_{i^*} \rangle}^{\{\beta_1, \dots, \beta_{i^*-1}, \beta\}}$.

Finally set

$$\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle} = \bigcup \{ \mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}^{\{\beta_1, \dots, \beta_k\}} \mid k < \omega, \beta_1 < \dots < \beta_k < \omega \}.$$

Define the direct extension order \leq^* over $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}$ to be the union of such order over every $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}^{\{\beta_1, \dots, \beta_k\}}$, for every $k < \omega, \beta_1 < \dots < \beta_k < \eta$.

Turn now to the definition of the forcing order \leq over $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}$.

Let $\beta < \eta, \beta \notin \{\beta_1, \dots, \beta_k\}$. Define a one element extension at coordinate β of a condition in $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}^{\{\beta_1, \dots, \beta_k\}}$.

Definition 2.4 Let $p \in \mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}^{\{\beta_1, \dots, \beta_k\} \cup \{\beta\}}$ and $q \in \mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}^{\{\beta_1, \dots, \beta_k\}}$. Set $p \geq q$ iff the following hold:

1. Suppose that $\beta = 0$.

Then $p(0) = f^0 \in \mathcal{P}_{E(0)}^*$ and $q(0) = \langle g^0, B^0 \rangle$ is a pure condition in $\mathcal{P}_{E(0)}$.

Set $p \geq q$ iff $f^0 \geq \langle g^0, B^0 \rangle$ in $\mathcal{P}_{E(0)}$ and $\langle p(\alpha) \mid 0 < \alpha < \eta \rangle \geq^* \langle q(\alpha) \mid 0 < \alpha < \eta \rangle$ in $\mathcal{P}_{\langle E(\alpha) \mid 0 < \alpha < \eta \rangle}$.

2. Suppose that $\beta > 0$.

Then $p(\beta) = f^\beta \in \mathcal{P}_{E(\beta)}^*$ and $q(\beta) = \langle g^\beta, B^\beta \rangle$ is a pure condition in $\mathcal{P}_{E(\beta)}$.

Set $p \geq q$ iff

(a) $f^\beta \geq \langle g^\beta, B^\beta \rangle$ in $\mathcal{P}_{E(\beta)}$ and $\langle p(\alpha) \mid \beta < \alpha < \eta \rangle \geq^* \langle q(\alpha) \mid \beta < \alpha < \eta \rangle$ in $\mathcal{P}_{\langle E(\alpha) \mid \beta < \alpha < \eta \rangle}$.

And

(b) $\langle p(\alpha) \mid \alpha < \beta \rangle \geq^* \langle q(\alpha) \mid \alpha < \beta \rangle^{ref}$ in $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \beta \rangle}$, where $\langle q(\alpha) \mid \alpha < \beta \rangle^{ref}$ - the reflection of $\langle q(\alpha) \mid \alpha < \beta \rangle$ below κ_β is defined as follows, where $q(\alpha) = \langle g^\alpha, B^\alpha \rangle$, if $\beta \notin \{\alpha_1, \dots, \alpha_k\}$ and $q(\alpha) = \langle g^\alpha \rangle$ otherwise.

i. Suppose first that $\alpha \in \{\beta_1, \dots, \beta_k\}$.

Then

- A. $\text{dom}((g^\alpha)^{ref}) = (\text{dom}(g^\alpha) \cap \kappa_\beta) \cup \{f^\beta(\xi) \mid \xi \in \text{dom}(g^\alpha) \setminus \kappa_\beta\}$,
- B. for every $\xi \in \text{dom}(g^\alpha) \cap \kappa_\beta = \text{dom}(g^\alpha) \cap \text{dom}((g^\alpha)^{ref})$, $(g^\alpha)^{ref}(\xi) = g^n(\xi)$,
- C. for every $\xi \in \text{dom}(g^\alpha) \setminus \kappa_\beta$, $(g^\alpha)^{ref}(f^\beta(\xi)) = g^\alpha(\xi)$.

It is crucial here that $f^m \upharpoonright (\text{dom}(g^n) \setminus \kappa_m)$ is one to one and the values there are above $\text{rng}(g^n) \cap \kappa_m$.

This follows by conditions (4),(5) of Definitions 1.10,1.11.

ii. Suppose now that $\alpha \notin \{\beta_1, \dots, \beta_k\}$.

Then

- A. $\text{dom}((g^\alpha)^{ref}) = (\text{dom}(g^\alpha) \cap \kappa_\beta) \cup \{f^\beta(\xi) \mid \xi \in \text{dom}(g^\alpha) \setminus \kappa_\beta\}$,
- B. for every $\xi \in \text{dom}(g^\alpha) \cap \kappa_\beta = \text{dom}(g^\alpha) \cap \text{dom}((g^\alpha)^{ref})$, $(g^\alpha)^{ref}(\xi) = g^\alpha(\xi)$,
- C. for every $\xi \in \text{dom}(g^\alpha) \setminus \kappa_\beta$, $(g^\alpha)^{ref}(f^\beta(\xi)) = g^\alpha(\xi)$.

Again, it is crucial here that $f^\beta \upharpoonright (\text{dom}(g^\alpha) \setminus \kappa_\beta)$ is one to one and the values there are above $\text{rng}(g^\alpha) \cap \kappa_\beta$, and this follows by conditions (3),(4) of Definition 2.1 and (4),(5) of Definition 2.2.

One more crucial observation here is that the measure $(E(\alpha))(\text{dom}(g^\alpha))$, to which B^α belongs, reflects to basically the same measure,

It follows by (4) of Definition 2.1.

- D. $A^\alpha \upharpoonright \text{dom}((g^\alpha)^{ref}) \subseteq \{(f^\beta(\xi, \zeta) \mid (\xi, \zeta) \in B^\alpha\}$.

Denote further in this subsection $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}$ by just \mathcal{P} .

The next lemma follows from the definitions:

Lemma 2.5 *For every $\beta < \eta$ and $p \in \mathcal{P}$ with $p(\beta) \in \mathcal{P}_{E(\beta)}^*$ (i.e. non-pure on the coordinate β), the part $\langle \mathcal{P}_{\langle E(\alpha) \mid \alpha < \beta \rangle}, \leq \rangle$ of \mathcal{P} above p has cardinality $f_\beta^\lambda(p(\beta)(\kappa_\beta) < \kappa_\beta$.*

Lemma 2.6 *For every $\beta < \eta$, the forcing $\langle \mathcal{P}_{\langle E(\alpha) \mid \beta \leq \alpha < \eta \rangle}, \leq^* \rangle$ is κ_β -closed.*

Lemma 2.7 *The forcing $\langle \mathcal{P}, \leq \rangle$ satisfies κ_η^{++} -c.c.*

Lemma 2.8 *$\langle \mathcal{P}, \leq, \leq^* \rangle$ is a Prikry type forcing notion.*

Proof. The proof proceeds by induction on the length of the sequence of extenders, i.e. on η . The argument repeats those of Lemma 1.25.

□

Denote for every limit α , $0 < \alpha \leq \eta$, $\bigcup_{\gamma < \alpha} \kappa_\gamma$ by $\bar{\kappa}_\alpha$.

It follows, by the previous lemmas, that the forcing $\langle \mathcal{P}, \leq \rangle$ preserves all the cardinals, but $\bar{\kappa}_\alpha^+$, $0 < \alpha \leq \eta$ a limit ordinal. Using the arguments of the previous lemma it is possible to show that all such cardinals are preserved as well.

Let G be a generic subset of $\langle \mathcal{P}, \leq \rangle$.

Lemma 2.9 *For every limit ordinal μ , $0 < \mu \leq \eta$, $\bar{\kappa}_\mu$ remains a strong limit cardinal in $V[G]$.*

Proof. Given $p \in \mathcal{P}$ and $\beta < \eta$. Suppose that $p(\beta)$ is non-pure. Then $p(\beta)(\kappa_\beta)$ is defined, and hence also the reflection $h_\lambda^\beta(p(\beta)(\kappa_\beta))$ of λ below κ_β . By the definition of the forcing, then the part $\mathcal{P}_{\langle E(\alpha) \mid \alpha < \beta \rangle}$ above p will act as $\mathcal{P}_{\langle E(\alpha) \mid h_\lambda^\beta(p(\beta)(\kappa_\beta)) \mid \alpha < \beta \rangle}$. In particular, $2^{\kappa_\alpha} \leq h_\lambda^\beta(p(\beta)(\kappa_\beta)) < \kappa_\beta$. The upper part of the forcing, i.e. $\mathcal{P}_{\langle E(\alpha) \mid \beta \leq \alpha < \eta \rangle}$, does not add new bounded subsets to κ_β .

So we are done.

□

As in the case $\eta = \omega$, the next lemma is just a variation of the Prikry condition of the forcing.

Lemma 2.10 *Let $p \in \mathcal{P}$ and ζ be a $\langle \mathcal{P}, \leq \rangle$ -name of an ordinal or just $p \Vdash_{\langle \mathcal{P}, \leq \rangle} \zeta$ is an ordinal.*

Then there are $p^ \geq^* p$ and $\alpha_1 < \dots < \alpha_k < \eta$, for some $k < \omega$, such that*

1. *for every i , $1 \leq i \leq k$, $p^*(\alpha_i) = \langle f_{\alpha_i}^{p^*}, A_{\alpha_i}^{p^*} \rangle$,*
2. *for every $\vec{v}_1 \in A_{\alpha_1}^{p^*}, \dots, \vec{v}_k \in A_{\alpha_k}^{p^*}$,
 $p^* \frown \vec{v}_1 \dots \frown \vec{v}_k$ decides ζ .*

Lemma 2.11 *For every limit ordinal μ , $0 < \mu \leq \eta$, $(\bar{\kappa}_\mu^+)^V$ remains a cardinal in $V[G]$.*

The proof of this lemma repeats those of Lemma 1.27.

Given $p \in \mathcal{P}$. Denote by $\text{np}(p)$ the set of all coordinates α of p such that $p(\alpha) \in \mathcal{P}_{E(\alpha)}^*$, i.e. a non-pure extension was made at the coordinate α .

Assume that η is a limit ordinal.

Let $\tau \in [\bar{\kappa}_\eta, \lambda)$ we define in $V[G]$ a function $t_\tau : \eta \rightarrow \bar{\kappa}_\eta$ as follows.
For every $\alpha < \eta$, find $p \in G$ such that $\alpha \in \text{np}(p)$ and if $\alpha_1 < \dots < \alpha_k$ is the increasing enumeration of $\text{np}(p) \setminus \alpha$ (i.e. $\alpha = \alpha_1$), then the following hold:

1. $\tau \in \text{dom}(p(\alpha_k))$.
Set $\tau_k = \tau$.
2. For every $i, 1 \leq i \leq k-1$, $\tau_i \in \text{dom}(p(\alpha_i))$,
where $\tau_i = p(\alpha_{i+1})(\tau_{i+1})$.

Set $t_\tau(\alpha) = p(\alpha)(\tau_1)$.

Lemma 2.12 *In $V[G]$, if $\tau, \rho \in [\bar{\kappa}_\eta, \lambda)$ and $\tau < \rho$, then there is $\alpha^* < \eta$ such that for every $\alpha, \alpha^* \leq \alpha < \eta$, $t_\tau(\alpha) < t_\rho(\alpha)$.*

Proof. Work in V . Let $p \in \mathcal{P}$ be any condition and $\tau, \rho \in [\bar{\kappa}_\eta, \lambda)$, $\tau < \rho$.

Let α^* be a coordinate above $\text{np}(p)$. Then $p(\alpha) = \langle f_\alpha^p, A_\alpha^p \rangle$, for every $\alpha, \alpha^* \leq \alpha < \eta$.

Extend p to p^* by adding τ, ρ to all $\text{dom}(f_\alpha^p)$ with $\alpha^* \leq \alpha < \eta$.

Now, by the definition of the order on \mathcal{P} , for every $\alpha, \alpha^* \leq \alpha < \eta$ and every $q \geq p^*$ such that q defines $t_\tau(\alpha)$ and $t_\rho(\alpha)$, we will have $t_\tau(\alpha) < t_\rho(\alpha)$.

So,

$$p^* \Vdash (\forall \alpha)(\alpha^* \leq \alpha < \eta \rightarrow \underset{\sim}{t}_\tau(\alpha) < \underset{\sim}{t}_\rho(\alpha)).$$

□

It is possible to say a bit more. Namely, let in $V[G]$, for every $\alpha < \eta$, λ_α be the reflection of λ below κ_α , i.e. for some $p \in G$ with $p(\alpha) = f_\alpha^p$, $\lambda_\alpha = h_\lambda^\alpha(f_\alpha^p(\kappa_\alpha))$. Then the following holds:

Lemma 2.13 *The sequence $\langle t_\tau \mid \tau \in [\bar{\kappa}_\eta, \lambda) \rangle$ is a scale in $\langle \prod_{\alpha < \eta} \lambda_\alpha, <_{J^{bd}} \rangle$.*

In particular, we obtain the following:

Corollary 2.14 *It is possible to blow up the power of a singular in the core model² cardinal of arbitrary cofinality in a cardinal preserving extension.*

²Core model with strong cardinals, but below o -hand grenade. It was defined and studied by Ralf Schindler in [8]

3 One generalization.

In the previous section we assumed that $\eta < \kappa_0$ in order to blow up the power of a singular cardinal of cofinality η .

Let us now take η to be an inaccessible cardinal.

Let $\langle \kappa_\alpha \mid \alpha < \eta \rangle$ be now an increasing sequence with limit η and each $E(\alpha)$, for $\alpha < \eta$, be a (κ_α, η) -extender.

Assume that η is the least inaccessible limit of κ_α 's.

We proceed as in the previous section and define $\langle \mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}, \leq, \leq^* \rangle$. It shares the properties of the forcing of the previous section.

Let G be a generic subset of $\langle \mathcal{P}_{\langle E(\alpha) \mid \alpha < \eta \rangle}, \leq \rangle$.

Denote $\bigcup_{\beta < \alpha} \kappa_\beta$ by $\bar{\kappa}_\alpha$, for every $\alpha < \eta$. Then the following holds:

Theorem 3.1 *$V[G]$ is a cofinality preserving extension of V such that for every $\alpha < \eta$, $\bar{\kappa}_\alpha$ is a strong limit singular cardinal with $2^{\bar{\kappa}_\alpha} > \bar{\kappa}_\alpha^+$.*

In addition η remains inaccessible.

By passing to $V[G]_\eta$ we obtain the following:

Corollary 3.2 *It is possible to blow up the power of a proper class club of singular cardinals in the core model in a cofinality preserving extension.*

References

- [1] M. Gitik, Prikry type forcings, in Handbook of Set Theory, Foreman, Kanamori, eds.
- [2] M. Gitik, Short extenders forcings,
- [3] M. Gitik and M. Magidor, Extender based forcings. *J.of Symbolic Logic* 59:2 (1994), 445-460.
- [4] M. Gitik and S. Shelah, On certain indestructibility of strong cardinals and a question of Hajnal. *Archive of Math. Logic* 28 (1989), 35-42.
- [5] M. Gitik and S. Shelah, On some configurations related to SWH. *Arch of Math. Logic* 40 (2001), 639-650.
- [6] C. Merimovich, Supercompact extender based Prikry forcing, *Annals of Pure and Applied Logic*, Volume 168 Number 8, August 2017, pages 1571-1587.
- [7] C. Merimovich, Extender based Magidor-Radin forcing, *Israel Journal of Mathematics*, Volume 182 Number 1, April 2011, pages 439-480.
- [8] R. Schindler, The core model for almost linear iterations, *Annals of Pure and Applied Logic*, Volume 116, Issues 13, 15 August 2002, Pages 205-272.
- [9] S. Shelah, Cardinal arithmetic for sceptics, *Bull. Amer. Math. Soc.* 26 (1992), 197-210.
- [10] S. Shelah, Logical Dreams, *Bull. Amer. Math. Soc.* 40 (2003), 203-228.