

NEW ALGEBRAS OF FUNCTIONS ON TOPOLOGICAL GROUPS ARISING FROM G -SPACES

E. GLASNER AND M. MEGRELISHVILI

ABSTRACT. For a topological group G we introduce the algebra $SUC(G)$ of *strongly uniformly continuous* functions. We show that $SUC(G)$ contains the algebra $WAP(G)$ of weakly almost periodic functions as well as the algebras $LE(G)$ and $Asp(G)$ of locally equicontinuous and Asplund functions respectively. For the Polish groups of order preserving homeomorphisms of the unit interval and of isometries of the Urysohn space of diameter 1, we show that $SUC(G)$ is trivial. We introduce the notion of fixed point on a class P of flows (P -fpp) and study in particular groups with the SUC -fpp. We study the Roelcke algebra ($= UC(G)$ = right and left uniformly continuous functions) and SUC compactifications of the groups $S(\mathbb{N})$, of permutations of a countable set, and $H(C)$, the group of homeomorphisms of the Cantor set. For the first group we show that $WAP(G) = SUC(G) = UC(G)$ and also provide a concrete description of the corresponding metrizable (in fact Cantor) semitopological semigroup compactification. For the second group, in contrast, we show that $SUC(G)$ is properly contained in $UC(G)$. We then deduce that for this group $UC(G)$ does not yield a right topological semigroup compactification.

CONTENTS

1. Introduction	2
2. Actions and G -compactifications	4
3. Cyclic G -systems and point-universality	7
4. Strong Uniform Continuity	10
5. SUC, homogeneity and the epimorphism problem	14
6. Representations of groups and G -spaces on Banach spaces	16
7. Dynamical complexity of functions	21
8. The group $H_+[0, 1]$	27
9. Matrix coefficient characterization of SUC and LE	29
10. Some conclusions about $H_+[0, 1]$ and $Iso(\mathbb{U}_1)$	37
11. Relative extreme amenability: SUC-fpp groups	39
12. The Roelcke compactification of the group $S(\mathbb{N})$	41
13. The homeomorphisms group of the Cantor set	44
References	48

Date: March 26, 2008.

Key words and phrases. Asplund function, fixed point property, G -compactification, G -space, locally equicontinuous, matrix coefficient, right topological semigroup compactification, strongly uniformly continuous.

Research partially supported by BSF (Binational USA-Israel) grant no. 2006119.

1. INTRODUCTION

In this paper we introduce the property of *Strong Uniform Continuity* (in short: SUC) of G -spaces and the associated notion of SUC functions. For every compact G -space X the corresponding orbit maps $\tilde{x} : G \rightarrow X$, $g \mapsto gx$ are right uniformly continuous for every $x \in X$. If all the maps $\{\tilde{x}\}_{x \in X}$ are also left uniformly continuous then we say that X is SUC. Every right uniformly continuous bounded real valued function $f : G \rightarrow \mathbb{R}$ comes from some compact G -space X . That is, there exist a compact G -space X , a continuous function $F : X \rightarrow \mathbb{R}$, and a point $x_0 \in X$ such that $f = F \circ \tilde{x}_0$. We say that f is SUC if it comes from a compact G -space X which is SUC. We denote by $SUC(G)$ the corresponding class of functions on G . The class $SUC(G)$ forms a uniformly closed G -invariant subalgebra of the algebra $UC(G) := RUC(G) \cap LUC(G)$ of (right and left) uniformly continuous functions. Of course we have $SUC(G) = UC(G) = RUC(G) = LUC(G)$ when G is either discrete or abelian so that the notion of strong uniform continuity can be useful only when one deals with non-abelian non-discrete topological groups. Mostly we will be interested in Polish non-locally compact large groups, but some of the questions we study are of interest in the locally compact case as well.

In our recent work [13] we investigated, among other topics, the algebras of *locally equicontinuous* $LE(G)$, and *Asplund* functions $Asp(G)$ on a topological group G . The inclusions $UC(G) \supset SUC(G) \supset LE(G) \supset Asp(G) \supset WAP(G)$ hold for an arbitrary topological group G . In the present article we provide a characterization of the elements of $SUC(G)$ and $LE(G)$ in terms of matrix coefficients for appropriate Banach representations of G by linear isometries.

Intuitively the dynamical complexity of a function $f \in RUC(G)$ can be estimated by the *topological complexity* of the *cyclic* G -flow X_f (the pointwise closure of the left G -orbit $\{gf\}_{g \in G}$ of f) treating it as a subset of the Banach space $RUC(G)$. This leads to a natural dynamical hierarchy (see Theorem 7.12) where $SUC(G)$ plays a basic role. In some sense $SUC(G)$ is the largest “nice subalgebra” of $UC(G)$. It turns out that $f \in SUC(G)$ iff X_f is a subset of $UC(G)$. Moreover the algebra $SUC(G)$ is *point-universal* in the sense of [13] and every other point-universal subalgebra of $UC(G)$ is contained in $SUC(G)$. Recall that a G -algebra $\mathcal{A} \subset RUC(G)$ is point-universal if and only if the associated G -compactification $G \rightarrow G^{\mathcal{A}}$ is a right topological semigroup compactification.

As an application we conclude that the algebra $UC(G)$ is point-universal if and only if it coincides with the algebra $SUC(G)$ and that the corresponding *Roelcke compactification* $G \rightarrow G^{UC}$ is in general not a right topological semigroup compactification of G (in contrast to the compactification $G \rightarrow G^{SUC}$ determined by the algebra $SUC(G)$).

For locally compact groups, $SUC(G)$ contains the subalgebra $C_0(G)$ consisting of the functions which vanish at infinity, and therefore determines the topology of G . The structure of $SUC(G)$ — in contrast to $RUC(G)$ which is always huge for non-precompact groups — is “computable” for several large groups like: $H_+[0, 1]$, $Iso(\mathbb{U}_1)$ (the isometry group of the Urysohn space of diameter one \mathbb{U}_1), $U(H)$ (the unitary group on an infinite dimensional Hilbert space), $S_\infty = S(\mathbb{N})$ (the Polish infinite symmetric group) and any noncompact connected simple Lie group with finite center (e.g., $SL_n(\mathbb{R})$). For instance, $SUC(G) = WAP(G)$ for $U(H)$, S_∞ and $SL_n(\mathbb{R})$. In the first case we use a result of Uspenskij [46] which identifies the Roelcke completion of $U(H)$ as the compact semigroup of contracting operators on the Hilbert space H . For S_∞ see Section 12, and for $SL_n(\mathbb{R})$ this follows from an old result of Veech [49].

The group $H_+[0, 1]$ of orientation preserving homeomorphisms of the closed unit interval, endowed with the compact open topology is a good test case in the class of “large” yet “computable” topological groups. See Section 8 for more details on this group. In particular recall the result from [28] which shows that $H_+[0, 1]$ is WAP-*trivial*: Every weakly almost periodic function on $G := H_+[0, 1]$ is a constant. Equivalently, G is *reflexively trivial*, that is, every continuous representation $G \rightarrow Iso(V)$ where V is a reflexive Banach space is trivial.

Here we show that G is even “SUC-trivial” — that is, the algebra $SUC(G)$ (and hence, also the algebras $LE(G)$ and $Asp(G)$) consists only of constant functions — and that every continuous representation of G into the group of linear isometries $Iso(V)$ of an Asplund Banach space V is trivial. Since in general $WAP(G) \subset Asp(G)$ and since every reflexive Banach space is Asplund these results strengthen the main results of [28]. SUC-triviality implies that every *adjoint continuous* (see Section 6) representation is trivial for $H_+[0, 1]$. The latter fact follows also from a recent unpublished result of Uspenskij (private communication).

From the WAP-triviality (equivalently, reflexive triviality) of $H_+[0, 1]$ and results of Uspenskij about $Iso(\mathbb{U}_1)$ Pestov deduces in [39, Corollary 1.4] the fact that the group $Iso(\mathbb{U}_1)$ is also WAP-trivial. Using a similar idea and the matrix coefficient characterization of SUC one can conclude that $Iso(\mathbb{U}_1)$ is SUC-trivial. It is an open question whether the group $H([0, 1]^\omega)$ is SUC-trivial (or, WAP-trivial).

The above mentioned description of $SUC(G)$ and $LE(G)$ in terms of matrix coefficients (Section 9) is nontrivial. The proof is based on a dynamical modification of a well known interpolation technique of Davis, Figiel, Johnson and Pelczyński [7].

In section 11 we introduce a notion of extreme amenability with respect to a class of flows. In particular we examine *extreme SUC-amenability* and *extreme*

SUC-amenable groups. Namely those groups which have a fixed point property on compact SUC G -spaces. Several natural groups, like $SL_2(\mathbb{R})$, S_∞ , $H(C)$ (the homeomorphisms group of the Cantor set), and $H_+(\mathbb{S})$, which fail to be extremely amenable, are however extremely SUC-amenable. In the last two sections we study the Roelcke and SUC compactifications of the groups S_∞ and $H(C)$. For the first group we show that $WAP(S_\infty) = SUC(S_\infty) = UC(S_\infty)$ and also provide a concrete description of the corresponding metrizable (in fact Cantor) semitopological semigroup compactification. For the latter group $G := H(C)$, in contrast, we have $SUC(G) \subsetneq UC(G)$ from which fact we deduce that the corresponding Roelcke compactification $G \rightarrow G^{UC}$ is not a right topological semigroup compactification of G .

Finally let us note that although in this work we consider, for convenience, algebras of real-valued functions, it seems that there should be no difficulty in extending our definitions and results to the complex case.

2. ACTIONS AND G -COMPACTIFICATIONS

Unless explicitly stated otherwise, all spaces in this paper are at least Tychonoff. A (*left*) *action* of a topological group G on a topological space X is defined by a function $\pi : G \times X \rightarrow X$, $\pi(g, x) := gx$ such that always $g_1(g_2x) = (g_1g_2)x$ and $ex = x$ hold, where $e = e_G$ is the neutral element of G . Every $x \in X$ defines an *orbit map* $\tilde{x} : G \rightarrow X$, $g \mapsto gx$. Also every $g \in G$ induces a *g -translation* $\pi^g : X \rightarrow X$, $x \mapsto gx$. If the action π is continuous then we say that X is a *G -space* (or a *G -system* or a *G -flow*). Sometimes we denote it as a pair (G, X) . If the orbit Gx_0 of x_0 is dense in X for some $x_0 \in X$ then the G -space X is *point transitive* (or just *transitive*) and the point x_0 is a *transitive point*. If X in addition is compact then the pair (X, x_0) is said to be a *pointed system* or a *G -ambit*. If every point x in a compact G -space X is transitive then X is said to be *minimal*.

Let G act on X_1 and on X_2 . A continuous map $f : X_1 \rightarrow X_2$ is a *G -map* (or a *homomorphism* of dynamical systems) if $f(gx) = gf(x)$ for every $(g, x) \in G \times X_1$.

A *right action* $X \times G \rightarrow X$ can be defined analogously. If G^{op} is the *opposite group* of G with the same topology then the right G -space (X, G) can be treated as a left G^{op} -space (G^{op}, X) (and vice versa). A map $h : G_1 \rightarrow G_2$ between two groups is a *co-homomorphism* (or, an *anti-homomorphism*) if $h(g_1g_2) = h(g_2)h(g_1)$. This happens iff $h : G_1^{op} \rightarrow G_2$ (the same assignment) is a homomorphism.

The Banach algebra (under the supremum norm) of all continuous real valued bounded functions on a topological space X will be denoted by $C(X)$. Let (G, X) be a left (not necessarily compact) G -space. Then it induces the right

action $C(X) \times G \rightarrow C(X)$, with $(fg)(x) = f(gx)$, and the corresponding cohomomorphism $h : G \rightarrow Iso(C(X))$. While the g -translations $C(X) \rightarrow C(X)$ (being isometric) are continuous, the orbit maps $\tilde{f} : G \rightarrow C(X)$, $g \mapsto fg$ are not necessarily continuous. The function $f \in C(X)$ is *right uniformly continuous* if the orbit map $G \rightarrow C(X)$, $g \mapsto fg$ is norm continuous. The set $RUC(X)$ of all right uniformly continuous functions on X is a uniformly closed G -invariant subalgebra of $C(X)$. Here and in the sequel “subalgebra” means a uniformly closed unital (containing the constants) subalgebra. A “ G -subalgebra” is an algebra which is invariant under the natural right action of G .

Every topological group G can be treated as a G -space under the left regular action of G on itself. In this particular case $f \in RUC(G)$ iff f is uniformly continuous with respect to the *right uniform structure* \mathcal{R} on G (furthermore this is also true for coset G -spaces G/H).

Thus, $f \in RUC(G)$ iff for every $\varepsilon > 0$ there exists a neighborhood V of the identity element $e \in G$ such that $\sup_{g \in G} |f(vg) - f(g)| < \varepsilon$ for every $v \in V$.

Analogously one defines *right translations* $(gf)(x) := f(xg)$, and the algebra $LUC(G)$ of *left uniformly continuous* functions. These are the functions which are uniformly continuous with respect to the *left uniform structure* \mathcal{L} on G .

A G -compactification of a G -space X is a G -map $\nu : X \rightarrow Y$ into a compact G -space Y with $\text{cl } \nu(X) = Y$. A compactification is *proper* when ν is a topological embedding. Given a compact G -space X and a point $x_0 \in X$ the map $\nu : G \rightarrow X$ defined by $\nu(g) = gx_0$ is a compactification of the G -space G (the left regular action) in the orbit closure $\text{cl } Gx_0 \subset X$.

We say that a G -compactification $\nu : G \rightarrow S$ of $X := G$ is a *right topological semigroup compactification* of G if S is a *right topological semigroup* (that is, S is a compact semigroup such that for every $p \in S$ the map $S \rightarrow S, s \mapsto sp$ is continuous) and ν is a homomorphism of semigroups.

There exists a canonical 1-1 correspondence between the G -compactifications of X and G -subalgebras of $RUC(X)$ (see for example [50]). The compactification $\nu : X \rightarrow Y$ induces an isometric G -embedding of G -algebras

$$j_\nu : C(Y) = RUC(Y) \hookrightarrow RUC(X), \quad \phi \mapsto \phi \circ \nu$$

and the algebra \mathcal{A}_ν is defined as the image $j_\nu(C(Y))$. Conversely, if \mathcal{A} is a G -subalgebra of $RUC(X)$, then denote by $X^\mathcal{A}$ or by $|\mathcal{A}|$ the corresponding Gelfand space treating it as a weak star compact subset of the dual space \mathcal{A}^* . It has a structure of a G -space $(G, |\mathcal{A}|)$ and the natural map $\nu_\mathcal{A} : X \rightarrow X^\mathcal{A}$, $x \mapsto \text{eva}_x$, where $\text{eva}_x(\varphi) := \varphi(x)$, is the evaluation at x (a multiplicative functional), defines a G -compactification. If $\nu_1 : X \rightarrow X^{\mathcal{A}_1}$ and $\nu_2 : X \rightarrow X^{\mathcal{A}_2}$ are two G -compactifications then $\mathcal{A}_{\nu_1} \subset \mathcal{A}_{\nu_2}$ iff $\nu_1 = \alpha \circ \nu_2$ for some G -map $\alpha : X^{\mathcal{A}_2} \rightarrow X^{\mathcal{A}_1}$.

The algebra \mathcal{A} determines the compactification $\nu_{\mathcal{A}}$ uniquely, up to the equivalence of G -compactifications. The G -algebra $RUC(X)$ defines the corresponding Gelfand space $|RUC(X)|$, which we denote by $\beta_G X$, and the *maximal G -compactification* $i_{\beta} : X \rightarrow \beta_G X$. Note that this map may not be an embedding even for Polish G and X (see [24]); it follows that there is no proper G -compactification for such X . If X is a compact G -space then $\beta_G X$ can be identified with X and $C(X) = RUC(X)$.

Denote by G^{RUC} the Gelfand space of the G -algebra $RUC(G)$. The canonical embedding $u : G \rightarrow G^{RUC}$ defines the *greatest ambit* $(G^{RUC}, u(e))$ of G .

It is easy to see that the intersection $UC(G) := RUC(G) \cap LUC(G)$ is a left and right G -invariant closed subalgebra of $RUC(G)$. We denote the corresponding compactification by G^{UC} . Denote by $\mathcal{L} \wedge \mathcal{R}$ the *lower uniformity* of G . It is the infimum (greatest lower bound) of the left and right uniformities on G ; we call it the *Roelcke uniformity*. Clearly, for every bounded function $f : G \rightarrow \mathbb{R}$ we have $f \in UC(G)$ iff $f : (G, \mathcal{L} \wedge \mathcal{R}) \rightarrow \mathbb{R}$ is uniformly continuous. Recall the following important fact (in general the infimum $\mu_1 \wedge \mu_2$ of two compatible uniform structures on a topological space X need not be compatible with the topology of X).

Lemma 2.1. (1) (Roelcke-Dierolf [41]) *For every topological group G the Roelcke uniform structure $\mathcal{L} \wedge \mathcal{R}$ generates the given topology of G .*
(2) *For every topological group G the algebra $UC(G)$ separates points from closed subsets in G .*

Proof. (1) See Roelcke-Dierolf [41, Proposition 2.5].

(2) Follows from (1). □

By a *uniform G -space* (X, μ) we mean a G -space (X, τ) where τ is the (completely regular) topology defined by the uniform structure μ and the g -translations ($g \in G$) are uniform isomorphisms.

Let $X := (X, \mu)$ be a uniform G -space. A point $x_0 \in X$ is a *point of equicontinuity* (notation: $x_0 \in Eq_X$) if for every entourage $\varepsilon \in \mu$, there is a neighborhood O of x_0 such that $(gx_0, gx) \in \varepsilon$ for every $x \in O$ and $g \in G$. The G -space X is *equicontinuous* if $Eq_X = X$. (X, μ) is *uniformly equicontinuous* if for every $\varepsilon \in \mu$ there is $\delta \in \mu$ such that $(gx, gy) \in \varepsilon$ for every $g \in G$ and $(x, y) \in \delta$. For compact X (equipped with the unique compatible uniformity), equicontinuity and uniform equicontinuity coincide. Compact (uniformly) equicontinuous G -space X is also said to be *Almost Periodic* (in short: AP); see also Section 7. If Eq_X is dense in X then (X, μ) is said to be an *almost equicontinuous* (AE) G -space [2].

The following definition is standard (for more details see for example [13]).

Definition 2.2. (1) A function $f \in C(X)$ on a G -space X comes from a compact G -system Y if there exist a G -compactification $\nu : X \rightarrow Y$ (so, ν is onto if X is compact) and a function $F \in C(Y)$ such that $f = F \circ \nu$ (i.e., if $f \in \mathcal{A}_\nu$). Then necessarily, $f \in RUC(X)$.

$$\begin{array}{ccc} X & \xrightarrow{\nu} & Y \\ & \searrow f & \downarrow F \\ & & \mathbb{R} \end{array}$$

- (2) A function $f \in RUC(G)$ comes from a pointed system (Y, y_0) if for some continuous function $F \in C(Y)$ we have $f(g) = F(gy_0)$, $\forall g \in G$. Notation: $f \in \mathcal{A}(Y, y_0)$. Defining $\nu : X = G \rightarrow Y$ by $\nu(g) = gy_0$ observe that this is indeed a particular case of 2.2.1.
- (3) Let Γ be a class of compact G -spaces. For a G -space X denote by $\Gamma(X)$ the class of all functions on X which come from a G -compactification $\nu : X \rightarrow Y$ where the G -system Y belongs to Γ .

Let P be a class of compact G -spaces which is preserved by G -isomorphisms, products and closed G -subspaces. It is well known (see for example, [13, Proposition 2.9]) that for every G -space X there exists a *universal* (maximal) G -compactification $X \rightarrow X^P$ such that X^P lies in P . More precisely, for every (not necessarily compact) G -space X denote by $\mathcal{P} \subset C(X)$ the collection of functions coming from G -spaces having property P . Then \mathcal{P} is a uniformly closed, G -invariant subalgebra of $RUC(X)$ and the maximal G -compactification of X with property P is the corresponding Gelfand space $X^P := |\mathcal{P}|$. If X is compact then (G, X^P) is the maximum factor of (G, X) with property P . In particular let P be one of the following natural classes of compact G -spaces: a) Almost Periodic (=equicontinuous); b) Weakly Almost Periodic; c) Hereditarily Not Sensitive; d) Locally Equicontinuous; e) all compact G -spaces. Then, in this way the following maximal (in the corresponding class) G -compactifications are: a) G^{AP} ; b) G^{WAP} ; c) G^{Asp} ; d) G^{LE} ; e) G^{RUC} .

For undefined concepts and more details see Section 7 and also [13].

3. CYCLIC G -SYSTEMS AND POINT-UNIVERSALITY

Here we give some background material about *cyclic* compact G -systems X_f defined for $f \in RUC(X)$. These G -spaces play a significant role in many aspects of topological dynamics and are well known at least for the particular case of $X := G$. We mostly use the presentation and results of [13] (see also [51]).

As a first motivation note a simple fact about Definition 2.2. For every G -space X a function $f : X \rightarrow \mathbb{R}$ lies in $RUC(X)$ iff it comes from a compact G -flow Y . We can choose Y via the maximal G -compactification $G \rightarrow \beta_G G = Y$ of G . This is the largest possibility in this setting. Among all possible G -compactifications

$\nu : X \rightarrow Y$ of X such that f comes from (ν, Y) there exists also the smallest one. Take simply the smallest G -subalgebra \mathcal{A}_f of $RUC(X)$ generated by the orbit fG of f in $RUC(X)$. Denote by X_f the Gelfand space $|\mathcal{A}_f| = X^{\mathcal{A}_f}$ of the algebra \mathcal{A}_f . Then the corresponding G -compactification $X \rightarrow Y := X_f$ is the desired one. We call \mathcal{A}_f and X_f the *cyclic G -algebra* and *cyclic G -system* of f , respectively. Next we provide an alternative construction and some basic properties of X_f .

Let X be a (not necessarily compact) G -space. Given $f \in RUC(X)$ let $I = [-\|f\|, \|f\|] \subset \mathbb{R}$ and $\Omega = I^G$, the product space equipped with the compact product topology. We let G act on Ω by $g\omega(h) = \omega(hg)$, $g, h \in G$. Define the continuous map

$$f_{\sharp} : X \rightarrow \Omega, \quad f_{\sharp}(x)(g) = f(gx)$$

and the closure $X_f := \text{cl}(f_{\sharp}(X))$ in Ω . Note that $X_f = f_{\sharp}(X)$ whenever X is compact.

Denoting the unique continuous extension of f to $\beta_G X$ by \tilde{f} (it exists because $f \in RUC(X)$) we now define a map

$$\psi : \beta_G X \rightarrow X_f \quad \text{by} \quad \psi(y)(g) = \tilde{f}(gy), \quad y \in \beta_G X, \quad g \in G.$$

Let $pr_e : \Omega \rightarrow \mathbb{R}$ denote the projection of $\Omega = I^G$ onto the e -coordinate and let $F_e := pr_e \upharpoonright_{X_f} : X_f \rightarrow \mathbb{R}$ be its restriction to X_f . Thus, $F_e(\omega) := \omega(e)$ for every $\omega \in X_f$.

As before denote by \mathcal{A}_f the smallest (closed and unital, of course) G -invariant subalgebra of $RUC(X)$ which contains f . There is then a naturally defined G -action on the Gelfand space $X^{\mathcal{A}_f} = |\mathcal{A}_f|$ and a G -compactification (morphism of dynamical systems if X is compact) $\pi_f : X \rightarrow |\mathcal{A}_f|$. Next consider the map $\pi : \beta_G X \rightarrow |\mathcal{A}_f|$, the canonical extension of π_f induced by the inclusion $\mathcal{A}_f \subset RUC(X)$.

The action of G on Ω is not in general continuous. However, the restricted action on X_f is continuous for every $f \in RUC(X)$. This follows from the second assertion of the next fact.

Proposition 3.1. (See for example [13])

- (1) Each $\omega \in X_f$ is an element of $RUC(G)$. That is, $X_f \subset RUC(G)$.
- (2) The map $\psi : \beta_G X \rightarrow X_f$ is a continuous homomorphism of G -systems. The dynamical system $(G, |\mathcal{A}_f|)$ is isomorphic to (G, X_f) and the diagram

$$\begin{array}{ccccc}
 X & \xrightarrow{i_{\beta}} & \beta_G X & & \\
 \pi_f \downarrow & \searrow \pi & \downarrow \psi & \searrow \tilde{f} & \\
 |\mathcal{A}_f| & \xleftrightarrow{\quad} & X_f & \xrightarrow{F_e} & \mathbb{R}
 \end{array}$$

commutes.

- (3) $f = F_e \circ f_{\sharp}$. Thus every $f \in RUC(X)$ comes from the system X_f . Moreover, if f comes from a system Y and a G -compactification $\nu : X \rightarrow Y$

then there exists a homomorphism $\alpha : Y \rightarrow X_f$ such that $f_{\sharp} = \alpha \circ \nu$. In particular, $f \in \mathcal{A}_f \subset \mathcal{A}_\nu$.

If $X := G$ with the usual left action then X_f is the pointwise closure of the G -orbit $Gf := \{gf\}_{g \in G}$ of f in $RUC(G)$. Hence (X_f, f) is a transitive pointed G -system.

As expected by the construction the cyclic G -systems X_f provide “building blocks” for compact G -spaces. That is, every compact G -space can be embedded into the G -product of G -spaces X_f .

Let us say that a topological group G is *uniformly Lindelöf* if for every nonempty open subset $O \subset G$ countably many translates $g_n O$ cover G (there are several alternative names for this notion: ω -bounded, ω -bounded, ω -narrow, ω -precompact). It is well known that G is uniformly Lindelöf iff G is a topological subgroup in a product of second countable groups. When G is uniformly Lindelöf (e.g. when G is second countable) the compactum X_f is metrizable.

The question “when is X_f a subset of $UC(G)$?” provides another motivation for introducing the notion of strongly uniformly continuous (SUC) functions (see Definition 4.2 and Theorem 4.12).

The *enveloping (or Ellis) semigroup* $E = E(X)$ of a compact G -space X is defined as the closure in X^X (with its compact pointwise convergence topology) of the set $\{\pi^g : X \rightarrow X\}_{g \in G}$ of translations considered as a subset of X^X . With the operation of composition of maps this is a right topological semigroup. Moreover, the map

$$j = j_X : G \rightarrow E(X), \quad g \mapsto \pi^g$$

is a right topological semigroup compactification of G . The compact space $E(X)$ becomes a G -space with respect to the natural action

$$G \times E(X) \rightarrow E(X), \quad (gp)(x) = gp(x).$$

Moreover the pointed G -system $(E(X), j(e))$ is point-universal in the following sense.

Definition 3.2. ([13]) A pointed G -system (X, x_0) is *point-universal* if it has the property that for every $x \in X$ there is a homomorphism $\pi_x : (X, x_0) \rightarrow (\text{cl}(Gx), x)$. The G -subalgebra $\mathcal{A} \subset RUC(G)$ is said to be *point-universal* if the corresponding G -ambit $(G^{\mathcal{A}}, u_{\mathcal{A}}(e))$ is point-universal.

We will use the following characterization of point-universality from [13].

Lemma 3.3. *Let (X, x_0) be a transitive compact G -system. The following conditions are equivalent:*

- (1) *The system (X, x_0) is point-universal.*
- (2) *The orbit map $G \rightarrow X, g \mapsto gx_0$ is a right topological semigroup compactification of G .*

- (3) (X, x_0) is G -isomorphic to its enveloping semigroup $(E(X), j(e))$.
- (4) $\mathcal{A}(X, x_0) = \bigcup_{x \in X} \mathcal{A}(\text{cl}(Gx), x)$.
- (5) $X_f \subset \mathcal{A}(X, x_0)$ for every $f \in \mathcal{A}(X, x_0)$ (where $\mathcal{A}(X, x_0)$ is the corresponding subalgebra of $RUC(G)$ coming from the G -compactification $\nu : G \rightarrow X$, $\nu(g) = gx_0$).

In particular, for every right topological semigroup compactification $\nu : G \rightarrow S$ the pointed G -space $(S, \nu(e))$ is point-universal. For other properties of point-universality see [13] and Remark 9.4.

4. STRONG UNIFORM CONTINUITY

Let G be a topological group. As before denote by \mathcal{L} and \mathcal{R} the left and right uniformities on G . We start with a simple observation.

Lemma 4.1. *For every compact G -space X the corresponding orbit maps*

$$\tilde{x} : (G, \mathcal{R}) \rightarrow (X, \mu_X), \quad g \mapsto gx$$

are uniformly continuous for every $x \in X$.

Proof. Let V be an open neighborhood of the diagonal $\Delta \subset X \times X$. In order to obtain a contradiction suppose that for every neighborhood U of $e \in G$ there is $u_U \in U$ and $x_U \in X$ such that $(x_U, u_U x_U) \notin V$. For a convergent subnet we have $\lim(x_U, u_U x_U) = (x, x') \notin V$ contradicting the joint continuity of the G -action. \square

In general, for non-commutative groups, one cannot replace \mathcal{R} by the left uniformity \mathcal{L} (see Remark 4.4). This leads us to the following definition.

Definition 4.2. Let G be a topological group.

- (1) We say that a uniform G -space (X, μ) is *strongly uniformly continuous at* $x_0 \in X$ (notation: $x_0 \in SUC_X$) if the orbit map $\tilde{x}_0 : G \rightarrow X$, $g \mapsto gx_0$ is (\mathcal{L}, μ) -uniformly continuous. Precisely, this means that for every $\varepsilon \in \mu$ there exists a neighborhood U of $e \in G$ such that

$$(gux_0, gx_0) \in \varepsilon$$

for every $g \in G$ and every $u \in U$. If $SUC_X = X$ we say that X is *strongly uniformly continuous*.

- (2) If X is a compact G -space then there exists a unique compatible uniformity μ_X on X . So, SUC_X is well defined. By Lemma 4.1 it follows that a compact G -space X is SUC at x_0 iff $\tilde{x}_0 : G \rightarrow X$ is $(\mathcal{L} \wedge \mathcal{R}, \mu_X)$ -uniformly continuous. We let SUC denote the class of all compact G -systems such that (X, μ_X) is SUC.
- (3) A function $f \in C(X)$ is *strongly uniformly continuous* (notation: $f \in SUC(X)$) if it comes from a SUC compact dynamical system.

- (4) Let x_1 and x_2 be points of a G -space X . Write $x_1 \overset{SUC}{\sim} x_2$ if these points cannot be separated by a SUC function on X . Equivalently, this means that these points have the same images under the *universal SUC compactification* G -map $X \rightarrow X^{SUC}$ (see Lemma 4.6.1 below).

Lemma 4.3. $SUC(G) \subset UC(G)$.

Proof. Let $f : G \rightarrow \mathbb{R}$ belong to $SUC(G)$. Then it comes from a function $F : X \rightarrow \mathbb{R}$, where $\nu : G \rightarrow X$ is a G -compactification of G such that $f(g) = F(\nu(g))$, and $F \in SUC(X)$. Clearly, F is uniformly continuous because X is compact. Then $f \in RUC(G)$ by Lemma 4.1. In order to see that $f \in LUC(G)$, choose $x_0 := \nu(e) \in X$ in the definition of SUC . \square

Remark 4.4. Recall that if $\mathcal{L} = \mathcal{R}$ then G is said to be SIN group. If G is a SIN group then $X \in SUC$ for every compact G -space X . It follows that for a SIN group G we have $SUC(X) = RUC(X)$ for every, not necessarily compact, G -space X and also $SUC(G) = UC(G) = RUC(G)$. For example this holds for abelian, discrete, and compact groups.

A special case of a SUC uniform G -space is obtained when the uniform structure μ is defined by a G -invariant metric. If μ is a metrizable uniformity then (X, μ) is uniformly equicontinuous iff μ can be generated by a G -invariant metric on X . A slightly sharper property is the local version: $SUC_X \supset Eq_X$ (see Lemma 7.8).

We say that a compactification of G is *Roelcke* if the corresponding algebra \mathcal{A} is a G -subalgebra of $UC(G)$, or equivalently, if there exists a natural G -morphism $G^{UC} \rightarrow G^{\mathcal{A}}$.

Lemma 4.5. (1) Let $f : (X, \mu) \rightarrow (Y, \eta)$ be a uniformly continuous G -map.

Then $f(SUC_X) \subset SUC_Y$.

(2) The class SUC is closed under products, subsystems and quotients.

(3) Let $\alpha : G \rightarrow Y$ be a Roelcke compactification. Then $\alpha(G) \subset SUC_Y$.

(4) Let X be a not necessarily compact G -space and $f \in SUC(X)$. Then for every $x_0 \in X$ and every $\varepsilon > 0$ there exists a neighborhood U of e such that

$$|f(gux_0) - f(gx_0)| < \varepsilon \quad \forall (g, u) \in G \times U.$$

Proof. (1) and (2) are straightforward.

(3): Follows directly from (1) because the left action of G on itself is uniformity equicontinuous with respect to the left uniformity and $\alpha : (G, \mathcal{L}) \rightarrow (Y, \mu_Y)$ is uniformly continuous for every Roelcke compactification.

(4): There exist: a compact SUC G -space Y , a continuous function $F : Y \rightarrow \mathbb{R}$ and a G -compactification $\nu : X \rightarrow Y$ such that $f = F \circ \nu$. Now our assertion

follows from the fact that $\nu(x_0) \in SUC_Y$ for every $x_0 \in X$ (taking into account that F is uniformly continuous). \square

By Lemmas 4.5.2, 3.3 and the standard subdirect product construction (see [13, Proposition 2.9.2]) we can derive the following facts.

- Lemma 4.6.** (1) *The collection $SUC(X)$ is a G -subalgebra of $RUC(X)$ for every G -space X and the corresponding Gelfand space $|SUC(X)| = X^{SUC}$ with the canonical compactification $j : X \rightarrow X^{SUC}$ is the maximal SUC -compactification of X .*
- (2) *A compact G -system X is SUC if and only if $C(X) = SUC(X)$.*
- (3) *For every $f \in RUC(X)$ we have $f \in SUC(X)$ if and only if X_f is SUC .*
- (4) *$SUC(G)$ is a point-universal closed G -subalgebra of $RUC(G)$.*
- (5) *The canonical compactification $j : G \rightarrow G^{SUC}$ is always a right topological semigroup compactification of G .*

We need also the following link between SUC functions and cyclic G -spaces.

Lemma 4.7. *Let X be a G -space, $f \in RUC(X)$ and $h \in X_f$. Then the following are equivalent:*

- (1) $h \in LUC(G)$.
- (2) $h \in UC(G)$.
- (3) h , as a point in the G -flow $Y := X_f$, is in SUC_Y .

Proof. We know by Proposition 3.1.1 that $X_f \subset RUC(G)$. Thus (1) \iff (2).

For (1) \iff (3) observe that

$$|h(gu) - h(g)| < \varepsilon \quad \forall g \in G \quad \iff \quad |h(t_k gu) - h(t_k g)| < \varepsilon \quad \forall g \in G, k = 1, \dots, n$$

for arbitrary finite subset $\{t_1, \dots, t_n\}$ of G . \square

The following result shows that $SUC(X)$ can be described by internal terms for every compact G -space X .

Proposition 4.8. *Let X be a compact G -space. The following are equivalent:*

- (1) $f \in SUC(X)$.
- (2) *For every $x_0 \in X$ and every $\varepsilon > 0$ there exists a neighborhood U of e such that*

$$|f(gux_0) - f(gx_0)| < \varepsilon \quad \forall (g, u) \in G \times U.$$

Proof. (1) \implies (2): Apply Lemma 4.5.4.

(1) \impliedby (2): By Lemma 4.6.3 we have to show that the cyclic G -space X_f is SUC . Fix an arbitrary element $\omega \in X_f$. According to Lemma 4.7 it is equivalent to verify that $\omega \in LUC(G)$. The G -compactification map $f_{\sharp} : X \rightarrow X_f$ is onto because X is compact. Choose $x_0 \in X$ such that $f_{\sharp}(x_0) = \omega$. Then $\omega(g) = f(gx_0)$

for every $g \in G$. By assertion (2) for $x_0 \in X$ and ε we can pick a neighborhood U of e such that

$$|f(gux_0) - f(gx_0)| \leq \varepsilon \quad \forall (g, u) \in G \times U.$$

holds. Now we can finish the proof by observing that

$$|f(gux_0) - f(gx_0)| = |\omega(gu) - \omega(g)| \leq \varepsilon \quad \forall (g, u) \in G \times U.$$

□

The following result emphasizes the differences between $RUC(G)$ and $SUC(G)$.

Theorem 4.9. *Let $\alpha = \alpha_{\mathcal{A}} : G \rightarrow S$ be a G -compactification of G such that the corresponding left G -invariant subalgebra \mathcal{A} of $RUC(G)$ is also right G -invariant. Consider the following conditions:*

- (1) $\mathcal{A} \subset UC(G)$ (that is, $\alpha : G \rightarrow S$ is a Roelcke compactification).
- (2) The induced right action $S \times G \rightarrow S$, $(s, g) \mapsto s\alpha(g)$ is jointly continuous.
- (3) $\mathcal{A} \subset SUC(G)$.

Then

- (a) always, $1 \Leftrightarrow 2$ and $3 \Rightarrow 1$.
- (b) if, in addition, S is a right topological semigroup and $\alpha : G \rightarrow S$ is a right topological semigroup compactification of G then $1 \Leftrightarrow 2 \Leftrightarrow 3$.

Proof. (a) $1 \Leftrightarrow 2$: By our assumption \mathcal{A} is G -invariant with respect to left and right translations (that is, the functions $(fg)(x) := f(gx)$ and $(gf)(x) := f(xg)$ lie in \mathcal{A} for every $f \in \mathcal{A}$ and $g \in G$). Then the corresponding (weak star compact) Gelfand space $S := X^{\mathcal{A}} \subset \mathcal{A}^*$, admits the natural *dual left and right actions* (see also Definition 6.2 and Remark 6.7) $\pi_l : G \times S \rightarrow S$ and $\pi_r : S \times G \rightarrow S$ such that $(g_1s)g_2 = g_1(sg_2)$ for every $(g_1, s, g_2) \in G \times S \times G$. It is easy to see that this right action $S \times G \rightarrow S$ is jointly continuous if and only if $\mathcal{A} \subset LUC(G)$. On the other hand, since $\alpha : G \rightarrow S$ is a G -compactification of left G -spaces we already have $\mathcal{A} \subset RUC(G)$.

$3 \Rightarrow 1$: By Lemma 4.3 we have $SUC(G) \subset UC(G)$.

(b) We have to verify that $1 \Rightarrow 3$ provided that $\alpha : G \rightarrow S$ is a right topological semigroup compactification of G . The latter condition is equivalent to the fact that the system (G, S) is point universal (Lemma 3.3) and thus for every $x_0 \in S$ there exists a homomorphism of G -ambits $\phi : (S, \alpha(e)) \rightarrow (\text{cl}(Gx_0), x_0)$. By Lemma 4.5 we conclude that the point $x_0 = \phi(\alpha(e)) \in S$ is a point of SUC in the G -system $\text{cl}(Gx_0)$ (and, hence, in S). Since x_0 is an arbitrary point in S we get that $SUC_S = S$ and hence S is an SUC system. Since every function $f \in \mathcal{A}$ on G comes from the compactification $\alpha : G \rightarrow S$ we conclude that $\mathcal{A} \subset SUC(G)$. □

Corollary 4.10. *The G -compactification $j : G \rightarrow G^{SUC}$ is a right topological semigroup compactification of G such that the right action $G^{SUC} \times G \rightarrow G^{SUC}$ is also jointly continuous.*

Proof. Apply Proposition 4.6.5 and Theorem 4.9. □

Corollary 4.11. *The following conditions are equivalent:*

- (1) $i : G \rightarrow G^{UC}$ is a right topological semigroup compactification.
- (2) $(G^{UC}, i(e))$ is a point universal G -system.
- (3) G^{UC} is SUC.
- (4) $SUC(G) = UC(G)$.

Proof. Apply assertion (b) of Theorem 4.9 to $\mathcal{A} = UC(G)$ taking into account Lemmas 3.3 and 4.3. □

Particularly interesting examples of groups G with $SUC(G) = UC(G)$ are the Polish groups $U(H)$ of all unitary operators (Example 7.13), and the group $S_\infty(\mathbb{N})$ (Theorem 12.2). In both cases we actually have $SUC(G) = UC(G) = WAP(G)$. Note that these groups are not SIN (compare Remark 4.4).

For the next result see also Veech [49, Section 5].

Theorem 4.12. *Let $f \in RUC(X)$. The following conditions are equivalent:*

- (1) $X_f \subset UC(G)$.
- (2) (G, X_f) is SUC.
- (3) $f \in SUC(X)$.

Proof. 1 \Rightarrow 2: Let $h \in X_f$. By our assumption we have $h \in UC(G) \subset LUC(G)$. Then by Lemma 4.7, h , as a point in the G -flow $Y := X_f$, is in SUC_Y . So $SUC_Y = Y$. This means that (G, X_f) is SUC.

2 \Rightarrow 3: Let (G, X_f) be SUC. By Proposition 3.1.3, the function $f : X \rightarrow \mathbb{R}$ comes from the G -compactification $f_\sharp : X \rightarrow X_f$. By Definition 4.2.3 this means that $f \in SUC(X)$.

3 \Rightarrow 1: Let $f \in SUC(X)$. Then Lemma 4.6.3 says that X_f is SUC. By Lemma 4.7 we have $X_f \subset UC(G)$. □

5. SUC, HOMOGENEITY AND THE EPIMORPHISM PROBLEM

We say that a G -space X is a *coset G -space* if it is G -isomorphic to the usual coset G -space G/H where H is a closed subgroup of G and G/H is equipped with the quotient topology. We say that a G -space X is *homogeneous* if for every $x, y \in X$ there exists $g \in G$ such that $gx = y$. A homogeneous G -space X is a coset G -space if and only if the orbit map $\tilde{x} : G \rightarrow X$ is open for some (equivalently, every) $x \in X$. Furthermore, $\tilde{x} : G \rightarrow X$ is open iff it is a quotient map. Recall that by a well known result of Effros every homogeneous G -space with Polish G and X is necessarily a coset G -space.

Proposition 5.1. *Let $X = G/H$ be a compact coset G -space.*

- (1) *If X is SUC then X is equicontinuous (that is, almost periodic).*
- (2) *$SUC(X) = AP(X)$.*

Proof. (1): Indeed let $x_0H \in G/H$ and let ε be an element of the uniform structure on the compact space X . By Definition 4.2 we can choose a neighborhood U of e such that

$$(gux_0H, gx_0H) \in \varepsilon \quad \forall (g, u) \in G \times U.$$

By the definition of coset space topology the set $O := Ux_0H$ is a neighborhood of the point x_0H in G/H . We obtain that $(gxH, gx_0H) \in \varepsilon$ whenever $xH \in O$. This proves that x_0H is a point of equicontinuity of $X = G/H$. Hence X is AP.

(2): Every equicontinuous compact G -space is clearly SUC. This implies that always, $SUC(X) \supset AP(X)$. Conversely, let $f \in SUC(X)$. This means that $f = \alpha \circ F$ for a G -compactification $\alpha : X \rightarrow Y$ where Y is SUC and $F \in C(Y) = SUC(Y)$. We can suppose that α is onto because X is compact. Then α is a quotient map. On the other hand X is a coset space G/H . It follows that the natural onto map $G \rightarrow Y$ is also a quotient map. Therefore, Y is also a coset space of G . Now we can apply (1). It follows that Y is almost periodic. Hence f comes from an AP G -factor Y of X . Thus, $f \in AP(X)$. \square

Next we discuss a somewhat unexpected connection between SUC, free topological G -groups and an *epimorphism problem*. Uspenskij has shown in [44] that in the category of Hausdorff topological groups epimorphisms need not have a dense range. This answers a longstanding problem by K. Hofmann. Pestov established [36, 38] that the question completely depends on the *free topological G -groups* $F_G(X)$ of a G -space X in the sense of Megrelishvili [25]. More precisely, the inclusion $i : H \hookrightarrow G$ of topological groups is epimorphism iff the free topological G -group $F_G(X)$ of the coset G -space $X := G/H$ is trivial. Triviality means, ‘as trivial as possible’, isomorphic to the cyclic discrete group.

For a G -space X and points $x_1, x_2 \in X$ we write $x_1 \overset{Aut}{\sim} x_2$ if these two points have the same image under the canonical G -map $X \rightarrow F_G(X)$. If d is a compatible G -invariant metric on a G -space X then $F_G(X)$ coincides with the usual free topological group $F(X)$ of X (see [25, Proposition 3.11]). Therefore, in this case $x_1 \overset{Aut}{\sim} x_2$ iff $x_1 = x_2$.

Theorem 5.2. *Let H be a closed subgroup of G .*

- (1) *If $x_1 \overset{Aut}{\sim} x_2$ for x_1, x_2 in the G -space $X := G/H$ then $x_1 \overset{SUC}{\sim} x_2$.*
- (2) *If the inclusion $H \hookrightarrow G$ is an epimorphism then the coset G -space G/H is SUC-trivial.*

Proof. (1) Assuming the contrary let $f : G/H \rightarrow \mathbb{R}$ be SUC function which separates the points $x_1 := a_1H$ and $x_2 := a_2H$. Then the G -invariant pseudometric

ρ_f on G/H defined by $\rho_f(xH, yH) := \sup_{g \in G} |f(gxH) - f(gyH)|$ also separates these points. We show that ρ_f is continuous. Indeed let $\varepsilon > 0$ and $x_0H \in G/H$. By virtue of Lemma 4.5.4 we can choose a neighborhood U of e such that

$$|f(gux_0H) - f(gx_0H)| < \varepsilon \quad \forall (g, u) \in G \times U.$$

By the definition of coset space topology the set $O := Ux_0H$ is a neighborhood of the point x_0H in G/H . We obtain that $\rho_f(xH, x_0H) < \varepsilon$ whenever $xH \in O$. This proves the continuity of ρ_f .

Consider the associated metric space (Y, d) and the canonical distance preserving onto G -map $X \rightarrow Y, x \mapsto [x]$. The metric d on Y (defined by $d([x], [y]) := \rho_f(x, y)$) is G -invariant. Then $F_G(Y)$ and $F(Y)$ are canonically equivalent (see the discussion above). Since $d([x_1], [x_2]) > 0$ we conclude that x_1 and x_2 have different images in $F_G(X)$. This contradicts the assumption $x_1 \stackrel{Aut}{\sim} x_2$.

(2) Assume that G/H is not SUC-trivial. By Assertion (1) we get that the free topological G -group $F_G(G/H)$ of G/H is not trivial. Therefore by the above mentioned result of Pestov [36] we can conclude that the inclusion $H \hookrightarrow G$ is not an epimorphism. \square

Remark 5.3. (1) The converse to Theorem 5.2.2 is not true (take $G := H_+[0, 1]$, $H := \{e\}$ and apply Theorem 8.3).

(2) As a corollary of Theorem 5.2.2 one can get several examples of SUC-trivial (compact) G -spaces. For example, by [25] the free topological G -group $F_G(X)$ of $X := G/H$ with $G := H(\mathbb{T})$, $H := St(z)$ (where $z \in \mathbb{T}$ is an arbitrary point of the circle \mathbb{T}) is trivial. In fact it is easy to see that the same is true for the smaller group $G := H_+(\mathbb{T})$ (and the subgroup $H := St(z)$) (cf., Proposition 5.1).

(3) It is a well known result by Nummela [35] that if G is a SIN group then the inclusion of a closed proper subgroup $H \hookrightarrow G$ is not an epimorphism. This result easily follows from Theorem 5.2.2. Indeed if G is SIN then by Remark 4.4 for the coset G -space G/H we have $SUC(G/H) = RUC(G/H)$. Hence if G/H is SUC-trivial then necessarily $H = G$ because $RUC(G/H)$ is non-trivial for every closed proper subgroup H of G .

6. REPRESENTATIONS OF GROUPS AND G -SPACES ON BANACH SPACES

For a real normed space V denote by B_V its closed unit ball $\{v \in V : \|v\| \leq 1\}$. Denote by $Iso(V)$ the topological group of all linear surjective isometries $V \rightarrow V$ endowed with the *strong operator topology*. This is just the topology of pointwise convergence inherited from V^V . Let V^* be the dual Banach space of V and

$$\langle \cdot, \cdot \rangle : V \times V^* \rightarrow \mathbb{R}, \quad (v, \psi) \mapsto \langle v, \psi \rangle = \psi(v)$$

is the canonical (always continuous) bilinear mapping.

A *representation* (*co-representation*) of a topological group G on a normed space V is a homomorphism (resp. co-homomorphism) $h : G \rightarrow Iso(V)$. Sometimes it is more convenient to describe a representation (co-representation) by the corresponding left (resp. right) linear isometric actions $\pi_h : G \times V \rightarrow V$, $(g, v) \mapsto gv = h(v)(g)$ (resp., $V \times G \rightarrow V$, $(v, g) \mapsto vg = h(v)(g)$). The (co)representation h is continuous if and only if the action π_h is continuous.

Remark 6.1. Many results formulated for co-representations remain true also for representations (and vice versa) taking into account the following simple fact: for every representation (co-representation) h there exists an associated co-representation (representation) $h^{op} : G \rightarrow Iso(V)$, $g \mapsto h(g^{-1})$.

Definition 6.2. Let $\pi : G \times V \rightarrow V$ be a continuous left action of G on V by linear operators. The *adjoint (or, dual) right action* $\pi^* : V^* \times G \rightarrow V^*$ is defined by $\psi g(v) := \psi(gv)$. The corresponding *adjoint (dual) left action* is $\pi^* : G \times V^* \rightarrow V^*$, where $g\psi(v) := \psi(g^{-1}v)$. Similarly, if $\pi : V \times G \rightarrow V$ is a continuous linear *right action* of G on V (e.g., induced by some co-representation), then the corresponding *adjoint (dual) action* $\pi^* : G \times V^* \rightarrow V^*$ is defined by $g\psi(v) := \psi(vg)$.

The main question considered in [26] was whether the dual action π^* of G on V^* is jointly continuous with respect to the norm topology on V^* . When this is the case we say that the action π (and, also the corresponding representation $h : G \rightarrow Iso(V)$, when π is an action by linear isometries) is *adjoint continuous*. (This name was suggested by V. Uspenskij).

Remark 6.3. In general, not every continuous representation is adjoint continuous (see for example [26]). A standard example is the representation of the circle group $G := \mathbb{T}$ on $V := C(\mathbb{T})$ by translations. Here the Banach space V is separable but with “bad” geometry. The absence of adjoint continuity may happen even for relatively “good” (for instance, for separable Radon-Nikodým) Banach spaces like $V := l_1$. Indeed if we consider the symmetric group $G := S_\infty$, naturally embedded into $Iso(V)$ (endowed with the strong operator topology) as the group of “permutation of coordinates” operators, then the dual action of G on $l_1^* = l_\infty$ is not continuous (see [27]).

It turns out that the situation in that respect is the best possible for the important class $\mathcal{A}sp$ of Asplund Banach spaces. The investigation of this class and the closely related *Radon-Nikodým property* is among the main themes in Banach space theory. Recall that a Banach space V is an *Asplund* space if the dual of every separable linear subspace is separable, iff every bounded subset A of the dual V^* is (weak*,norm)-*fragmented*, iff V^* has the Radon-Nikodým property. Reflexive spaces and spaces of the type $c_0(\Gamma)$ are Asplund. For more

details cf. [6, 10]. For the readers convenience we recall also the definition of fragmentability.

Definition 6.4. (Jane and Rogers [23]) Let (X, τ) be a topological space and ρ be a metric on the set X . Then X is said to be (τ, ρ) -fragmented if for every nonempty $A \subset X$ and every $\varepsilon > 0$ there exists a τ -open subspace O of X such that $O \cap A$ is nonempty and ε -small in (X, ρ) .

Namioka's Joint continuity theorem implies that every weakly compact set in a Banach space is norm fragmented. This explains why every reflexive space is Asplund.

We will use the following result.

Theorem 6.5. [26, Corollary 6.9] *Let V be an Asplund Banach space. If a (not necessarily isometric) linear action $\pi : G \times V \rightarrow V$ is continuous then the dual right action $\pi^* : V^* \times G \rightarrow V^*$ is also continuous.*

Certainly, this result remains true for dual left actions $\pi^* : G \times V^* \rightarrow V^*$, where $g\psi(v) := \psi(g^{-1}v)$, as well as for dual actions defined by a right action $\pi : V \times G \rightarrow V$. The obvious reason is the continuity of the map $G \rightarrow G$, $g \mapsto g^{-1}$.

The following definition provides a flow version of the group representation definitions discussed above. It differs from the usual notion of G -linearization in that here we represent the phase space of the flow as a subset of the dual space V^* (with respect to the dual action and *weak star topology*) rather than as a subset of V .

Definition 6.6. [30] Let X be a G -space. A continuous (*proper*) representation of (G, X) on a Banach space V is a pair

$$(h, \alpha) : G \times X \rightrightarrows Iso(V) \times B^*$$

where $h : G \rightarrow Iso(V)$ is a strongly continuous co-homomorphism and $\alpha : X \rightarrow B^*$ is a weak star continuous G -map (resp. *embedding*) with respect to the *dual action* $G \times V^* \rightarrow V^*$, $(g\varphi)(v) := \varphi(h(g)(v))$. Here B^* is the weak star compact unit ball of the dual space V^* .

Alternatively, one can define a representation in such a way that h is a *homomorphism* and the dual action $G \times V^* \rightarrow V^*$ is defined by $(g\varphi)(v) := \varphi(h(g^{-1})(v))$.

Remark 6.7. Let X be a G -space and let \mathcal{A} be a Banach (closed, unital) subalgebra of $C(X)$. Associated with \mathcal{A} we have the canonical \mathcal{A} -compactification $\nu_{\mathcal{A}} : X \rightarrow X^{\mathcal{A}}$ of X , where $X^{\mathcal{A}} = |\mathcal{A}|$ is the Gelfand space of \mathcal{A} . Here $X^{\mathcal{A}}$ is canonically embedded into the weak star compact unit ball B^* of the dual space \mathcal{A}^* . If \mathcal{A} is G -invariant (that is, the function $(fg)(x) := f(gx)$ lies in \mathcal{A} for every $f \in \mathcal{A}$ and $g \in G$) then $X^{\mathcal{A}}$ admits the natural adjoint action $G \times X^{\mathcal{A}} \rightarrow X^{\mathcal{A}}$ with

the property that all translations $\check{g} : X^{\mathcal{A}} \rightarrow X^{\mathcal{A}}$ are continuous and such that $\alpha_{\mathcal{A}} : X \rightarrow X^{\mathcal{A}} \subset B^*$ is G -equivariant. We obtain in this way a representation (where h is not necessarily continuous)

$$(h, \alpha_{\mathcal{A}}) : (G, X) \rightrightarrows (Iso(\mathcal{A}), B^*)$$

on the Banach space \mathcal{A} , where $h(g)(f) := fg$ (and $\alpha_{\mathcal{A}}(x)(f) := f(x)$). We call it the *canonical* (or, *regular*) \mathcal{A} -*representation* of (G, X) . It is continuous iff $\mathcal{A} \subset RUC(X)$ (see for example [30, Fact 2.2] and [31, Fact 7.2]). The regular $RUC(X)$ -representation leads to the maximal G -compactification $X \rightarrow \beta_G X$ of X . It is proper if and only if X is G -compactifiable.

The following observation due to Teleman is well known (see also [38]).

Fact 6.8. (Teleman [43]) *Every topological group can be embedded into $Iso(V)$ for some Banach space V .*

Proof. It is well known that $RUC(G)$ determines the topology of G . Hence the regular $V := RUC(G)$ -representation $(h, \alpha) : (G, G) \rightrightarrows (Iso(V), B^*)$ is proper. That is, the map α is an embedding. In fact it is easy to see that the cohomomorphism h is an embedding of topological spaces. The representation $h^{op} : G \rightarrow Iso(V), g \mapsto h(g^{-1})$ is then a topological group embedding. \square

Definition 6.9. Let \mathcal{K} be a “well behaved” subclass of the class \mathcal{Ban} of all Banach spaces. Typical and important particular cases for such \mathcal{K} are: \mathcal{Hilb} , \mathcal{Ref} or \mathcal{Asp} , the classes of Hilbert, reflexive or Asplund Banach spaces respectively.

- (1) A topological group G is \mathcal{K} -*representable* if there exists a (co)representation $h : G \rightarrow Iso(V)$ for some $V \in \mathcal{K}$ such that h is topologically faithful (that is, an embedding). Notation: $G \in \mathcal{K}_r$.
- (2) In the opposite direction, we say that G is \mathcal{K} -*trivial* if every continuous \mathcal{K} -representation (or, equivalently, *co-representation*) $h : G \rightarrow Iso(V)$ is trivial.
- (3) We say that a topological group G is *SUC-trivial* if $SUC(G) = \{constants\}$. Analogously can be defined *WAP-trivial* groups. G is WAP-trivial iff G is *reflexively trivial* (\mathcal{Ref} -trivial in the sense of Definition 6.9.2). Similarly, $Asp(G) = \{constants\}$ iff G is \mathcal{Asp} -trivial. These equivalences follow for instance from Theorem 9.3 below.

Remark 6.10. (1) By Teleman’s theorem (Fact 6.8) every topological group is “Banach representable”. Hence, $\{Topological\ Groups\} = \mathcal{Ban}_r$.

- (2) $\{Topological\ Groups\} = \mathcal{Ban}_r \supset \mathcal{Asp}_r \supset \mathcal{Ref}_r \supset \mathcal{Hilb}_r$.
- (3) By Herer and Christensen [22] (see also Banaszczyk [4]) abelian (even monothetic) groups can be \mathcal{Hilb} -trivial. Note also that $c_0 \notin \mathcal{Hilb}_r$, [29].
- (4) [29] The additive group $L_4[0, 1]$ is reflexively but not Hilbert representable.

- (5) [28] $H_+[0, 1] \notin \mathcal{R}ef_r$. It was shown in [28] that every weakly almost periodic function on the topological group $G := H_+[0, 1]$ is constant and that G is $\mathcal{R}ef$ -trivial. By Pestov's observation (see [39, Corollary 1.4] and Lemma 10.2) the same is true for the group $Iso(\mathbb{U}_1)$.
- (6) Theorem 10.3.3 shows that $H_+[0, 1]$ is even $\mathcal{A}sp$ -trivial. In fact we show that every "adjoint continuous" representation of that group is trivial (Theorem 10.3.2). This result was obtained also by Uspenskij (unpublished). Furthermore, we prove a stronger result by showing that $H_+[0, 1]$ (and also $Iso(\mathbb{U}_1)$) are SUC-trivial.

Problem 6.11. (See also [31] and [28])

- (1) Distinguish $\mathcal{A}sp_r$ and $\mathcal{R}ef_r$ by finding $G \in \mathcal{A}sp_r$ such that $G \notin \mathcal{R}ef_r$.
- (2) Find an abelian $G \notin \mathcal{R}ef_r$.

Now we turn to the "well behaved actions". Recall the dynamical versions of Eberlein and Radon-Nikodým compact spaces.

Definition 6.12. [13, 30] Let X be a G -space.

- (1) (G, X) is a *Radon-Nikodým system* (RN for short) if there exists a proper representation of (G, X) on an Asplund Banach space V . If we can choose V to be reflexive, then (G, X) is called an *Eberlein system*. The classes of Radon-Nikodým and Eberlein compact systems will be denoted by RN and Eb respectively.
- (2) (G, X) is called an *RN-approximable system* (RN_{app}) if it can be represented as a subdirect product of RN systems.

Note that compact spaces which are not Eberlein are necessarily non-metrizable, while even for $G := \mathbb{Z}$, there are many natural *metric* compact G -systems which are not RN.

- Definition 6.13.**
- (1) A representation (h, α) of a G -space X on V is *adjoint continuous* if the dual action $G \times V^* \rightarrow V^*$ is also continuous (or, equivalently, if the group corepresentation $h : G \rightarrow Iso(V)$ is adjoint continuous).
 - (2) Denote by Adj the class of compact G -systems which admit a proper adjoint continuous representation on some Banach space V . Theorem 6.5 implies that $\text{RN} \subset Adj$.
 - (3) Denote by $\text{adj}(G)$ the collection of functions on G which come from a compact G -space X such that (G, X) is in the class Adj . In fact this means that f can be represented as a generalized matrix coefficient (see Section 9) of some adjoint continuous representation of G .

Proposition 6.14. $\mathcal{A}sp(G) \subset \text{adj}(G)$ for every topological group G .

Proof. By [30, Theorem 7.11] (or Proposition 7.5) $f \in Asp(G)$ iff f comes from a G -compactification $G \rightarrow X$ of G with $X \in RN$. Now observe (as in Definition 6.13.2) that $RN \subset Adj$ by Theorem 6.5. \square

7. DYNAMICAL COMPLEXITY OF FUNCTIONS

In this section we introduce a hierarchy of dynamical complexity of functions on a topological group G which reflects the complexity of the G -systems from which they come. Our main tool is the cyclic G -system X_f corresponding to a function $f : X \rightarrow \mathbb{R}$. Recall that when $X := G$, the space X_f is the pointwise closure of the orbit Gf in $RUC(G)$. The topological nature of X_f in the Banach space $RUC(G)$ relates to the dynamical complexity of f and leads to a natural hierarchy of complexity (see Theorem 7.12 below). In particular we will examine the role that SUC functions play in this hierarchy.

Periodic orbits and the profinite compactification

The most elementary dynamical system is a finite (periodic) orbit. It corresponds to a clopen subgroup $H < G$ of finite index. These subgroups form a directed set and the corresponding compact inverse limit G -system

$$X^{PF} = \lim_{\leftarrow} G/H$$

is the *profinite* compactification of G .

Almost Periodic functions and the Bohr compactification

The weaker requirement that X_f be norm compact in $RUC(G)$ leads to the well known definition of almost periodicity. A function $f \in C(X)$ on a G -space X is *almost periodic* if the orbit $fG := \{fg\}_{g \in G}$ forms a precompact subset of the Banach space $C(X)$. The collection $AP(X)$ of AP functions is a G -subalgebra in $RUC(X)$. The universal almost periodic compactification of X is the Gelfand space X^{AP} of the algebra $AP(X)$. When X is compact this is the classical *maximal equicontinuous* factor of the system X . A compact G -space X is equicontinuous iff X is almost periodic (AP), that is, iff $C(X) = AP(X)$. For a G -space X the collection $AP(X)$ is the set of all functions which come from equicontinuous (AP) G -compactifications.

For every topological group G , treated as a G -space, the corresponding universal AP compactification is the well known *Bohr compactification* $b : G \rightarrow bG$, where bG is a compact topological group.

Theorem 7.1. *Let X be a G -space. For $f \in RUC(X)$ the following conditions are equivalent:*

- (1) $f \in AP(X)$.
- (2) (G, X_f) is equicontinuous.
- (3) X_f is norm compact in $RUC(G)$.

Proof. (1) \iff (2): $f \in AP(X)$ iff the cyclic algebra \mathcal{A}_f (which, by Proposition 3.1, generates the compactification $X \rightarrow X_f$) is a subalgebra of $AP(X)$.

(2) \iff (3): It is easy to see that the G -space X_f is equicontinuous iff the norm and pointwise topologies coincide on $X_f \subset RUC(G)$. \square

Weakly Almost Periodic functions

A function $f \in C(X)$ on a G -space X is called *weakly almost periodic* (WAP for short; notation: $f \in WAP(X)$) if the orbit $fG := \{fg\}_{g \in G}$ forms a weakly precompact subset of $C(X)$. A compact G -space X is said to be *weakly almost periodic* [8] if $C(X) = WAP(X)$. For a G -space X the collection $WAP(X)$ is the set of all functions which come from WAP G -compactifications. The universal WAP G -compactification $X \rightarrow X^{WAP}$ is well defined. The algebra $WAP(G)$ is a point-universal G -algebra containing $AP(G)$. The compactification $G \rightarrow G^{WAP}$ (for $X := G$) is the universal semitopological semigroup compactification of G .

A compact G -space X is WAP iff it admits sufficiently many representations on reflexive Banach spaces [30]. Furthermore if X is a metric compact G -space then X is WAP iff X admits a proper representation on a reflexive Banach space. That is, iff X is an Eberlein G -space.

Theorem 7.2. *Let X be a G -space. For $f \in RUC(X)$ the following conditions are equivalent:*

- (1) $f \in WAP(X)$.
- (2) (G, X_f) is WAP.
- (3) X_f is weak compact in $RUC(G)$.
- (4) (G, X_f) is Eberlein (i.e., reflexively representable).

Proof. (1) \iff (2): $f \in WAP(X)$ iff the algebra \mathcal{A}_f is a subalgebra of $WAP(X)$.

(2) \implies (3): Let $F_e : X_f \rightarrow \mathbb{R}$, $F_e(\omega) = \omega(e)$ be as in the definition of X_f . Consider the weak closure $Y := \text{cl}_w(F_e G)$ of the orbit $F_e G$. Then Y is weakly compact in $C(X_f)$ because $F_e \in C(X_f) = WAP(X_f)$ is weakly almost periodic. If ω_1 and ω_2 are distinct elements of X_f then $(F_e g)(\omega_1) = \omega_1(g) \neq \omega_2(g) = (F_e g)(\omega_2)$ for some $g \in G$. This means that the separately continuous evaluation map $Y \times X_f \rightarrow \mathbb{R}$ separates points of X_f . Now X_f can be treated as a pointwise compact bounded subset in $C(Y)$. Hence by Grothendieck's well known theorem [21] we get that X_f is weakly compact in $C(Y)$. Since $G \rightarrow Y$, $g \mapsto gF_e$ is a G -compactification of G , we have a natural embedding of Banach algebras

$j : C(Y) \hookrightarrow RUC(G)$. It follows that $X_f = j(X_f)$ is also weakly compact as a subset of $RUC(G)$.

(3) \implies (4): The isometric action $G \times RUC(G) \rightarrow RUC(G)$, $(g, f) \mapsto gf$ induces a representation $h : G \rightarrow Iso(RUC(G))$. If the G -subset X_f is weakly compact in $RUC(G)$ then one can apply Theorem 4.11 (namely, the equivalence between (i) and (ii)) of [30] which guarantees that the G -space X_f is Eberlein.

(4) \implies (1): $f \in WAP(X)$ because it comes from (G, X_f) (Proposition 3.1) which is WAP (being reflexively representable). \square

Asplund functions, “sensitivity to initial conditions” and Banach representations

The following definition of “sensitivity to initial conditions” is essential in several definitions of chaos in dynamical systems, mostly for $G := \mathbb{Z}$ or \mathbb{R} actions on metric spaces (see for instance papers of Guckenheimer, Auslander and Yorke, Devaney, Glasner and Weiss).

Definition 7.3. [13] Let (X, μ) be a uniform G -space.

- (1) We say that X is *sensitive to initial conditions* (or just *sensitive*) if there exists an $\varepsilon \in \mu$ such that for every nonempty open subset O of X the set gO is not ε -small for some $g \in G$. Otherwise, X is *non-sensitive* (for short: NS).
- (2) X is *Hereditarily Non Sensitive* (HNS) if every closed G -subspace of X is NS.

Denote by HNS the class of all compact HNS systems. The following result says that a compact G -system X is HNS iff (G, X) admits sufficiently many representations on Asplund spaces.

Theorem 7.4. [13]

- (1) $HNS = RN_{app}$.
- (2) If X is a compact metric G -space then X is HNS iff X is RN (that is, Asplund representable).

A function $f : X \rightarrow \mathbb{R}$ on a G -space X is *Asplund* (notation: $f \in Asp(X)$) [30] if it satisfies one of the following equivalent conditions.

Proposition 7.5. Let $f : X \rightarrow \mathbb{R}$ be a function on a G -space X . The following conditions are equivalent:

- (1) f comes from a G -compactification $\nu : X \rightarrow Y$ where (G, Y) is HNS.
- (2) f comes from a G -compactification $\nu : X \rightarrow Y$ where (G, Y) is RN.

- (3) f comes from a G -compactification $\nu : X \rightarrow Y$ and a function $F : Y \rightarrow \mathbb{R}$ where the pseudometric space $(Y, \rho_{H,F})$ with

$$\rho_{H,F}(x, x') = \sup_{h \in H} |F(hx) - F(hx')|.$$

is separable for every countable (equivalently, second countable) subgroup $H \subset G$.

The collection $Asp(X)$ is always a G -subalgebra of $RUC(X)$. It defines the maximal HNS-compactification $X \rightarrow X^{Asp} = |Asp(X)|$ of X . For every topological group G the algebra $Asp(G)$ (as usual, $X := G$ is a left G -space) is point-universal.

Theorem 7.6. *Let X be a G -space. For every $f \in RUC(X)$ the following conditions are equivalent:*

- (1) $f \in Asp(X)$.
- (2) (G, X_f) is RN.
- (3) X_f is norm fragmented in $RUC(G)$.

Proof. If X is compact then the proof follows directly from [13, Theorem 9.12]. Now observe that one can reduce the case of an arbitrary G -space X to the case of a compact G -space X_f by considering the cyclic G -system $(X_f)_{F_e}$ (defined for $X := X_f$ and $f := F_e$) which can be naturally identified with X_f . \square

Explicitly the fragmentability of X_f means that for every $\varepsilon > 0$ every nonempty (closed) subset A of $X_f \subset \mathbb{R}^G$ contains a relatively open (in the pointwise topology) nonempty subset $O \cap A$ which is ε -small in the Banach space $RUC(G)$.

As we already mentioned every weakly compact set is norm fragmented so that $WAP(X) \subset Asp(X)$ for every G -space X . In particular, $WAP(G) \subset Asp(G)$.

Locally equicontinuous functions

During the last decade various conditions weakening the classical notion of equicontinuity were introduced and studied (see e.g. [16], [1], [2], [3]). The following definition first appears in a paper of Glasner and Weiss [16].

Definition 7.7. [16] Let (X, μ) be a uniform G -space. A point $x_0 \in X$ is a point of *local equicontinuity* (notation: $x_0 \in LE_X$) if x_0 is a point of equicontinuity in the uniform G -subspace $\text{cl}(Gx_0)$. We have $x_0 \in LE_X$ iff $x_0 \in LE_Y$ iff $x_0 \in Eq_Y$ where Y is the orbit Gx_0 of x_0 (see Lemma 7.8.1). If $LE_X = X$, then X is *locally equicontinuous* (LE).

Lemma 7.8. (1) *Let Y be a dense G -subspace of (X, μ) and $y_0 \in Y$. Then $y_0 \in Eq_X$ if and only if $y_0 \in Eq_Y$.*
 (2) $SUC_X \supset LE_X \supset Eq_X$.

(3) $SUC(X) \supset LE(X)$.

Proof. (1) Let $\varepsilon \in \mu$. There exists $\delta \in \mu$ such that δ is a closed subset of $X \times X$ and $\delta \subset \varepsilon$. If $y_0 \in Eq_Y$ there exists an open set U in X such that $y_0 \in U$ and $(gy, gy_0) \in \delta$ for all $y \in U \cap Y$ and $g \in G$. Since Y is dense in X and U is open we have $U \subset \text{cl}(U \cap Y)$. Since every g -translation $X \rightarrow X$, $x \mapsto gx$ is continuous and δ is closed we get $(gx, gy_0) \in \delta \subset \varepsilon$ for every $g \in G$ and $x \in U$.

(2): Let $x_0 \in LE_X$. For every $\varepsilon \in \mu$ there exists a neighborhood $O(x_0)$ such that (gx, gx_0) is ε -small for every $x \in O(x_0)$ and $g \in G$. Choose a neighborhood $U(e)$ such that $Ux_0 \subset O$. Then (gux_0, gx_0) is ε -small, too. This proves the non-trivial part $SUC_X \supset LE_X$.

(3): Directly follows from (2). □

The collection $LE(X)$ forms a G -subalgebra of $RUC(X)$. Always, $Asp(X) \subset LE(X)$. The algebra $LE(X)$ defines the maximal LE-compactification $X \rightarrow X^{LE}$ of X . For every topological group G the algebra $LE(G)$ is point-universal [13].

Theorem 7.9. [13] *Let X be a G -space. For $f \in RUC(X)$ the following conditions are equivalent:*

- (1) $f \in LE(X)$.
- (2) (G, X_f) is LE.
- (3) X_f is orbitwise light in $RUC(G)$ (that is, for every function $\psi \in X_f$ the pointwise and norm topologies coincide on the orbit $G\psi$).

Proof. (1) \iff (2) Directly follows from [13, Theorem 5.15.1]. On the other hand by [13, Lemma 5.18] we have (2) \iff (3). □

The dynamical hierarchy

Theorem 7.10. (1) *Let X be a (not necessarily compact) G -space. We have the following chain of inclusions of G -subalgebras:*

$$RUC(X) \supset SUC(X) \supset LE(X) \supset Asp(X) \supset WAP(X) \supset AP(X)$$

and the corresponding chain of G -factor maps

$$\beta_G X \rightarrow X^{SUC} \rightarrow X^{LE} \rightarrow X^{Asp} \rightarrow X^{WAP} \rightarrow X^{AP}.$$

- (2) *For every topological group G we have the following chain of inclusions of G -subalgebras:*

$$RUC(G) \supset UC(G) \supset SUC(G) \supset LE(G) \supset Asp(G) \supset WAP(G) \supset AP(G)$$

and the corresponding chain of G -factor maps

$$G^{RUC} \rightarrow G^{UC} \rightarrow G^{SUC} \rightarrow G^{LE} \rightarrow G^{Asp} \rightarrow G^{WAP} \rightarrow G^{AP}.$$

Proof. For the assertions concerning $SUC(X)$ and $SUC(G)$ see Lemmas 4.3 and 7.8. For the other assertions see [13]. \square

Remark 7.11. The compactifications G^{AP} and G^{WAP} of G are respectively a *topological group* and a *semitopological semigroup*. The compactifications G^{RUC} and G^{Asp} are right topological semigroup compactifications of G (see [13]). The same is true for the compactification $j : G \rightarrow G^{SUC}$ (Lemma 4.6.5). Below (Theorem 10.3.5) we show that the *Roelcke compactification* $i : G \hookrightarrow G^{UC}$ (which is always proper by Lemma 2.1) is not in general a right topological semigroup compactification. That is, $UC(G)$ is not in general point-universal.

We sum up our results in the following dynamical hierarchy theorem where we list dynamical properties of $f \in RUC(X)$ and the corresponding topological properties of $X_f \subset RUC(G)$ (cf. [13, Remark 9.13]).

Theorem 7.12. *For every G -space X and a function $f \in RUC(X)$ we have*

$$\begin{aligned} X_f \text{ is norm compact} &\iff f \text{ is AP} \\ X_f \text{ is weakly compact} &\iff f \text{ is WAP} \\ X_f \text{ is norm fragmented} &\iff f \text{ is Asplund} \\ X_f \text{ is orbitwise light} &\iff f \text{ is LE} \\ X_f \subset UC(G) &\iff f \text{ is SUC} \end{aligned}$$

Example 7.13. Let G be the unitary group $U(H) = Iso(H)$ where H is an infinite dimensional Hilbert space. Then $UC(G) = SUC(G) = LE(G) = Asp(G) = WAP(G)$. Indeed the completion of $(G, \mathcal{L} \wedge \mathcal{R})$ can be identified with the compact semitopological semigroup $\Theta(H)$ of all nonexpansive linear operators (Uspenskij [46]). It follows that G^{UC} can be identified with $\Theta(H)$. The latter is a reflexively representable G -space (see for example [30, Fact 5.2]). Therefore $UC(G) \subset WAP(G)$. The reverse inclusion is well known (see for instance Theorem 7.10). Hence, $UC(G) = WAP(G)$.

Let $\mathcal{C}_u = \{f \in UC(G) : X_f \subset UC(G)\}$. The collection of functions \mathcal{C}_u was studied by Veech in [49]. He notes there that $WAP(G) \subset \mathcal{C}_u$ and proves the following theorem.

Theorem 7.14. (Veech [49, Proposition 5.4]) *Let G be a semisimple analytic Lie group with finite center and without compact factors. If $f \in \mathcal{C}_u$ then every limit point of Gf in X_f ; i.e. any function of the form $h(g) = \lim_{g_n \rightarrow \infty} f(gg_n)$, is a constant function.*

(A sequence $g_n \in G$ “tends to ∞ ” if each of its “projections” onto the simple components of G tends to ∞ in the usual sense.) He then deduces the fact that

for G which is a direct product of simple groups the algebra $WAP(G)$ coincides with the algebra \mathcal{W}^* of continuous functions on G which “extend continuously” to the product of the one-point compactification of the simple components of G ([49, Theorem 1.2]). By our Proposition 4.12, $\mathcal{C}_u = SUC(G)$. Taking this equality into account, Veech’s theorem implies now the following result.

Corollary 7.15. *For every simple noncompact connected Lie group G with finite center (e.g., $SL_n(\mathbb{R})$) we have $SUC(G) = WAP(G) = \mathcal{W}^*$. In particular the corresponding universal SUC (and hence WAP) compactification is equivalent to the one point compactification of G .*

8. THE GROUP $H_+[0, 1]$

Consider the Polish topological group $G := H_+[0, 1]$ of all orientation preserving homeomorphisms of the closed unit interval, endowed with the compact open topology. Here is a list of some selected known results about this group:

- (1) G is topologically simple.
- (2) G is not Weil-complete; that is, the right uniform structure \mathcal{R} of G is not complete. The completion of the uniform space (G, \mathcal{R}) can be identified with the semigroup of all continuous, nondecreasing and surjective maps $[0, 1] \rightarrow [0, 1]$ endowed with the uniform structure of uniform convergence (Roelcke-Dierolf [41, p. 191]).
- (3) G is *Roelcke precompact* (that is the Roelcke uniformity $\mathcal{L} \wedge \mathcal{R}$ on G is precompact) [41].
- (4) The completion of $(G, \mathcal{L} \wedge \mathcal{R})$ can be identified with the curves that connect the points $(0, 0)$ and $(1, 1)$ and “never go down” (Uspenskij [48], see Lemma 8.4 below).
- (5) Every weakly almost periodic function on G is constant and every continuous representation $G \rightarrow Iso(V)$, where V is a reflexive Banach space, is trivial (Megrelishvili [28]).
- (6) G is *extremely amenable*; that is every compact Hausdorff G -space has a fixed point property (Pestov [37]).

We are going to show that $H_+[0, 1]$ is SUC-trivial and hence also $\mathcal{A}sp$ -trivial. Since every reflexive Banach space is Asplund these results strengthen the main results of [28] (results mentioned in item (5) above).

Definition 8.1. Let (X, μ) be a compact G -space. We say that two points $a, b \in X$ are *SUC-proximal* if there exist nets s_i and g_i in G and a point $x_0 \in X$ such that s_i converges to the neutral element e of G , the net $g_i x_0$ converges to a and the net $g_i s_i x_0$ converges to b .

Lemma 8.2. *If the points a and b are SUC-proximal in a G -space X then $a \overset{SUC}{\sim} b$.*

Proof. A straightforward consequence of our definitions using Lemma 4.5.4. \square

Theorem 8.3. *Let $G = H_+[0, 1]$ be the topological group of orientation-preserving homeomorphisms of $[0, 1]$ endowed with the compact open topology. Then G is SUC-trivial.*

Proof. Denote by $j : G \rightarrow G^{SUC}$ and $i : G \rightarrow G^{UC}$ the G -compactifications (i necessarily is proper by Lemma 2.1) induced by the Banach G -algebras $SUC(G) \subset UC(G)$. There exists a canonical onto G -map $\pi : G^{UC} \rightarrow G^{SUC}$ such that the following diagram of G -maps is commutative:

$$\begin{array}{ccc} G & \xrightarrow{i} & G^{UC} \\ & \searrow j & \downarrow \pi \\ & & G^{SUC} \end{array}$$

We have to show that G^{SUC} is trivial for $G = H_+[0, 1]$. One of the main tools for the proof is the following identification.


Lemma 8.4. [48, Uspenskiĭ] *The dynamical system G^{UC} is isomorphic to the G -space (G, Ω) . Here Ω denotes the compact space of all curves in $[0, 1] \times [0, 1]$ which connect the points $(0, 0)$ and $(1, 1)$ and “never go down”, equipped with the Hausdorff metric. These are the relations $\omega \subset [0, 1] \times [0, 1]$ where for each $t \in [0, 1]$, $\omega(t)$ is either a point or a vertical closed segment. The natural action of $G = H_+[0, 1]$ on Ω is $(g\omega)(t) = g(\omega(t))$ (by composition of relations on $[0, 1]$).*

We first note that every “zig-zag curve” (i.e. a curve z which consists of a finite number of horizontal and vertical pieces) is an element of Ω . In particular the curves γ_c with exactly one vertical segment defined as $\gamma_c(t) = 0$ for every $t \in [0, c)$, $\gamma_c(c) = \{c\} \times [0, 1]$ and $\gamma_c(t) = 1$ for every $t \in (c, 1]$, are elements of $\Omega = G^{UC}$. Note that the curve γ_1 is a fixed point for the left G action. We let $\theta = \pi(\gamma_1)$ be its image in G^{SUC} . Of course θ is a fixed point in G^{SUC} . We will show that $\theta = j(e)$ and since the G -orbit of $j(e)$ is dense in G^{SUC} this will show that G^{SUC} is a singleton.


The idea is to show that zig-zag curves are SUC-proximal in G^{UC} . Then Lemma 8.2 will ensure that their images in G^{SUC} coincide. Choosing a sequence z_n of zig-zag curves which converges in the Hausdorff metric to $i(e)$ in G^{UC} we will have $\pi(z_n) = \pi(\gamma_1) = \theta$ for each n . This will imply that indeed $j(e) = \pi(i(e)) = \pi(\lim_{n \rightarrow \infty} z_n) = \lim_{n \rightarrow \infty} \pi(z_n) = \theta$.

First we show that $\pi(\gamma_1) = \pi(\gamma_c)$ for any $0 < c < 1$. As indicated above, since G^{SUC} is the Gelfand space of the algebra $SUC(G)$, by Lemma 8.2, it suffices to show that the pair γ_1, γ_c is SUC-proximal in G^{UC} . Since $X^{SUC} = X$ for $X := G^{SUC}$ we conclude that $\pi(\gamma_1) = \pi(\gamma_c)$.

Let $p \in G^{UC}$ be the zig-zag curve defined by $p(t) = t$ in the interval $[0, c]$ and by $p(t) = c$ for every $t \in [c, 1]$. Pick a sequence s_n of elements in G such that s_n converges to e and $s_n c < c$. It is easy to choose a sequence g_n in G such that $g_n s_n c$ converges to 0 and $g_n c$ converges to 1. Then the sequences s_n and g_n are the desired sequences; that is, $g_n p \rightarrow \gamma_c$, $g_n s_n p \rightarrow \gamma_1$ (see the picture below).



Denote $\theta = \pi(\gamma_1) = \pi(\gamma_c)$. Using similar arguments (see the picture below, where $a \overset{\pi}{\sim} b$ means $\pi(a) = \pi(b)$) construct a sequence $z_n \in G^{UC}$ of zig-zag curves which converges to $i(e)$ and such that $\pi(z_n) = \theta$ for every n .



In view of the discussion above this construction completes the proof of the theorem. \square

9. MATRIX COEFFICIENT CHARACTERIZATION OF SUC AND LE

Definition 9.1. Let $h : G \rightarrow Iso(V)$ be a co-representation of G on a normed space V and let

$$V \times G \rightarrow V, \quad (v, g) \mapsto vg := h(g)(v)$$

be the corresponding right action. For a pair of vectors $v \in V$ and $\psi \in V^*$ the associated *matrix coefficient* is defined by

$$m_{v,\psi} : G \rightarrow \mathbb{R}, \quad g \mapsto \psi(vg) = \langle vg, \psi \rangle = \langle v, g\psi \rangle.$$

If $h : G \rightarrow Iso(V)$ is a representation then the matrix coefficient $m_{v,\psi}$ is defined similarly by

$$m_{v,\psi} : G \rightarrow \mathbb{R}, \quad g \mapsto \psi(gv) = \langle gv, \psi \rangle = \langle v, \psi g \rangle.$$

For example, if $V = H$ is a Hilbert space then $f = m_{u,\psi}$ is the *Fourier-Stieltjes transform*. In particular, for $u = \psi$ we get the *positive definite functions*.

We say that a vector $v \in V$ is *G-continuous* if the corresponding orbit map $\tilde{v} : G \rightarrow V$, $\tilde{v}(g) = vg$, defined through $h : G \rightarrow Iso(V)$, is norm continuous. The *continuous G-vector* $\psi \in V^*$ are defined similarly with respect to the dual action.

Lemma 9.2. [30] *Let $h : G \rightarrow Iso(V)$ be a co-representation of G on V . If ψ (resp.: $v \in V$) is norm G -continuous, then $m_{v,\psi}$ is left (resp.: right) uniformly continuous on G . Hence, if v and ψ are both G -continuous then $m_{v,\psi} \in UC(G)$.*

In the next theorem we list characterizations of several subalgebras of $RUC(G)$ in terms of matrix coefficients. These characterizations also provide an alternative way to establish the inclusions in Theorem 7.10.2 for $X := G$.

Theorem 9.3. *Let G be a topological group and $f \in C(G)$.*

- (1) $f \in RUC(G)$ iff $f = m_{v,\psi}$ for some continuous co-representation $h : G \rightarrow Iso(V)$, where $V \in Ban$.
- (2) $f \in UC(G)$ iff $f = m_{v,\psi}$ for some co-representation where v and ψ are both G -continuous iff $f = m_{v,\psi}$ for some continuous co-representation where ψ is G -continuous.
- (3) $f \in SUC(G)$ iff $f = m_{v,\psi}$ for some continuous co-representation $h : G \rightarrow Iso(V)$, where φ is norm G -continuous in V^* for every φ from the weak star closure $cl_{w^*}(G\psi)$.
- (4) $f \in adj(G)$ iff $f = m_{v,\psi}$ for some adjoint continuous co-representation.
- (5) $f \in LE(G)$ iff $f = m_{v,\psi}$ for some continuous co-representation $h : G \rightarrow Iso(V)$, where weak star and norm topologies coincide on each orbit $G\varphi$ where φ belongs to the weak star closure $Y := cl_{w^*}(G\psi)$.
- (6) $f \in Asp(G)$ iff f is a matrix coefficient of some continuous Asplund co-representation of G .
- (7) $f \in WAP(G)$ iff f is a matrix coefficient of some continuous reflexive co-representation (or, representation) of G .

Proof. Claim (1) follows by taking in the regular $RUC(G)$ -corepresentation $V := RUC(G)$, $v := f$ and $\psi := \alpha(e)$. Claim (4) is a reformulation of Definition 6.13.3.

The remaining assertions are essentially nontrivial. Their proofs are based on an equivariant generalization of the Davis-Figiel-Johnson-Pelczyński interpolation technic [7]. For detailed proofs of (6) and (7) (for co-representations) see

[30, Theorem 7.17] and [30, Theorem 5.1]. As to the “representations case” in (7) observe that a matrix coefficient of a (continuous) co-representation on a reflexive space V can be treated as matrix coefficient of a (continuous) representation on the dual space V^* . The continuity of the dual action follows by Theorem 6.5.

For (2) see [32]. Below we provide the proof of the new assertions (3) and (5). See Theorems 9.8 and 9.10 respectively. \square

Remark 9.4. Let $\mathcal{A} \subset RUC(G)$ be a point-universal G -subalgebra. Then \mathcal{A} is *left m -introverted* in the sense of [33], [5, Definition 1.4.11]. Indeed, we have only to check that every matrix coefficient $m_{v,\psi}$ of the regular \mathcal{A} -representation of the action (G, G) on the Banach space \mathcal{A} belongs again to \mathcal{A} whenever $v \in \mathcal{A}$ and $\psi \in |\mathcal{A}| \subset \mathcal{A}^*$. Let $X := |\mathcal{A}|$. Then the G -system $(X, \text{eva}(e))$ is point universal and $\mathcal{A} = \mathcal{A}(X, \text{eva}(e))$. The matrix coefficient $m_{v,\psi}$ comes from the subsystem $(\text{cl}(G\psi), \psi)$. In other words $m_{v,\psi} \in \mathcal{A}(\text{cl}(G\psi), \psi)$. By Lemma 3.3 we have $\mathcal{A}(\text{cl}(G\psi), \psi) \subset \mathcal{A}(X, \text{eva}(e))$. Thus $m_{v,\psi} \in \mathcal{A}(X, \text{eva}(e)) = \mathcal{A}$.

Definition 9.5. (1) Let G be a topological group and $G \times X \rightarrow X$ and $Y \times G \rightarrow Y$ be respectively left and right actions. A map $\langle \cdot, \cdot \rangle : Y \times X \rightarrow \mathbb{R}$ is *G -compatible* if

$$\langle yg, x \rangle = \langle y, gx \rangle \quad \forall (y, g, x) \in Y \times G \times X.$$

(2) We say that a subset $M \subset Y$ is *SUC-small at $x_0 \in X$* if for every $\varepsilon > 0$ there exists a neighborhood U of e such that

$$|\langle v, ux_0 \rangle - \langle v, x_0 \rangle| \leq \varepsilon \quad \forall (v, u) \in M \times U.$$

If M is SUC-small at every $x \in X$ then we say that M is *SUC-small for X* .

(3) Let $h : G \rightarrow Iso(V)$ be a continuous co-representation on a normed space V and $h^* : G \rightarrow Iso(V^*)$ be the dual representation. Then we say that $M \subset V$ is *SUC-small at $x_0 \in X \subset V^*$* if this happens in the sense of (2) regarding the canonical bilinear G -compatible map $\langle \cdot, \cdot \rangle : V \times V^* \rightarrow \mathbb{R}$.

For example, a vector $\psi \in V^*$ in the dual space V^* is G -continuous iff the unit ball B_V of V is SUC-small at ψ (see Lemma 9.7.3).

We give some useful properties of SUC-smallness.

Lemma 9.6. (1) Let $Y_1 \times X_1 \rightarrow \mathbb{R}$ and $Y_2 \times X_2 \rightarrow \mathbb{R}$ be two G -compatible maps. Suppose that $\gamma_1 : X_1 \rightarrow X_2$ and $\gamma_2 : Y_2 \rightarrow Y_1$ are G -maps such that

$$\langle y, \gamma_1(x) \rangle = \langle \gamma_2(y), x \rangle \quad \forall (y, x) \in Y_2 \times X_1.$$

Then for every nonempty subset $M \subset Y_2$ the subset $\gamma_1(M) \subset Y_1$ is SUC-small at $x \in X_1$ if and only if M is SUC-small at $\gamma_1(x) \in X_2$.

(2) Let X be a (not necessarily compact) G -space. If $f \in SUC(X)$ then:

(a) fG is SUC-small for X with respect to the G -compatible evaluation map

$$fG \times X \rightarrow \mathbb{R}, \quad (fg, x) \mapsto f(gx).$$

(b) The subset $F_e G$ of $C(X_f)$ is SUC-small for X_f considered as a subset of V^* where $V := C(X_f)$ (with respect to the canonical map $V \times V^* \rightarrow \mathbb{R}$ and the natural co-representation $G \rightarrow \text{Iso}(V)$).

Proof. (1): Observe that for every triple $(m, u, x_0) \in M \times G \times X$ we have

$$\langle \gamma_1(m), ux_0 \rangle - \langle \gamma_1(m), x_0 \rangle = \langle m, \gamma_2(ux_0) \rangle - \langle m, \gamma_2(x_0) \rangle = \langle m, u\gamma_2(x_0) \rangle - \langle m, \gamma_2(x_0) \rangle.$$

(2): (a) Directly follows by Lemma 4.5.4.

(b): Let $f \in \text{SUC}(X)$. Then it comes by Proposition 3.1.3 from a compact G -system X_f and the G -compactification $f_{\sharp} : X \rightarrow X_f$. As we know there exists $F_e \in C(X_f)$ such that $f = F_e \circ f_{\sharp}$. Theorem 4.12 implies that X_f is SUC and $F_e \in \text{SUC}(X_f)$. By claim (a) it follows that $F_e G$ is SUC-small for X_f . The G -compatible map $F_e G \times X_f \rightarrow \mathbb{R}$ can be treated as a restriction of the canonical form $V \times V^* \rightarrow \mathbb{R}$, where $V := C(X_f)$ (considered X_f as a subset of $C(X_f)^*$). \square

Lemma 9.7. *Let $h : G \rightarrow \text{Iso}(V)$ be a continuous co-representation.*

- (1) *For every subset X of V^* the family of SUC-small sets for X in V is closed under taking subsets, norm closures, finite linear combinations, finite unions and convex hulls.*
- (2) *If $M_n \subset V$ is SUC-small at $x_0 \in V^*$ for every $n \in \mathbb{N}$ then so is the set $\bigcap_{n \in \mathbb{N}} (M_n + \delta_n B_V)$ for every positive decreasing sequence $\delta_n \rightarrow 0$.*
- (3) *For every $\psi \in V^*$ the following are equivalent:*
 - (i) *The orbit map $\tilde{\psi} : G \rightarrow V^*$ is norm continuous.*
 - (ii) *\mathcal{B} is SUC-small at ψ , where $\mathcal{B} := \{\check{v} : V^* \rightarrow \mathbb{R}, x \mapsto \check{v}(x) := \langle v, x \rangle\}_{v \in B_V}$.*

Proof. Assertion (1) is straightforward.

(2): We have to show that the set $\bigcap_{n \in \mathbb{N}} (M_n + \delta_n B_V)$ is SUC-small at x_0 . Let $\varepsilon > 0$ be fixed. Since Gx_0 is a bounded subset of V^* one can choose $n_0 \in \mathbb{N}$ such that $|v(gx_0)| < \frac{\varepsilon}{4}$ for every $g \in G$ and every $v \in \delta_{n_0} B_V$. Since M_{n_0} is SUC-small at x_0 we can choose a neighborhood $U(e)$ such that $|m(ux_0) - m(x_0)| < \frac{\varepsilon}{2}$ for every $u \in U$ and every $m \in M_{n_0}$. Now every element $w \in \bigcap_{n \in \mathbb{N}} (M_n + \delta_n B_V)$ has a form $w = m + v$ for some $m \in M_{n_0}$ and $v \in \delta_{n_0} B_V$. Then for every $u \in U$ we have

$$|w(ux_0) - w(x_0)| \leq |m(ux_0) - m(x_0)| + |v(ux_0)| + |v(x_0)| < \frac{\varepsilon}{2} + \frac{\varepsilon}{4} + \frac{\varepsilon}{4} = \varepsilon.$$

(3): Use that $\|u\psi - \psi\| = \sup_{v \in B_V} |\langle v, u\psi \rangle - \langle v, \psi \rangle|$ and B_V is G -invariant. \square

Theorem 9.8. *The following conditions are equivalent:*

- (1) $f \in \text{SUC}(G)$.

- (2) $f = m_{v,\psi}$ for some continuous Banach co-representation $h : G \rightarrow Iso(V)$, $v \in V$ and $\psi \in V^*$, with the property that φ is norm G -continuous for every φ in the weak $*$ closure $cl_{w^*}(G\psi)$.

Moreover, one can assume in (2) that $cl_{w^*}(G\psi)$ separates points of V .

Proof. (2) \implies (1): Let $h : G \rightarrow Iso(V)$ be a continuous co-homomorphism such that f is a matrix coefficient of h . That is, we can choose $v \in V$ and $\psi \in V^*$ such that $f(g) = \langle vg, \psi \rangle = \langle v, g\psi \rangle$ for every $g \in G$. One can assume that $\|\psi\| = 1$. The strong continuity of h ensures that the dual restricted (left) action of G on the weak star compact unit ball (V_1^*, w^*) is jointly continuous. Consider the orbit closure $X := cl_{w^*}(G\psi)$ in the compact G -space (V_1^*, w^*) . Define the continuous function $\hat{v} : X \rightarrow \mathbb{R}$ induced by the vector v . Precisely, $\hat{v}(x) = \langle v, x \rangle$ and in particular, $f(g) = \hat{v}(g\psi)$. So f comes from X and the compactification $\nu : G \rightarrow X$, $g \mapsto g\psi$. It suffices to show that the G -system X is SUC. Let x_0 be an arbitrary point in X and let w be an arbitrary vector in V . By the definition of the weak star topology and the corresponding uniformity on the compact space X it suffices to show that for every $\varepsilon > 0$ there exists a neighborhood $U(e)$ such that $|\hat{w}(gux_0) - \hat{w}(gx_0)| \leq \varepsilon$ for every $g \in G$ and every $u \in U$. By simple computations we get

$$\begin{aligned} |\hat{w}(gux_0) - \hat{w}(gx_0)| &= |\langle w, gux_0 \rangle - \langle w, gx_0 \rangle| = |\langle wg, ux_0 \rangle - \langle wg, x_0 \rangle| \\ &= |\langle wg, ux_0 - x_0 \rangle| \leq \|wg\| \cdot \|ux_0 - x_0\|. \end{aligned}$$

Take into account that $\|wg\| = \|w\|$. Since $x_0 \in cl(G\psi)$, by our assumption the orbit map $\tilde{x}_0 : G \rightarrow V^*$ is norm continuous with respect to the dual action of G on V^* . Therefore, given $\varepsilon > 0$, there exists a neighborhood U of e in G such that $\|ux_0 - x_0\| < \|w\|^{-1}\varepsilon$ for every $u \in U$. Thus, $|\hat{w}(gux_0) - \hat{w}(gx_0)| \leq \varepsilon$. This shows that X is SUC. Hence, $f \in SUC(G)$.

(1) \implies (2): Let $f \in SUC(X)$. Then it comes by Proposition 3.1.3 from a compact transitive G -system X_f and the G -compactification $f_{\sharp} : G \rightarrow X_f$. There exists $F := F_e \in C(X_f)$ such that $f = F \circ f_{\sharp}$. By Lemma 9.6.2 we conclude that FG is SUC-small for $X_f \subset C(X_f)^*$.

Let $M := co(-FG \cup FG)$ be the convex hull of the symmetric set $-FG \cup FG$. Then M is a convex symmetric bounded G -invariant subset in $C(X_f)$. By Lemma 9.7.1 we know that M is also SUC-small for X_f .

For brevity of notation let E denote the Banach space $C(X_f)$. Since X_f is a compact G -space the natural right action of G on $E = C(X_f)$ (by linear isometries) is continuous.

Consider the sequence $K_n := 2^n M + 2^{-n} B_E$, where B_E is the unit ball of E . Since M is convex and symmetric, we can apply the construction of [7] (we mostly use the presentation and the development given by Fabian in the book [10]). Let

$\|\cdot\|_n$ be the Minkowski's functional of the set K_n . That is,

$$\|v\|_n = \inf \{ \lambda > 0 \mid v \in \lambda K_n \}$$

Then $\|\cdot\|_n$ is a norm on E equivalent to the given norm of E for every $n \in \mathbb{N}$. For $v \in E$, let

$$N(v) := \left(\sum_{n=1}^{\infty} \|v\|_n^2 \right)^{1/2} \quad \text{and} \quad V := \{v \in E \mid N(v) < \infty\}$$

Denote by $j : V \hookrightarrow E$ the inclusion map. Then (V, N) is a Banach space, $j : V \rightarrow E$ is a continuous linear injection and

$$M \subset j(B_V) = B_V$$

Indeed, if $v \in M$ then $2^n v \in K_n$. Therefore, $\|v\|_n \leq 2^{-n}$ and $N(v)^2 \leq \sum_{n \in \mathbb{N}} 2^{-2n} < 1$.

By our construction M and B_E are G -invariant. This implies that the natural right action $V \times G \rightarrow V$, $(v, g) \mapsto vg$ is isometric, that is $N(vg) = N(v)$. Moreover, by the definition of the norm N on V (use the fact that the norm $\|\cdot\|_n$ on E is equivalent to the given norm of E for each $n \in \mathbb{N}$) we can show that this action is norm continuous. Therefore, the co-representation $h : G \rightarrow \text{Iso}(V)$, $h(g)(v) := vg$ on the Banach space (V, N) is well defined and continuous.

Let $j^* : E^* \rightarrow V^*$ be the adjoint map of $j : V \rightarrow E$. Now our aim is to check the G -continuity of every vector $\varphi \in j^*(X_f) = cl_{w^*}(G\psi)$, where $\psi := j^*(z)$ and z denotes the point $f_{\sharp}(e) \in X_f$. By Lemma 9.7.3 we have to show that B_V is SUC-small for $j^*(X_f)$.

Claim : $j(B_V) \subset \bigcap_{n \in \mathbb{N}} K_n = \bigcap_{n \in \mathbb{N}} (2^n M + 2^{-n} B_E)$.

Proof. The norms $\|\cdot\|_n$ on E are equivalent to each other. It follows that if $v \in B_V$ then $\|v\|_n < 1$ for all $n \in \mathbb{N}$. That is, $v \in \lambda_n K_n$ for some $0 < \lambda_n < 1$ and $n \in \mathbb{N}$. By the construction K_n is a convex subset containing the origin. This implies that $\lambda_n K_n \subset K_n$. Hence $j(v) = v \in K_n$ for every $n \in \mathbb{N}$. \square

Recall now that FG is SUC-small for $X_f \subset C(X_f)^*$. By Lemma 9.7.1 we know that also $M := co(-FG \cup FG)$ is SUC-small for $X_f \subset C(X_f)^*$. Moreover by Lemma 9.7.2 we obtain that $A := \bigcap_{n \in \mathbb{N}} (2^n M + 2^{-n} B_E) \subset C(X_f)$ is SUC-small for $X_f \subset C(X_f)^*$. The linear continuous operator $j : V \rightarrow C(X_f)$ is a G -map. Then by Lemma 9.6.1 it follows that $j^{-1}(A) \subset V$ is SUC-small for $j^*(X_f) \subset V^*$. The same is true for B_V because by the above claim we have $j(B_V) \subset A$ (and hence, $B_V \subset j^{-1}(A)$). That is B_V is SUC-small for $j^*(X_f)$. Now Lemma 9.7.3 shows that the orbit map $\tilde{\varphi} : G \rightarrow V^*$ is G -continuous for every $\varphi \in j^*(X_f) = cl_{w^*}(G\psi)$. By our construction $F \in j(V)$ (because $F \in M \subset j(B_V)$). Since j is injective the element $v := j^{-1}(F)$ is uniquely determined in V . We show that $f = m_{v, \psi}$

for the co-representation h . Using the equality $F \circ \alpha_f = f$ and the fact that α_f is a G -map we get

$$\langle Fg, z \rangle = F(g\alpha_f(e)) = (F \circ \alpha_f)(g) = f(g).$$

On the other hand,

$$m_{v,\psi}(g) = \langle vg, \psi \rangle = \langle j^{-1}(F)g, j^*(z) \rangle = \langle j(j^{-1}(F))g, z \rangle = \langle Fg, z \rangle.$$

Hence, $f = m_{v,\psi}$, as required. Therefore we have proved that (1) \iff (2).

Finally we show that one can assume in (2) that $cl_{w^*}(G\psi) = j^*(X_f)$ separates points of V . If v_1, v_2 are different elements in V then $j(v_1) \neq j(v_2)$. Since X_f separates $C(X_f)$ then $\langle j(v_1), \phi \rangle \neq \langle j(v_2), \phi \rangle$ for some $\phi \in X_f$. Now observe that $\langle j(v), \phi \rangle = \langle v, j^*(\phi) \rangle$ for every $v \in V$. \square

Corollary 9.9. $Adj(G) \subset SUC(G)$.

Next we show how one can characterize $LE(G)$ in terms of matrix coefficients.

Theorem 9.10. *The following conditions are equivalent:*

- (1) $f \in LE(G)$.
- (2) $f = m_{v,\psi}$ for some continuous co-representation $h : G \rightarrow Iso(V)$, $v \in V$ and $\psi \in V^*$, for a Banach space V , with the property that the weak $*$ and the norm topologies coincide on the orbit $G\varphi$ of every φ in the weak $*$ closure $Y := cl_{w^*}(G\psi)$.

Moreover, one can assume in (2) that Y separates points of V .

Proof. (2) \implies (1): By definition f comes from $Y := cl_{w^*}(G\psi)$. Hence it suffices to show that Y is LE. Equivalently, we need to show that Y is orbitwise light (see [13, Lemma 5.8.2]). Let μ_Y be the uniform structure on the compact space Y . Denote by $(\mu_Y)_G$ the corresponding uniform structure of uniform convergence inherited from Y^G (see [13]). We have to show that $top(\mu_Y)|_{G\varphi} = top((\mu_Y)_G)|_{G\varphi}$ for every $\varphi \in Y$. Observe that the topology $(\mu_Y)_G$ on the orbit $G\varphi$ is weaker than the norm topology. Since the latter is the same as the weak star topology (that is, $top(\mu_Y)|_{G\varphi}$) we get that indeed $top(\mu_Y)|_{G\varphi} = top((\mu_Y)_G)|_{G\varphi}$.

(1) \implies (2): The proof uses again the interpolation technique of [7], as in Theorem 9.8. The proof is similar so we omit the details. However we provide necessary definition and two lemmas (Definition 9.11 and Lemmas 9.12 and 9.13). They play the role of Lemmas 9.6 and 9.7.

For every set M denote by \mathbb{R}^M the set of all real valued functions $M \rightarrow \mathbb{R}$. The topologies of pointwise and uniform convergence on \mathbb{R}^M will be denoted by τ_p and τ_u respectively.

Definition 9.11. Let $\langle \cdot, \cdot \rangle : Y \times X \rightarrow \mathbb{R}$ be a G -compatible map (as in Definition 9.5) and M be a nonempty subset of Y . Denote by $j : X \rightarrow \mathbb{R}^M$, $j(x)(m) := \langle m, x \rangle$ the associated map.

- (1) We say that a subset A of X is M -light if the pointwise and uniform topologies coincide on $j(A) \subset \mathbb{R}^M$.
- (2) M is LE -small at $x_0 \in X$ if the orbit Gx_0 is M -light.
- (3) M is LE -small for X if the orbit Gx is M -light at every $x \in X$ (compare Theorem 7.9).

We are going to examine this definition in a particular case of the canonical bilinear map $V \times V^* \rightarrow \mathbb{R}$ which is G -compatible for every co-representation $h : G \rightarrow Iso(V)$.

We collect here some useful properties of LE -smallness.

Lemma 9.12. (1) Let $Y_1 \times X_1 \rightarrow \mathbb{R}$ and $Y_2 \times X_2 \rightarrow \mathbb{R}$ be two G -compatible maps. Suppose that $\gamma_1 : X_1 \rightarrow X_2$ and $\gamma_2 : Y_2 \rightarrow Y_1$ are G -maps such that

$$\langle y, \gamma_1(x) \rangle = \langle \gamma_2(y), x \rangle \quad \forall (y, x) \in Y_2 \times X_1.$$

Then for every nonempty subset $M \subset Y_2$ the subset $\gamma_1(M) \subset Y_1$ is LE -small at $x \in X_1$ if and only if M is LE -small at $\gamma_1(x) \in X_2$.

- (2) Let X be a (not necessarily compact) G -space. If $f \in LE(X)$ then the subset $F_e G$ of $C(X_f)$ is LE -small for X_f considered as a subset of V^* where $V := C(X_f)$ (with respect to the canonical map $V \times V^* \rightarrow \mathbb{R}$ and the natural co-representation $G \rightarrow Iso(V)$).

Proof. (1): Similar to Lemma 9.6.1.

(2): Let $f \in LE(X)$. Then it comes by Proposition 3.1.3 from a compact G -system X_f and the G -compactification $f_{\sharp} : X \rightarrow X_f$. As we know there exists $F_e \in C(X_f)$ such that $f = F_e \circ f_{\sharp}$. Theorem 7.9 implies that X_f is LE and $F_e \in LE(X_f)$. By the same theorem, X_f is orbitwise light in $RUC(G)$. This means that pointwise and norm topologies in $RUC(G)$ agree on every G -orbit in X_f . On the other hand it is straightforward to see that for the G -compatible map

$$F_e G \times X_f \rightarrow \mathbb{R}$$

(Definition 9.11 with $M := F_e G$) the corresponding pointwise topology τ_p on X_f coincides with the pointwise topology inherited from $RUC(G)$ and the uniform topology τ_u on X_f coincides with the norm topology of $RUC(G)$. \square

Lemma 9.13. Let $h : G \rightarrow Iso(V)$ be a continuous co-representation.

- (a) For every $X \subset V^*$ the family of LE -small sets for X in V is closed under taking: subsets, norm closures, finite linear combinations, finite unions and convex hulls.

- (b) If $M_n \subset V$ is LE-small at $x_0 \in V^*$ for every $n \in \mathbb{N}$ then so is $\bigcap_{n \in \mathbb{N}} (M_n + \delta_n B_V)$ for every positive decreasing sequence $\delta_n \rightarrow 0$.
- (c) The following are equivalent:
- (i) The pointwise and norm topologies agree on the G -orbit Gx for every $x \in X \subset V^*$.
 - (ii) \mathcal{B} is LE-small for X , where $\mathcal{B} := \{\check{v} : V^* \rightarrow \mathbb{R}, x \mapsto \check{v}(x) := \langle v, x \rangle\}_{v \in B_V}$.

□

Question 9.14. Do we have $\text{adj}(G) = \text{SUC}(G)$ for every topological group G ? The question seems to be open even for abelian non-discrete G (say $G = \mathbb{R}$). The equivalent question for abelian G is whether $\text{adj}(G) = \text{UC}(G)$? Also, what is the relation between the algebras $\text{LE}(G)$ and $\text{adj}(G)$?

10. SOME CONCLUSIONS ABOUT $H_+[0, 1]$ AND $\text{Iso}(\mathbb{U}_1)$

From the reflexive triviality of $H_+[0, 1]$ and results of Uspenskij about $\text{Iso}(\mathbb{U}_1)$ Pestov deduces in [39, Corollary 1.4] the fact that the group $\text{Iso}(\mathbb{U}_1)$ is also reflexively trivial. Using a similar idea and the matrix coefficient characterization of SUC and LE one can conclude that $\text{Iso}(\mathbb{U}_1)$ is SUC-trivial and LE-trivial.

Recall the following results of Uspenskij.

Theorem 10.1. (Uspenskij [45, 47]) *The group $\text{Iso}(\mathbb{U}_1)$ is topologically simple and contains a copy of every second countable topological group (e.g., $H_+[0, 1]$).*

Lemma 10.2 below is a generalized version of Pestov's observation. Of course it is important here that the corresponding property admits a reformulation in terms of Banach space representations, which is the case, for instance, for SUC and LE.

Lemma 10.2. *Let G_1 be a topological subgroup of a group G_2 . Suppose that G_2 is G_1 -simple, in the sense that, every non-trivial normal subgroup N in G_2 containing G_1 is necessarily dense in G_2 . Then if G_1 is either: 1) SUC-trivial, 2) LE-trivial, 3) adjoint continuous trivial or 4) \mathcal{K} -trivial (where \mathcal{K} is a class of Banach spaces) then the same is true for G_2 .*

Proof. We consider only the case of SUC. Other cases are similar (and even easier for (3) and (4)).

We use Theorem 9.8. Let $h : G_2 \rightarrow \text{Iso}(V)$ be a continuous co-representation where $Y := \text{cl}_{w^*}(G\psi)$ separates points of V and φ is norm G_2 -continuous in V^* for every $\varphi \in Y$. It is enough to show that any such co-representation of G_2 is trivial. By Theorem 9.8 this will show that G_2 is SUC-trivial. First observe that the restriction $h|_{G_1}$ of h to G_1 is trivial. In fact, otherwise $vg \neq v$ for some $(v, g) \in$

$V \times G_1$ and by our assumption there exists $\varphi \in Y$ such that $\varphi(v) \neq \varphi(vg)$. Then the restriction $m_{v,\varphi}|_{G_1}$ of the corresponding matrix coefficient $m_{v,\varphi} : G_2 \rightarrow \mathbb{R}$ to G_1 is not constant. However, by Theorem 9.8, $m_{v,\varphi}|_{G_1} \in SUC(G_1)$, contradicting our assumption that G_1 is SUC-trivial. Therefore, $h|_{G_1} : G_1 \rightarrow Iso(V)$ is trivial. Hence G_1 is a subgroup of the normal closed subgroup $N := \ker(h)$ of G_2 . Since G_2 is G_1 -simple it follows that $N = G_2$. Hence h is trivial. \square

Note that if G_2 is *topologically simple*, that is $\{e\}$ -simple, then it is G_1 -simple for every subgroup.

The following theorem sums up some of our results concerning the topological groups $H_+[0, 1]$ and $Iso(\mathbb{U}_1)$.

Theorem 10.3. *Let G be one of the groups $H_+[0, 1]$ or $Iso(\mathbb{U}_1)$.*

- (1) *The compactifications $G^{SUC}, G^{LE}, G^{Asp}, G^{WAP}$ are trivial.*
- (2) *Every adjoint continuous (co)representation of the group G is trivial.*
- (3) *Every continuous Asplund (co)representation of the group G is trivial.*
- (4) *Every (co)representation $h : G \rightarrow Iso(V)$ on a separable Asplund space V is trivial.*
- (5) *The algebra $UC(G)$ and the ambit $(G^{UC}, i(e))$ are not point universal. In particular, the map $i : G \rightarrow G^{UC}$ is not a right topological compactification of G .*

Proof. (1): $H_+[0, 1]$ is SUC-trivial by Theorem 8.3. By results of Uspenskij (see Theorem 10.1) the group $Iso(\mathbb{U}_1)$ is topologically simple and also a universal second countable group. In particular it contains a copy G_1 of $H_+[0, 1]$ as a topological subgroup. It follows that $G_2 := Iso(\mathbb{U}_1)$ is G_1 -simple. Applying Lemma 10.2 we conclude that $G_2 := Iso(\mathbb{U}_1)$ is also SUC-trivial. The rest follows by the inclusions of Theorem 7.10.

(2): Every adjoint continuous (co)representation of $H_+[0, 1]$ must be trivial. Otherwise, by Theorem 9.8 (or Corollary 9.9), it contains a nonconstant SUC function. Now Lemma 10.2 implies that $Iso(\mathbb{U}_1)$ is also adjoint continuous trivial.

(3): By Theorem 6.5 every continuous Asplund (co)representation of G is adjoint continuous. Now apply (2).

(4): By a recent result of Rosendal and Solecki [42, Corollary 3] every homomorphism of $G = H_+[0, 1]$ into a separable group is necessarily continuous. Combining this result and our assertion (3) we obtain a proof in the case of $G = H_+[0, 1]$. The case of $G = Iso(\mathbb{U}_1)$ now follows by using again the $H_+[0, 1]$ -simplicity of $Iso(\mathbb{U}_1)$.

(5): Take a non-constant uniformly continuous function on G (such a function necessarily exists by Lemma 2.1). Since $SUC(G) = \{\text{constants}\}$ we get $SUC(G) \neq UC(G)$. Now Corollary 4.11 finishes the proof. \square

By Theorems 10.1 and 10.3 we get

Corollary 10.4. *Every second countable group G_1 is a subgroup of a Polish SUC-trivial group G_2 .*

However the following questions are open (see also [31]).

Question 10.5. (1) *Find a nontrivial Polish group which is SUC-trivial (Ref-trivial, Asp-trivial) but does not contain a subgroup topologically isomorphic to $H_+[0, 1]$.*

(2) *Is the group $H(I^\omega)$ SUC-trivial (Ref-trivial, Asp-trivial)?*

And, a closely related question (see Lemma 10.2):

(3) *Is the group $G_2 := H(I^\omega)$, G_1 -simple for a subgroup $G_1 < G_2$ where G_1 is a copy of either $H_+[0, 1]$ or of $\text{Iso}(\mathbb{U}_1)$?*

Theorem 10.6. *Let G be an Asplund trivial (e.g. $H_+[0, 1]$ or $\text{Iso}(\mathbb{U}_1)$) group. Then every metrizable right topological semigroup compactification of G is trivial.*

Proof. By Theorem 10.3.1, G^{Asp} is trivial, so that every RN transitive G -space is trivial. If $G \rightarrow S$ is a right topological semigroup compactification of G , then the natural induced G -space (G, S) is isomorphic to its own enveloping semigroup. By a recent work [14], a metric dynamical system (G, X) is RN iff its enveloping semigroup is metrizable. Now if S is metrizable then it follows that the transitive system (G, S) is RN and therefore trivial. \square

Recall, in contrast, that for every topological group G the algebra $RUC(G)$ separates points and closed subsets on G and therefore the maximal right topological semigroup compactification $G \hookrightarrow G^{\text{RUC}}$ is faithful.

11. RELATIVE EXTREME AMENABILITY: SUC-FPP GROUPS

Recall that a topological group G has the *fixed point on compacta property* (*fpp*) (or is *extremely amenable*) if every compact (transitive) G -space X has a fixed point. It is well known that locally compact extremely amenable groups are necessarily trivial (see for example [19]). Gromov and Milman [20] proved that the unitary group $U(H)$ is extremely amenable. Pestov has shown that the groups $H_+[0, 1]$ and $\text{Iso}(\mathbb{U}_1)$ are extremely amenable (see [37, 40] for more information).

Consider the following relativization.

Definition 11.1. Let P be a class of compact G -spaces.

- (1) A G -space X is P -*fpp* (or is *extreme P -amenable*) if every G -compactification $X \rightarrow Y$ such that Y is a member of P has a fixed point.
- (2) A topological group G is P -*fpp* (or is *extremely P -amenable*) if the G -space $X := G$ is P -fpp or equivalently, if every G -space Y in P has a fixed point.

Taking \mathcal{P} as the collection of all compact flows we get extreme amenability. With the class \mathcal{P} of compact affine flows we recover amenability. When \mathcal{P} is taken to be the collection of equicontinuous (that is, almost periodic) flows we obtain the old notion of minimal almost periodicity (MAP). Minimal almost periodicity was first studied by von Neumann and Wigner [34] who showed that $PSL(2, \mathbb{Q})$ has this property. See also Mitchell [33] and Berglund, Junghenn and Milnes [5].

Lemma 11.2. *Let \mathcal{P} be a class of compact G -spaces which is preserved by isomorphisms, products subsystems and quotients. Let \mathcal{P} and $X^{\mathcal{P}}$ be as in Section 2. The following conditions are equivalent:*

- (1) G is \mathcal{P} -fpp.
- (2) The compact G -space $G^{\mathcal{P}}$ is \mathcal{P} -fpp.
- (3) Any minimal compact G -space in \mathcal{P} is trivial.
- (4) For every $f \in \mathcal{P}$ the G -system X_f has a fixed point.
- (5) The algebra \mathcal{P} is extremely left amenable (that is it admits a multiplicative left invariant mean).

Proof. Clearly each of the conditions (1) and (3) implies all the others. Use the fact that the G -space $G^{\mathcal{P}}$ is point universal to deduce that each of (2) and (5) implies (3). Finally (4) implies (2) because $G^{\mathcal{P}}$ has a presentation as a subsystem of the product of all the X_f , $f \in \mathcal{P}$. Note that $f \in \mathcal{P}$ iff X_f has property \mathcal{P} (see [13, Proposition 2.9.3]). \square

Remark 11.3. (1) The smaller the class \mathcal{P} is one expects the property of being \mathcal{P} -fpp to be less restrictive; however even when one takes \mathcal{P} to be the class of equicontinuous \mathbb{Z} -spaces (that is, *cascades*) it is still an open question whether \mathcal{P} -fpp, that is minimal almost periodicity, is equivalent to extreme amenability (see [11]).

- (2) A minimal compact G -space X is LE iff X is AP. It follows by the inclusions $LE \supset RN_{\text{app}} \supset WAP \supset AP$ (cf. also Theorem 7.10) that G is minimally almost periodic iff G is \mathcal{P} -fpp for each of the following classes: WAP, RN_{app} or LE.

Here we point out two examples of topological groups G which are SUC-extremely amenable (equivalently, SUC-fpp) but not extremely amenable. In the next two sections we will show that S_{∞} as well as the group $H(C)$ of homeomorphisms of the Cantor set C are also SUC-fpp (both groups are not extremely amenable). See Corollary 12.3 and Theorem 13.8 below.

Example 11.4. For every $n \geq 2$ the simple Lie group $SL_n(\mathbb{R})$, being locally compact, is not extremely amenable. However it is SUC-extremely amenable. This follows easily from Corollary 7.15.

For the second example we need the following proposition.

Proposition 11.5. *Let G be a uniformly Lindelöf (e.g. second countable) group and X a not necessarily compact G -space. Assume that the action is 2-homogeneous on some uncountable dense orbit $Y = Gx_0$. Then X is SUC-trivial (and hence also SUC-fpp).*

Proof. It is enough to show that every $f \in \text{SUC}(X)$ is constant. We can suppose that $Y = X = Gx_0$. Let $St(x_0) < G$ be the stabilizer of x_0 . We observe that $St(x_0)$ is not open in G . Indeed, otherwise the space $G/St(x_0)$ is discrete and since G is uniformly Lindelöf it is easy to see that $G/St(x_0)$ is countable. However, by assumption X is uncountable and the cardinality of X is the same as that of the coset space $G/St(x_0)$. It follows that $Ux_0 \neq \{x_0\}$ for every neighborhood U of the identity $e \in G$. Now assume to the contrary that f is a nonconstant SUC function and let $\varepsilon := |f(a) - f(b)| > 0$ for some $a, b \in X$. By the definition of SUC there exists a neighborhood $U(e)$ such that

$$|f(gux_0) - f(gx_0)| < \varepsilon \quad \forall (g, u) \in G \times U.$$

As was already established there exists $x_1 = u_0x_0 \in Ux_0$ such that $x_1 \neq x_0$. Since the action is 2-homogeneous we can choose $g_0 \in G$ such that $g_0x_1 = a$ and $g_0x_0 = b$. Then $|f(g_0u_0x_0) - f(g_0x_0)| = \varepsilon$, a contradiction. \square

Now for many concrete homogeneous uncountable metric compact spaces X the natural action of the topological group $G = H(X)$ on X is 2-homogeneous. By Proposition 11.5 the flow (G, X) admits only constant SUC functions and the corresponding SUC G -compactification X^{SUC} is trivial. This is the case, to mention some concrete examples, for X the Cantor set (see Remark 13.5), the Hilbert cube and the circle \mathbb{T} .

In the latter case even the subgroup $G := H_+(\mathbb{T}) < H(\mathbb{T})$ of all orientation-preserving homeomorphisms of the circle acts 2-homogeneously on \mathbb{T} . Pestov has shown [37, 40] that the universal minimal dynamical G -system $M(G)$ for $G := H_+(\mathbb{T})$ coincides with the natural action of G on \mathbb{T} . Combining these results with Proposition 11.5 we obtain our second example.

Corollary 11.6. *The Polish group $G = H_+(\mathbb{T})$ of orientation preserving homeomorphisms of the circle is SUC-extremely amenable (but it is not extremely amenable).*

An alternative proof follows easily from Proposition 5.1.

12. THE ROELCKE COMPACTIFICATION OF THE GROUP $S(\mathbb{N})$

Let $G = S(\mathbb{N})$ be the Polish topological group of all permutations of the set \mathbb{N} of natural numbers (equipped with the topology of pointwise convergence). Consider the one point compactification $X^* = \mathbb{N} \cup \{\infty\}$ and the associated natural

G action (G, X^*) . For any subset $A \subset \mathbb{N}$ and an injection $\alpha : A \rightarrow \mathbb{N}$ let p_α be the map in $(X^*)^{X^*}$ defined by

$$p_\alpha(x) = \begin{cases} \alpha(x) & x \in A \\ \infty & \text{otherwise} \end{cases}$$

We have the following simple claim.

Claim 12.1. *The enveloping semigroup $E = E(G, X^*)$ of the G -system (G, X^*) consists of the maps $\{p_\alpha : \alpha : A \rightarrow \mathbb{Z}\}$ as above. Every element of E is a continuous function so that by the Grothendieck-Ellis-Nerurkar theorem [9], the system (G, X^*) is WAP.*

Proof. Let π_ν be a net of elements of $S(\mathbb{N})$ with $p = \lim_\nu \pi_\nu$ in E . Let $A = \{n \in \mathbb{N} : p(n) \neq \infty\}$ and $\alpha(n) = p(n)$ for $n \in A$. Clearly $\alpha : A \rightarrow \mathbb{N}$ is an injection and $p = p_\alpha$.

Conversely given $A \subset \mathbb{N}$ and an injection $\alpha : A \rightarrow \mathbb{N}$ we construct a sequence π_n of elements of $S(\mathbb{N})$ as follows. Let $A_n = A \cap [1, n]$ and $M_n = \max\{\alpha(i) : i \in A_n\}$. Next define an injection $\beta_n : [1, n] \rightarrow \mathbb{N}$ by

$$\beta_n(j) = \begin{cases} \alpha(j) & j \in A \\ j + M_n + n & \text{otherwise.} \end{cases}$$

Extending the injection β_n to a permutation π_n of \mathbb{N} , in an arbitrary way, we now observe that $p_\alpha = \lim_{n \rightarrow \infty} \pi_n$ in E . The last assertion is easily verified. \square

Theorem 12.2. (1) *The two algebras $UC(G)$ and $WAP(G)$ coincide.*

(2) *The universal WAP compactification G^{WAP} of G (and hence also G^{UC}), is isomorphic to $E = E(G, X^*)$. Thus the universal WAP (and Roelcke) compactification of G is homeomorphic to the Cantor set.*

Proof. Given $f \in UC(G)$ and an $\varepsilon > 0$ there exists $k \in \mathbb{N}$ such that — with $H = H(1, \dots, k) = \{g \in G : g(j) = j, \forall 1 \leq j \leq k\}$ —

$$\sup_{u, v \in H} |f(ugv) - f(g)| < \varepsilon.$$

Set $\hat{f}(g) = \sup_{u, v \in H} f(ugv)$, then $\|\hat{f} - f\| \leq \varepsilon$. Clearly \hat{f} , being H -biinvariant, is both right and left uniformly continuous; i.e. $\hat{f} \in UC(G)$. Let

$$\mathbb{N}_*^k = \{(n_1, n_2, \dots, n_k) : n_j \in \mathbb{N} \text{ are distinct}\} = \{\text{injections} : \{1, 2, \dots, k\} \rightarrow \mathbb{N}\}$$

and let G act on \mathbb{N}_*^k by

$$g(n_1, n_2, \dots, n_k) = (g^{-1}n_1, g^{-1}n_2, \dots, g^{-1}n_k).$$

The stability group of the point $(1, \dots, k) \in \mathbb{N}_*^k$ is just H and we can identify the discrete G -space G/H with \mathbb{N}_*^k . Under this identification, to a function $f \in$

$UC(G)$ which is right H -invariant (that is $f(gh) = f(g)$, $\forall g \in G, h \in H$), corresponds a bounded function $\omega_f \in \Omega_k = \mathbb{R}^{\mathbb{N}_*^k}$, namely

$$\omega_f(n_1, n_2, \dots, n_k) = f(g) \quad \text{iff} \quad g(j) = n_j, \quad \forall 1 \leq j \leq k.$$


If we now assume that $f \in UC(G)$ is both right and left H -invariant (so that $f = \hat{f}$) then, as we will see below, f and accordingly its corresponding ω_f , admits only finitely many values, corresponding to the finitely many double H cosets $\{HgH : g \in G\}$.

We set $Y_f = Y = \text{cl}\{g\omega_f : g \in G\} \subset \Omega_k = \mathbb{R}^{\mathbb{N}_*^k}$, where the closure is with respect to the pointwise convergence topology. (G, Y_f) is a compact G -system which is isomorphic, via the identification $G/H \cong \mathbb{N}_*^k$, to $X_f \subset \mathbb{R}^G$. We will refer to elements of $\Omega_k = \mathbb{R}^{\mathbb{N}_*^k}$ as *configurations*. Consider first the case $k = 2$.

In the following figure we have a representation of the configuration $f = \omega_f = \omega_{1,2}$ and three other typical elements of Y_f . The configuration $\omega_{2,7} = \sigma\omega_{1,2}$ — where σ is the permutation $(\begin{smallmatrix} 1 & 2 & 7 \\ 7 & 1 & 2 \end{smallmatrix})$ —, admits seven values (the maximal number it can possibly have): “blank” at points (m, n) with $m, n \notin \{2, 7\}$, the values \diamond and $*$ at $(2, 7)$ and $(7, 2)$ respectively, and four more constant values on the two horizontal and two vertical lines. (The circled diagonal points $(2, 2)$ and $(7, 7)$ are by definition not in \mathbb{N}_*^2 .) If we let π_n be the permutation

$$\pi_n(j) = \begin{cases} j & j \notin \{1, n\} \\ n & \text{for } j = 1 \\ 1 & \text{for } j = n \end{cases}$$

and denote by $p = \lim_{n \rightarrow \infty} \pi_n$, the corresponding element of $E(G, X)$ then, e.g. $\omega_{1,\infty} = p\omega_{1,2} = \lim_{n \rightarrow \infty} \pi_n\omega_{1,2} = \lim_{n \rightarrow \infty} \omega_{1,n}$.



The functions $\omega_{1,2}$, $\omega_{1,\infty}$, $\omega_{2,7}$ and $\omega_{2,\infty}$ in Ω_2

Now it is not hard to see that the G action on Y naturally extends to an action of $E = E(G, X^*)$ on Y where each $p \in E$ acts continuously. (Show that

the map $(a, b) \mapsto \omega_{a,b}$ is an isomorphism of G -systems from $X^* \times X^* \setminus \Delta$ onto Y , where $\Delta = \{(n, n) : n \in \mathbb{N}\}$.) It then follows that $E = E(G, X^*)$ coincides with $E(G, Y)$.

By the Grothendieck-Ellis theorem (see e.g. [12, Theorem 1.45]) these observations show that the G -space Y_f is WAP and therefore the function f , which comes from (G, Y_f) , is a WAP function.

These considerations are easily seen to hold for any positive integer k . For example, an easy calculation shows that for $H_k = H(1, 2, \dots, k)$ the number of double cosets $\{H_k g H_k : g \in G\}$ is

$$\sum_{j=0}^k \binom{k}{j} \frac{k!}{(k-j)!}$$

Since the subgroups $H_k = H(1, \dots, k)$ form a basis for the topology at e , as we have already seen, the union of the H_k -biinvariant functions for $k = 2, 3, \dots$ is dense in $UC(G)$ and we conclude that indeed $UC(G) = WAP(G)$.

Since for each H -biinvariant function f the enveloping semigroup of the dynamical system (G, Y_f) is isomorphic to $E(G, X^*)$ and since the Gelfand compactification of $UC(G)$ is isomorphic to a subsystem of the direct product

$$\prod \{Y_f : f \text{ is } H\text{-biinvariant for some } H(1, \dots, k)\}$$

we deduce that $E(G, X^*)$ serves also as the enveloping semigroup of the universal dynamical system $|WAP(G)| = |UC(G)|$. Finally, since $|WAP(G)|$ is point universal we conclude (by Lemma 3.3) that $(G, |WAP(G)|)$ and $(G, E(G, X^*))$ are G -isomorphic. \square

Corollary 12.3. *The Polish group $G = S(\mathbb{N})$ is SUC-extremely amenable but not extremely amenable.*

Proof. It was shown by Pestov [37] that G is not extremely amenable and the nontrivial universal minimal G -system was described in [17]. On the other hand the G -system G^{UC} described in Theorem 12.2 admits a unique minimal set which is a fixed point. Thus the SUC G -system G^{SUC} , being G -isomorphic to G^{UC} (see Theorems 12.2 and 7.10.2), has a fixed point. \square

13. THE HOMEOMORPHISMS GROUP OF THE CANTOR SET

In this section let C denote the classical Cantor set — i.e. the ternary subset of the interval $[0, 1]$. Thus C has the following representation:

$$C = \bigcap_{n=0}^{\infty} I^n,$$

where $I^n = \bigcup_{j=1}^{2^n} I_j^n$, is the disjoint union of the 2^n closed intervals obtained by removing from $I = [0, 1]$ the appropriate $2^n - 1$ open ‘middle third’ intervals. In

the sequel we will write I_j^m for the clopen subset $I_j^m \cap C$ of C . For each integer $m \geq 1$, $\mathbf{I}^m = \{I_j^m : 1 \leq j \leq 2^m\}$ denotes the basic partition of C into 2^m clopen “intervals”.

We let $G = H(C)$ be the Polish group of homeomorphisms of C equipped with the topology of uniform convergence. For $n \in \mathbb{N}$ we let

$$H_n = \{g \in G : gI_j^n = I_j^n, \forall 1 \leq j \leq 2^n\}.$$

Each H_n is a clopen subgroup of G and we note that the system of clopen subgroups $\{H_n : n = 2, 3, \dots\}$ forms a basis for the topology of G at the identity $e \in G$.

For any fixed integer $k \geq 1$ consider the collection

$$\mathcal{A}^k = \{\mathbf{a} = \{A_1, A_2, \dots, A_k\} : \text{a partition of } C \text{ into } k \text{ nonempty clopen sets}\}.$$

In particular note that for $k = 2^n$, \mathbf{I}^n is an element of \mathcal{A}^k .

The discrete homogeneous space G/H_n can be identified with $\mathcal{A}^k = \mathcal{A}^{2^n}$: an element $gH_n \in G/H_n$ is uniquely determined by the partition

$$\mathbf{a} = \{gI_j^n : 1 \leq j \leq 2^n = k\},$$

and conversely to every partition $\mathbf{a} \in \mathcal{A}^k$ corresponds a coset $gH_n \in G/H_n$. In fact, if $\mathbf{a} = \{A_1, A_2, \dots, A_k\}$ we can choose g to be any homeomorphism of C with $A_j = gI_j^n$.

Thus for $k = 2^n$ we have a parametrization of \mathcal{A}^k by the discrete homogeneous space G/H_n .

Let

$$\Omega^k = \mathbb{R}^{\mathcal{A}^k} \cong \mathbb{R}^{G/H_n}.$$

Via the quotient map $G \rightarrow G/H_n$, $g \mapsto gH_n$, the Banach space $\ell^\infty(\mathcal{A}^k)$ canonically embeds into the Banach space $RUC(G)$ where the image consists of all the right H_n -invariant functions in $RUC(G)$. Thus if $f \in RUC(G)$ satisfies $f(gh) = f(g)$ for all $g \in G$ and $h \in H_n$ then $\omega_f(gI_1^n, \dots, gI_k^n) = \omega_f(A_1, \dots, A_k) = f(g)$, where $A_j = gI_j^n$, is the corresponding configuration in $\ell^\infty(\mathcal{A}^k)$.

We equip $\Omega^k = \mathbb{R}^{\mathcal{A}^k}$ with its product topology. The group G acts on the space Ω^k as follows. For $\omega \in \Omega^k$ and $g \in G$ let

$$g\omega(\mathbf{a}) = \omega(g^{-1}A_1, g^{-1}A_2, \dots, g^{-1}A_k),$$

for any $\mathbf{a} = \{A_1, A_2, \dots, A_k\} \in \mathcal{A}^k$. Equivalently $g\omega(g'H_n) = \omega(g^{-1}g'H_n)$, for every $g'H_n \in G/H_n$. For each right H_n -invariant functions f in $RUC(G)$ we denote the compact orbit closure of $f = \omega_f$ in Ω^k by Y_f .

First let us consider the case $n = 1$, where $k = 2$,

$$\mathcal{A}^2 = \{\mathbf{a} = \{A, A^c\} : \text{a partition of } C \text{ into 2 nonempty clopen sets}\},$$

and

$$H = H_1 = \{g \in G : gI_j^1 = I_j^1, \quad j = 1, 2\}.$$

Claim 13.1. *There are exactly seven double cosets HgH , $g \in G$.*

Proof. For a partition $(A, A^c) \in \mathcal{A}^2$ exactly one of the following five possibilities holds: (1) $A = I_1^1$, (2) $A = I_2^1$, (3) $A \subsetneq I_1^1$, (4) $A \subsetneq I_2^1$, (5) $A \supsetneq I_1^1$, (6) $A \supsetneq I_2^1$, (7) $A \cap I_1^1 \neq \emptyset \neq A \cap I_2^1$, and $A^c \cap I_1^1 \neq \emptyset \neq A^c \cap I_2^1$.

Clearly for any two partitions $(A, A^c), (B, B^c)$ we have $(B, B^c) = (hA, hA^c)$ for some $h \in H$ iff they belong to the same class. Our claim follows in view of the correspondence $G/H \cong \mathcal{A}^2$. \square

Define an element $\omega_f \in \Omega^2$ and the corresponding function $f \in UC(G)$ as follows:

$$\omega(A, A^c) = j \quad \text{if } (A, A^c) \text{ is of type } (j), \quad j = 1, \dots, 7,$$

and $f(g) = \omega(gI_1^1, gI_2^1)$. Clearly f is H_1 -biinvariant and in particular an element of $UC(G)$. Let X_f denote the (pointwise) orbit closure of f in $RUC(G)$. Via the natural lift of Ω^2 to $RUC(G)$ we can identify X_f with $Y_f = \text{cl} \{g \cdot \omega_f : g \in G\} \subset \Omega^2$.

Next consider a sequence of homeomorphisms $h_n \in G$ satisfying the conditions

- (i) $h_n(I_1^n) = (I_{2^n}^n)^c$,
- (ii) $h_n((I_1^n)^c) = I_{2^n}^n$ and
- (iii) h_n is order preserving.

It is then easy to check that the limit $\lim_{n \rightarrow \infty} h_n \omega_f = \omega_0$ exists in Ω^2 where ω_0 is defined by

$$\omega_0(A, A^c) = \begin{cases} 5 & \text{if } 0 \in A \\ 4 & \text{if } 0 \notin A. \end{cases}$$

Now for any $g \in G$ we have

$$(g \cdot \omega_0)(A, A^c) = \omega_0(g^{-1}A, g^{-1}A^c) = \begin{cases} 5 & \text{if } g(0) \in A \\ 4 & \text{if } g(0) \notin A. \end{cases}$$

For $x \in C$ set

$$(\omega_x)(A, A^c) = \begin{cases} 5 & \text{if } x \in A \\ 4 & \text{if } x \notin A. \end{cases}$$

Then for $g \in G$ we have $g\omega_0 = \omega_{g0}$. Moreover denoting $Y_0 = \text{cl} \{g\omega_0 : g \in G\} \subset \Omega^2$ we have $Y_0 = G\omega_0$ and the map $\phi : (G, C) \rightarrow (G, Y_0)$ defined by $\phi(x) = \omega_x$ is an isomorphism of G -spaces. We get the following lemma.

Lemma 13.2. *Let $Y_0 = \text{cl} \{g\omega_0 : G \in G\}$ be the orbit closure of ω_0 in Ω^2 , then the G -space (G, Y_0) is isomorphic to (G, C) , the natural action of $G = H(C)$ on the Cantor set C .*

Remark 13.3. An argument analogous to that of Lemma 13.2 will show that for every n the number of H_n double cosets is finite. As in the case of $S(\mathbb{N})$ in

the previous section this shows the well known fact that $G = H(C)$ is Roelcke precompact (see [48]).

In contrast to Theorem 12.2 we obtain the following result.

Theorem 13.4. *For $G = H(C)$ we have $UC(G) \not\cong SUC(G)$.*

Proof. Consider the function

$$f_0(g) = \omega_0(g^{-1}I_1^1, g^{-1}I_2^1) = g\omega_0(I_1^1, I_2^1) = \omega_{g0}(I_1^1, I_2^1)$$

and let $h_n \in G$ be defined as above. Let u_n be a sequence of elements of G which converges to $e \in G$ and for which $h_n u_n 0 = 2/3$. Then, as $h_n 0 = 0$ for every n , we have

$$f_0(h_n) = \omega_0(h_n^{-1}I_1^1, h_n^{-1}I_2^1) = 5,$$

but as $h_n u_n 0 = 2/3$,

$$f_0(h_n u_n) = \omega_0(u_n^{-1}h_n^{-1}I_1^1, u_n^{-1}h_n^{-1}I_2^1) = \omega_{h_n u_n 0}(I_1^1, I_2^1) = 4.$$

Thus f_0 is not left uniformly continuous. Since $f_0 \in X_f \cong Y_0$, we conclude, by Theorem 4.12, that f is not a SUC function. \square

Remark 13.5. A similar argument will show that any two points $a, b \in C$ are SUC-proximal for the G -space (G, C) . Thus this G -space is SUC-trivial by Lemma 8.2. Letting $F : Y_0 \rightarrow \{4, 5\} \subset \mathbb{R}$ be the evaluation function $F(\omega) = \omega(I_1^1, I_2^1)$, we observe that

$$f_0(g) = \omega_0(g^{-1}I_1^1, g^{-1}I_2^1) = g\omega_0(I_1^1, I_2^1) = F(g\omega_0) = F(g\phi 0) = (F \circ \phi)(g0).$$

Thus the function f_0 comes from the G space C , via the continuous function $F \circ \phi : C \rightarrow \mathbb{R}$ and the point $0 \in C$. This is another way of showing that f_0 and hence also f are not SUC.

Lemma 13.6. *Let $\mathfrak{A} \subset RUC(G)$ be a closed G -invariant subalgebra containing the constants. Given $f \in \mathfrak{A}$ and $\varepsilon > 0$ there exist an n and an H_n -invariant function $\hat{f} \in \mathfrak{A}$ with $\|f - \hat{f}\| < \varepsilon$.*

Proof. By right uniform continuity there exists an n such that $\sup\{|f(hg) - f(g)| : g \in G\} < \varepsilon$ for every $h \in H_n$. Set $\hat{f}(g) = \sup_{h \in H_n} f(hg)$. This function is continuous because it is defined via the quotient map $G \rightarrow G/H_n$, where G/H_n is discrete. Since for every finite set $\{h_1, \dots, h_t\} \subset H_n$ the function

$$f_{h_1, \dots, h_t}(g) = \max_{1 \leq j \leq t} f(h_j g)$$

is in \mathfrak{A} , Dini's theorem implies that also \hat{f} is an element of \mathfrak{A} . \square

Remark 13.7. (1) By Theorem 13.4 and Corollary 4.11 we obtain, in particular, that the algebra $UC(G)$ is not point universal and the corresponding Roelcke compactification $G \rightarrow G^{UC}$ is not a right topological semigroup compactification of G . The same is true for $G := H_+[0, 1]$ because $UC(G) \neq SUC(G)$. This follows from Theorem 8.3 and Lemma 2.1.

(2) Using Lemma 13.6 one can try to determine, in analogy with the previous section, the structure of the universal WAP G -space, which is apparently much smaller than the Roelcke compactification of G .

Theorem 13.8. *The Polish group $G = H(C)$ of homeomorphisms of the Cantor set C has the SUC-fpp.*

Proof. Let f be a minimal, SUC function (that is, f is in $SUC(G)$ and X_f is a minimal G -space). Let \mathcal{A} be the smallest uniformly closed G -invariant algebra of $RUC(G)$ containing f and the constant functions. As we have seen above for every $\varepsilon > 0$ there exist n and $\hat{f} \in RUC(G)$ such that, with

$$H_n = \{g \in G : gI_j^n = I_j^n, \forall 1 \leq j \leq 2^n\},$$

\hat{f} is H biinvariant and $\|f - \hat{f}\| < \varepsilon$. By Lemma 13.6, \hat{f} is still an element of the algebra $\mathcal{A} \cap SUC(G)$. It clearly suffices to show that \hat{f} is a constant. For convenience of notation we will therefore assume that $f = \hat{f}$. Let now $\omega_f(A_1, A_2, \dots, A_{2^n})$ be the corresponding configuration in Ω^{2^n} . As we have seen the function $f(g) = \omega_f(g^{-1}I_1^n, \dots, g^{-1}I_{2^n}^n)$ has a finite range and again an application of the sequence h_n to ω_f will yield the limit configuration

$$\omega_0(A_1, A_2, \dots, A_{2^n}) = \begin{cases} a & \text{if } 0 \in A_1 \\ b & \text{if } 0 \notin A_1 \end{cases}$$

for some $a, b \in \mathbb{R}$. Since f is a SUC function $X_f \subset SUC(G)$, by Theorem 4.12, and the function $f_0(g) = \omega_0(g^{-1}\mathbf{I}^n)$, which is an element of X_f , is SUC. Since f_0 comes from the coset G -space C (see Lemma 13.2 and Remark 13.5) Theorem 5.1 implies that it is constant (i.e. $a = b$). However, by assumption X_f is a minimal G -space and it follows that $f = f_0$ is constant. \square

REFERENCES

1. E. Akin, J. Auslander and K. Berg, *When is a transitive map chaotic*, Convergence in Ergodic Theory and Probability, Walter de Gruyter & Co. 1996, pp. 25-40.
2. E. Akin, J. Auslander and K. Berg, *Almost equicontinuity and the enveloping semigroup*, Topological dynamics and applications, Contemporary Mathematics **215**, a volume in honor of R. Ellis, 1998, pp. 75-81.
3. E. Akin and E. Glasner, *Residual properties and almost equicontinuity*, J. d'Anal. Math. **84**, (2001), 243-286.
4. W. Banaschzyk, *Additive subgroups of topological vector spaces*, Lecture Notes in Math. **1466**, Springer-Verlag, 1991.
5. J.F. Berglund, H.D. Junghenn and P. Milnes, *Compact right topological semigroups and generalizations of almost periodicity*, Lecture Notes in Math. **663**, Springer-Verlag, 1978.

6. R.D. Bourgin, *Geometric Aspects of Convex Sets with the Radon-Nikodým Property*, Lecture Notes in Math. **993**, Springer-Verlag, 1983.
7. W.J. Davis, T. Figiel, W.B. Johnson and A. Pelczyński, *Factoring weakly compact operators*, *J. of Funct. Anal.* **17** (1974), 311-327.
8. R. Ellis, *Equicontinuity and almost periodic functions*, Proc. Amer. Math. Soc. **10**, (1959), 637-643.
9. R. Ellis and M. Nerurkar, *Weakly almost periodic flows*, Trans. Amer. Math. Soc. **313** (1989), 103-119.
10. M. Fabian, *Gateaux differentiability of convex functions and topology. Weak Asplund spaces*, Canadian Math. Soc. Series of Monographs and Advanced Texts, A Wiley-Interscience Publication, New York, 1997.
11. E. Glasner, *On minimal actions of Polish groups*, Topology Appl. **85** (1998), 119-125.
12. E. Glasner, *Ergodic theory via joinings*, AMS, Surveys and Monographs, **101**, 2003.
13. E. Glasner and M. Megrelishvili, *Linear representations of hereditarily non-sensitive dynamical systems*, Colloq. Math. **104** (2006), no. 2, 223-283.
14. E. Glasner, M. Megrelishvili and V.V. Uspenskij, *On metrizable enveloping semigroups*, Preprint, June, 2006. ArXiv e-print, Article math. DS/0606373, to appear in Israel J. Math.
15. E. Glasner and B. Weiss, *Sensitive dependence on initial conditions*, Nonlinearity **6** (1993), 1067-1075.
16. E. Glasner and B. Weiss, *Locally equicontinuous dynamical systems*, Colloquium Mathematicum, part 2, **84/85** (2000), 345-361.
17. E. Glasner and B. Weiss, *Minimal actions of the group $S(Z)$ of permutations of the integers*, Geom. and Funct. Anal. **12** (2002), 964-988.
18. E. Glasner and B. Weiss, *The universal minimal system for the group of homeomorphisms of the Cantor set*, Fund. Math. **176** (2003), no. 3, 277-289.
19. E. Granirer and A.T. Lau, *Invariant means on locally compact groups*, Ill. J. Math. **15** (1971), 249-257.
20. M. Gromov and V.D. Milman, *A topological application of the isoperimetric inequality*, Amer. J. Math. **105** (1983), 843-854.
21. A. Grothendieck, *Critères de compacité dans les espaces fonctionelles généraux*, Amer. J. Math. **74** (1952), 168-186.
22. W. Herer and J.P.R. Christensen, *On the existence of pathological submeasures and the construction of exotic topological groups*, Math. Ann. **213** (1975), 203-210.
23. J.E. Jayne and C.A. Rogers, *Borel selectors for upper semicontinuous set-valued maps*, Acta Math. **155**, (1985), 41-79.
24. M. Megrelishvili, *A Tychonoff G -space not admitting a compact G -extension or a G -linearization*, Russ. Math. Surv. **43:2** (1988), 177-178.
25. M. Megrelishvili, *Free topological G -groups*, New Zealand J. of Math. **25** (1996), 59-72.
26. M. Megrelishvili, *Fragmentability and continuity of semigroup actions*, Semigroup Forum **57** (1998), 101-126.
27. M. Megrelishvili, *Operator topologies and reflexive representability*, In: "Nuclear groups and Lie groups" Research and Exposition in Math. series, vol. **24**, Heldermann Verlag Berlin, 2001, 197-208.
28. M. Megrelishvili, *Every semitopological semigroup compactification of the group $H_+[0, 1]$ is trivial*, Semigroup Forum **63:3** (2001), 357-370.
29. M. Megrelishvili, *Reflexively but not unitarily representable topological groups*, Topology Proceedings **25** (2002), 615-625.
30. M. Megrelishvili, *Fragmentability and representations of flows*, Topology Proceedings **27:2** (2003), 497-544. See also: www.math.biu.ac.il/~megereli.
31. M. Megrelishvili, *Topological transformation groups: selected topics*, in: Open Problems In Topology II (Elliott Pearl, ed.), Elsevier Science, 2007, pp. 423-438.
32. M. Megrelishvili, *Every topological group is a group retract of a minimal group*, preprint, October, 2006. ArXiv e-print, article math.GN/0610226, to appear in Topology Appl.
33. T. Mitchell, *Function algebras, means and fixed points*, Trans. AMS **130** (1968), 117-126.
34. J. von Neumann and E.P. Wigner, *Minimally almost periodic groups*, Ann. of Math. **41** (1940), 746-750.

35. E.C. Nummela, On epimorphisms of topological groups, *Gen. Top. and its Appl.* **9** (1978), 155-167.
36. V. Pestov, *Epimorphisms of Hausdorff groups by way of topological dynamics*, *New Zealand J. of Math.* **26** (1997), 257-262.
37. V. Pestov, *On free actions, minimal flows and a problem by Ellis*, *Trans. Amer. Math. Soc.* **350** (1998), 4149-4165.
38. V. Pestov, *Topological groups: where to from here?* *Topology Proceedings* **24** (1999), 421-502. <http://arXiv.org/abs/math.GN/9910144>.
39. V. Pestov, *The isometry group of the Urysohn space as a Lévy group*, *Topology Appl.* **154** (2007), 2173-2184.
40. V. Pestov, *Dynamics of infinite-dimensional groups. The Ramsey-Dvoretzky-Milman phenomenon*. Revised edition of *Dynamics of infinite-dimensional groups and Ramsey-type phenomena* [Inst. Mat. Pura. Apl. (IMPA), Rio de Janeiro, 2005; MR2164572]. University Lecture Series, **40**. American Mathematical Society, Providence, RI, 2006. viii+192 pp.
41. W. Roelcke and S. Dierolf, *Uniform structures on topological groups and their quotients*, Mc Graw-hill, New York, 1981.
42. C. Rosendal and S. Solecki, *Automatic Continuity of Homomorphisms and Fixed Point on Metric Compacta*, *Israel J. Math.* **162** (2007), 349-371.
43. S. Teleman, *Sur la représentation linéaire des groupes topologiques*, *Ann. Sci. Ecole Norm. Sup.* **74** (1957), 319-339.
44. V.V. Uspenskij, *The epimorphism problem for Hausdorff topological groups*, *Topology Appl.* **57** (1994), 287-294.
45. V.V. Uspenskij, *On subgroups of minimal topological groups*, Accepted, to appear in Proceedings of the International Workshop on the Urysohn metric space (Ben-Gurion University, Israel, May 21-14, 2006), a special Volume of *Topology and its Applications*.
46. V.V. Uspenskij, *The Roelcke compactification of unitary groups*, in: *Abelian groups, module theory, and topology*, Proceedings in honor of Adalberto Orsatti's 60th birthday (D. Dikranjan, L. Salce, eds.), *Lecture notes in pure and applied mathematics*, Marcel Dekker, New York e.a. **201** (1998), 411-419.
47. V.V. Uspenskij, *Compactifications of topological groups*, Proceedings of the Ninth Prague Topological Symposium (Prague, August 19-25, 2001). Edited by Petr Simon. Published April 2002 by *Topology Atlas* (electronic publication). Pp. 331-346, arXiv:math.GN/0204144.
48. V.V. Uspenskij, *The Roelcke compactification of groups of homeomorphisms*, *Topology Appl.* **111** (2001), 195-205.
49. W.A. Veech, *Weakly almost periodic functions on semisimple Lie groups*, *Monat. Math.* **88**, (1979), 55-68.
50. J. de Vries, *Equivariant embeddings of G-spaces*, in: J. Novak (ed.), *General Topology and its Relations to Modern Analysis and Algebra IV*, Part B, Prague, 1977, 485-493.
51. J. de Vries, *Elements of Topological Dynamics*, Kluwer Academic Publishers, 1993.

DEPARTMENT OF MATHEMATICS, TEL-AVIV UNIVERSITY, RAMAT AVIV, ISRAEL
 E-mail address: glasner@math.tau.ac.il
 URL: <http://www.math.tau.ac.il/~glasner>

DEPARTMENT OF MATHEMATICS, BAR-ILAN UNIVERSITY, 52900 RAMAT-GAN, ISRAEL
 E-mail address: megereli@math.biu.ac.il
 URL: <http://www.math.biu.ac.il/~megereli>