

A SUBJECTIVE APPROACH TO QUANTUM PROBABILITY

EHUD LEHRER AND ERAN SHMAYA

May 17, 2005

ABSTRACT. A likelihood order is defined over linear subspaces of a finite dimensional Hilbert space. The question arises as to when such an order can be represented by a quantum probability. We introduce a few behaviorally plausible axioms that provide the answer in two cases: pure state and uniform measure. The general problem is answered by using duality-like conditions.

The general problem of characterizing the partial orders that admit a quantum representation by behaviorally justified axioms remains open.

Keywords: subjective probability, decision theory, quantum probability

1. INTRODUCTION

According to the subjective approach probabilities are merely degrees-of-belief of a rational agent. These degrees-of-belief might be indicated by the agent's willingness to bet or take other actions (see [6]). Savage ([15]) derives both probabilities and utilities from rational preferences (i.e., that satisfy some putative properties) alone. Such preferences induce, in particular, a preference order over events. That is, an agent who holds rational preferences could indicate which of two events is more likely, and moreover, this likelihood order is transitive. Savage's first step is to derive a (finitely additive) probability that represents the likelihood order.

In this paper we adopt a similar approach and apply it to the quantum framework without going beyond probabilities. While classical probability is defined over subsets (events) of a state space, quantum probability is defined over subspaces of Hilbert space. Furthermore, disjointness of the classical model is replaced by orthogonality.

Formally, let \mathcal{H} be a separable Hilbert space. A quantum probability measure μ over \mathcal{H} assigns a number between 0 and 1 to every closed

School of Mathematical Sciences, Tel Aviv University, Tel Aviv 69978, Israel.
e-mails: lehrer@post.tau.ac.il ; gawain@post.tau.ac.il .

subspace that satisfies $\mu(A \oplus B) = \mu(A) + \mu(B)$ whenever $A \perp B$ and $\mu(\mathcal{H}) = 1$. Gleason's Theorem (Gleason (1957)) states that, if $\dim(\mathcal{H}) \geq 3$ every quantum measure μ is induced by a self-adjoint nonnegative operator T with trace 1 in the following way: $\mu(A) = \text{tr}(\Pi_A T)$ for every subspace A , where Π_A is the orthogonal projection over A .

We assume the existence of a likelihood order \preceq over subspaces of a given finite-dimensional Hilbert space. The statement 'A is less likely than B in one's eyes' could be understood operationally: one would prefer betting that B occurs than that A occurs (in the corresponding physical measurements).

We say that the likelihood order \preceq can be *represented by a quantum probability* μ if $A \preceq B$ if and only if $\mu(A) \leq \mu(B)$. The goal of the line of research presented here is to find plausible properties (axioms, in the jargon of decision theory), preferably rationality-motivated, that ensure that \preceq is representable by a quantum probability.

Throughout, it is assumed that \preceq possesses three properties. The first is that every subspace is more likely than the zero-dimensional one. The second is that a subspace B is more likely than A if and only if $B + C$ is more likely than $A + C$, whenever $C \perp A$ and $C \perp B$. That is, adding or deleting a subspace which is orthogonal to both A and B would preserve the likelihood order.

The classical counterpart of the third property is a consequence of the second. However, in the quantum model it has to be explicitly assumed. It states that if B is more likely than A , then the orthogonal complement of B is less likely than that of A .

Savage ([15]) also assumes these three axioms but, in order to obtain a representation by a measure, he needs an additional, less motivated, property that concerns with the richness of the state space. This one dictates that the state space could be split into mutually disjoint arbitrarily small (with respect to the likelihood order) subsets. The lack of a quantum counterpart (in the case of a finite-dimensional Hilbert space) of such an Archimedean property makes our study completely different from that of Savage.

Our main results refer to likelihood orders that can be represented by two types of quantum measures. The first is the most important from a physical point of view. The probabilities of this type are called *pure states* and are of the form $\mu(A) = \|\Pi_A(p)\|^2$ for some unit vector $p \in \mathcal{H}$. That is, the probability of a subspace A is the length squared of the projection of the vector p . By Gleason's Theorem these measures are the extreme points of the convex set of all quantum probabilities.

We characterize the likelihood orders that can be represented by a quantum measure.

The second main result characterizes the likelihood orders that can be represented by the uniform distribution, defined by $\mu(A) = \frac{\dim(A)}{\dim(\mathcal{H})}$. This is the only quantum measure that obtains a discrete set of values.

We also provide a full characterizing of the partial orders that admit a quantum representation using a duality-like condition, which is less behaviorally justified. However, the general representation problem in terms of axioms that are well-motivated from a behavioral viewpoint remains open.

Subjective analysis of quantum probability has been treated in the literature by several authors. Deutsch ([7], [18]) assumes that an agent assigns a value to any possible outcome of any possible measurement. Deutsch's analysis hinges on what he calls the 'principle of substitutibility', which is similar to the Independence Axiom of von Neumann and Morgenstern ([17]). Barnum et al. ([1]) claimed that Deutsch's proof relies on a tacit symmetry assumption. Deutsch's approach has been defended by Wallace ([18]) who argued that the symmetry assumption is in fact a direct consequence of a 'physicality' assumption saying that an agent assigns the same value to two games instantiated by the same physical process.

While the physical state is the primitive in Deutsch's and Wallace's analysis, which is motivated by the Everett interpretation of quantum mechanics, in ours it is the logic structure – the structure of subspaces of a Hilbert space, as formulated by von Neumann ([16]). Gyntelberg and Hansen ([10]) applied a general event-lattice theory (with axioms that resemble those of von Neumann and Morgenstern) to a similar framework. Pitowsky ([13]) assumed that for every possible measurement the agent has a certain probability over the corresponding outcomes. From some natural axioms he derives the probabilistic structure over quantum mechanics. Caves et al. ([2]) assume that the agent has degrees-of-belief that determine the odds under which he is willing to take a bet. Under the assumption that the agent cannot be attacked by a Dutch book, and an assumption about 'maximal information', they showed that these degrees-of-belief must be given by a pure state.

The main difference between the aforementioned approaches and ours is that we do not assume that the agent has quantitative assignments of probabilities (i.e., numerical degrees-of-belief) to subspaces. Instead, the primitive of our model is a qualitative belief given by the

likelihood order. This study is motivated by Savage's theory of subjective probability ([15]) in classical decision theory.

The approach is behavioral in nature. An agent chooses one alternative from a finite set of choices and an outside observer records the choices made. When the possible alternatives are bets on outcomes of various measurements, the choices of the decision maker reveal his opinion about which outcome of such measurements is more likely. This partial order (i.e., 'being more likely than') over outcomes of different measurements is the primitive of our study. A representation of the partial order by a quantum probability would mean that the agent acts as if he has quantitative degrees-of-belief that obey the rules of quantum probability.

The paper is structured as follows. The next section characterizes the likelihood orders that admit a quantum probability representation in terms of continuity and a duality-like condition, called the cancelation condition. Section 3 introduces the main axioms. Sections 4 and 5 are devoted to the main results: representation by a pure state and by a uniform distribution. Section 6 provides an example of a likelihood order that satisfies the main axioms except for continuity, and cannot be represented by a quantum measure. The paper is concluded with Section 7 that provides some final comments and open problems.

2. THE CANCELATION CONDITION AND CONTINUITY

Let \mathcal{H} be a finite dimensional Hilbert space and let \preceq be a weak order over linear subspaces of \mathcal{H} , that is \preceq is reflexive (i.e., for every A , $A \preceq A$), transitive (i.e., for every A, B, C , if $A \preceq B$ and $B \preceq C$ then $A \preceq C$) and complete (i.e., for every A, B , $A \preceq B$ or $B \preceq A$ or both). We call \preceq the *likelihood order*, and when $A \preceq B$ we say that B is *more likely than* A . Denote by \sim the equivalence relation induced by \preceq (i.e., $A \sim B$ if $A \preceq B$ and $B \preceq A$) and by \prec the corresponding strict order (i.e., $A \prec B$ if $A \preceq B$ and $B \not\preceq A$.)

2.1. The cancelation condition. Cancelation condition (see, for example [8]) is a well-known property of a weak order in the classical framework:

Proposition 1. *Let \preceq be a weak order over subsets of a finite set Ω . For $A \subseteq \Omega$, denote by $\mathbf{1}_A$ the indicator function of A . Then there exists an additive probability measure μ over Ω such that $A \preceq B \leftrightarrow \mu(A) \leq \mu(B)$ for every $A, B \subseteq \Omega$ if and only if the following conditions hold:*

- (1) For every $A \subseteq \Omega$, $\Phi \preceq A$.

- (2) $\Phi \prec \Omega$.
- (3) For every n , if $A_1, \dots, A_n, B_1, \dots, B_n$ are subsets of Ω such that $\sum_{i=1}^n \mathbf{1}_{A_i} = \sum_{i=1}^n \mathbf{1}_{B_i}$ and $A_i \preceq B_i$ for every i , then $A_i \sim B_i$ for every i .

Definition 1. Let μ be a quantum probability measure over \mathcal{H} . We say that μ represents \preceq if $A \preceq B \leftrightarrow \mu(A) \leq \mu(B)$ for every two subspaces A, B of \mathcal{H} .

In the quantum framework, orthogonal projection will replace the indicator functions that appear in Lemma 1.

Definition 2. The likelihood order \preceq satisfies the *cancelation condition* if, for every $2n$ subspaces of \mathcal{H} , $A_1, \dots, A_n, B_1, \dots, B_n$, and n positive numbers α_i , $i = 1, \dots, n$, if $\sum_{i=1}^n \alpha_i \Pi_{A_i} = \sum_{i=1}^n \alpha_i \Pi_{B_i}$ and $A_i \preceq B_i$ for every $i = 1, \dots, n$, then $A_i \sim B_i$ for every $i = 1, \dots, n$.

2.2. Continuity. The cancelation condition by itself is not sufficient to ensure the existence of a representative measure (see Example 2 below.) Similar examples appear in the classical framework, when one tries to extend Proposition 1 to an infinite Ω . In the current framework, in order to obtain a characterization of the likelihood orders that can be represented by a quantum measure, we need the additional assumption that \preceq is continuous w.r.t. the natural topology over subspaces.

Let A be subspace and r be a positive number. Denote by U the unit ball, $U = \{x; \|x\| \leq 1\}$. By $N_r(A)$ we denote the r -neighborhood of A restricted to the unit ball, $\cup\{V(x, r); x \in A \text{ and } x \in U\}$, where $V(x, r)$ is the ball of radius r around x . For two subspaces A and B we denote $\delta_*(A, B) = \inf\{r > 0 : A \cap U \subset N_r(B)\}$ and $\delta^*(A, B) = \inf\{r > 0 : B \cap U \subset N_r(A)\}$. The Hausdorff metric is defined as

$$\delta(A, B) = \max\{\delta^*(A, B), \delta_*(A, B)\}.$$

Definition 3. The likelihood order \preceq is *lower semi-continuous* if, for every subspace B , the set of the subspaces A such that $A \prec B$ is open with respect to δ .

Theorem 1. Let \preceq be a likelihood order. There exists a quantum probability measure that represents \preceq if and only if the following conditions are satisfied:

- (1) $\{0\} \preceq A$ For every subspace A of \mathcal{H} ;
- (2) $\{0\} \prec \mathcal{H}$;
- (3) \preceq is lower semi-continuous;
- (4) \preceq satisfies the cancelation condition.

Proof. Assume first that \preceq is represented by a quantum probability μ . Then by Gleason's Theorem there exists a nonnegative operator T with trace 1 such that $\mu(A) = \text{tr}(\Pi_A T)$ for every subspace A of V . In particular, the function $A \mapsto \mu(A)$ is continuous and therefore the order that it represents is lower semi-continuous. As for the cancellation condition, let $A_1, \dots, A_n, B_1, \dots, B_n$ be subspaces such that $\sum_{i=1}^n \alpha_i \Pi_{A_i} = \sum_{i=1}^n \alpha_i \Pi_{B_i}$ and $A_i \preceq B_i$, where $\alpha_i, i = 1, \dots, n$, are positive numbers. It follows that

$$\begin{aligned} \sum_i \alpha_i \mu(A_i) &= \sum_i \alpha_i \text{tr}(\Pi_{A_i} T) = \text{tr}\left(\left(\sum_i \alpha_i \Pi_{A_i}\right) T\right) = \\ &= \text{tr}\left(\left(\sum_i \alpha_i \Pi_{B_i}\right) T\right) = \sum_i \alpha_i \text{tr}(\Pi_{B_i} T) = \sum_i \alpha_i \mu(B_i). \end{aligned}$$

Since $\mu(A_i) \leq \mu(B_i)$ for every i , it follows that $\mu(A_i) = \mu(B_i)$, which implies $A_i \sim B_i$.

Consider the finite dimensional Hilbert space of Hermitian operators over \mathcal{H} with the inner product of two Hermitian operators S and T being $\text{tr}(ST)$. Denote $\mathcal{C} = \text{Conv}\{\Pi_B - \Pi_A; A \prec B\}$ and $\mathcal{D} = \text{span}\{\Pi_B - \Pi_A; A \sim B\}$. \mathcal{C} is a convex set and \mathcal{D} is a linear subspace. From the cancellation condition it follows that \mathcal{C} and \mathcal{D} are disjoint.

The separation theorem (see, [14]) ensures that there is a non-zero linear function, represented in this case by the Hermitian operator T , such that $\text{tr}(DT) = 0$ for every $D \in \mathcal{D}$ (since \mathcal{D} is a subspace) and $\text{tr}(CT) \geq 0$ for every $C \in \mathcal{C}$.

Since $\text{tr}(DT) = 0$ for $D \in \mathcal{D}$, it follows that $\text{tr}(AT) = \text{tr}(BT)$ if $A \sim B$. Let $A \prec B$. By the definition of \mathcal{C} , $\Pi_A - \Pi_B \in \mathcal{C}$. Due to lower semi-continuity, for B' sufficiently close to B , $A \prec B'$ and therefore $\Pi_A - \Pi_{B'} \in \mathcal{C}$. However, the set $\text{Conv}\{\Pi_A - \Pi_{B'}\}$, where B' is sufficiently close to B , contains $\Pi_A - \Pi_B$ as an interior point. Therefore, $\text{tr}((\Pi_A - \Pi_B)T)$ is strictly positive. We conclude that T represents \preceq .

Finally, for every A , $\text{tr}(\Pi_A T) \geq 0$ since $\{0\} \preceq A$. Therefore, T is positive semidefinite. Moreover, since $\{0\} \prec \mathcal{H}$, it follows that $\text{tr}(T) > 0$. Define $T' = \frac{T}{\text{tr}(T)}$. We obtain that \preceq is represented by $\mu(A) = \text{tr}(T'A)$, T' is positive semidefinite and $\text{tr}(T') = 1$, as desired. \square

The cancellation condition (even in the classical framework) is difficult to justify. It is desirable to derive a probability representation of a likelihood order over linear subspaces from more plausible assumptions.

3. DE-FINETTI'S AND OTHER AXIOMS

The most natural condition is de-Finetti's. When applied to classical probability it states that C is disjoint of $A \cup B$, then B is preferred to A iff $B \cup C$ is preferred to $A \cup C$. In the quantum framework it takes following form:

de-Finetti's Axiom: For every linear subspaces A, B, C of \mathcal{H} , if $A \perp C$ and $B \perp C$, then $A \preceq B$ iff $A + C \preceq B + C$.

In the classical framework it easily follows from **de-Finetti's axiom** that if $A \preceq B$ then $B^c \preceq A^c$. In the quantum framework, we need to require it explicitly.

Negation: For every two linear subspaces A, B of \mathcal{H} , if $A \preceq B$ then $B^\perp \preceq A^\perp$.

As illustrated by the following example there might be weak orders that satisfy **de-Finetti's axiom** and not **Negation**.

Example 1. Let \mathcal{H} be \mathbb{R}^2 . Any monotonic (w.r.t set inclusion) weak order on \mathcal{H} satisfies **de-Finetti's axiom** but it might not satisfy **Negation**. Let \succeq' be such a weak order. As for a higher dimensional Hilbert space, let \mathcal{H} be \mathbb{R}^3 . Let p be the northern pole of the unit ball, E be the equator, and define $\mu(u) = |\langle p, u \rangle|^2$ for any unit vector u .

Define \preceq as follows: If A and B are two subspaces of different dimensions, then $A \succ B$ if the dimension of A is greater than that of B . If $A = \text{span}\{u\}$ and $B = \text{span}\{v\}$, where u and v are unit vectors, then $A \succeq B$ either when $u \notin E$ and $\mu(u) \geq \mu(v)$ or when $u, v \in E$ and $u \succeq' v$. Finally, if A and B are two-dimensional subspaces, then $A \succeq B$ either when $p \notin B$ and $\|\Pi_A(p)\|^2 > \|\Pi_B(p)\|^2$ or when $p \in A \cap B$ and $A \cap E \succeq' B \cap E$. The weak order \preceq preserves **de-Finetti's axiom** but since \succeq' does not preserve **Negation** on E , so \preceq does not on \mathcal{H} .

We will also need the obvious assumption that any subspace is as preferred as the zero-dimensional one. Formally,

Monotonicity: For every subspace A of \mathcal{H} , $\{0\} \preceq A$.

Note that **Monotonicity** and **de-Finetti's axiom** together imply that if $A \subseteq B$ then $A \preceq B$ for every pair A, B of subspaces. Thus, \preceq is monotonic with respect to set inclusion.

In the sequel, we will say that a weak order \preceq satisfies the *standard assumptions* if it satisfies **de-Finetti's axiom**, **Negation** and **Monotonicity**.

Do the standard assumptions guarantee that \preceq can be represented by a measure? The following is a counterexample.

Example 2. Let μ_1, μ_2 be two different quantum probability measures over \mathcal{H} and define the *lexicographic order* induced by μ_1 and μ_2 as follows. $A \preceq B$ if either $\mu_1(A) < \mu_1(B)$ or $\mu_1(A) = \mu_1(B)$ and $\mu_2(A) < \mu_2(B)$. Then, \preceq satisfies the standard assumptions. Furthermore, it satisfies the cancelation condition.

The lexicographic order cannot be represented by a measure since it lacks the following property:

Separability: There is a countable set of subspaces, \mathcal{A} , such that for any two subspaces B and C such that $B \prec C$, there is $A \in \mathcal{A}$ that satisfies $B \preceq A \preceq C$.

As indicated by Debreu ([5]), **Separability** is necessary for \preceq in order to be represented by a real function (not necessarily a measure).

4. PURE STATES IN \mathbb{R}^3

The most important probabilities from the physical point of view are those of the form $\mu(A) = \|\Pi_A(p)\|^2$ for some unit vector $p \in A$. These distributions are sometimes called *pure states*. It follows from Gleason's Theorem that pure states are the extreme points of the convex set of all quantum probabilities.

It is clear that if μ is a pure state and \preceq is the induced likelihood order, then the one-dimensional subspace spanned by p is equivalent (under \sim) to \mathcal{H} . In this section we prove the inverse statement. We say that \preceq is *non-trivial* if there exists a subspace A such that $\{0\} \prec A$.

Theorem 2. *Let \preceq be a weak order over subspaces of a finite dimensional real-Hilbert space that satisfies the standard assumptions and **Separability**. Assume that there exists a one-dimensional subspace P such that $P \sim \mathcal{H}$. Let p be a unit vector in P . If \preceq is non-trivial, then \preceq is represented by the pure state $\mu(A) = \|\Pi_A(p)\|^2$.*

Proof. Let E be the orthogonal complement of $P = \text{span}\{p\}$. By **Negation**, since $P \sim \mathcal{H}$ it follows that $E \sim \{0\}$. Let A be a subspace of \mathcal{H} , A_0 be $A_0 = A \cap E$ and A_1 be the one-dimensional subspace of \mathcal{H} that is spanned by $\Pi_A(p)$ (the orthogonal projection of p over A .) Then, since $A_0 \subset E$, it follows from **Monotonicity** that $A_0 \sim 0$. Since $A_1 \perp$

A_0 and $A = A_1 + A_0$, it follows from **de-Finetti's axiom** that $A \sim A_1$. Thus, the likelihood order \preceq is determined by its restriction to one-dimensional subspaces. Moreover, since $\mu(A) = \mu(A_1)$, it is sufficient to prove that the likelihood order over one-dimensional subspaces is represented by μ . Slightly abusing notation, we will identify a unit vector u in \mathcal{H} with the one-dimensional subspace spanned by u . With this convention, for every unit vector u , $\mu(u) = |\langle p, u \rangle|^2$.

Let u, v be two unit vectors. We need to show that $u \preceq v$ iff $|\langle p, u \rangle|^2 \leq |\langle p, v \rangle|^2$. By looking at the three-dimensional space \mathcal{H}_{uv} spanned by p, u, v , with its two-dimensional subspace $\mathcal{H}_{uv} \cap E$ we can assume w.l.o.g. that $\dim \mathcal{H} = 3$. In this case the theorem will follow directly from the following proposition. Let S^2 be the unit sphere in \mathbb{R}^3 . We say that \preceq is *uniform* if all the one-dimensional subspaces are equivalent.

Proposition 2. *Let \preceq be a weak order over \mathbb{R}^3 that satisfies the standard assumptions, and such that the restriction of \preceq to one-dimensional subspaces is separable. Assume that \preceq is not uniform and it attains its minimum over S^2 at m . Furthermore, assume that there exists a two-dimensional subspace E such that $m \sim u$ for every $u \in E$. Let $p \in S^2$ such that $p \perp E$. Then, for every pair $u, v \in S^2$, $u \preceq v$ iff $|\langle p, u \rangle|^2 \leq |\langle p, v \rangle|^2$.*

Proof of Proposition 2. The proof of the proposition is broken into five claims. As usual, we identify elements of S^2 with their corresponding one-dimensional subspaces.

Claim 1. Let $q, r \in S^2$ be such that $q \preceq r$. If q' and r' are, respectively, the orthogonal complements of q and r in the plane $\text{span}\{q, r\}$, then $r' \preceq q'$.

Proof. Let $n \in S^2$ such that $n \perp \text{span}\{q, r\}$. Then $q^\perp = \text{span}\{n, q'\}$ and $r^\perp = \text{span}\{n, r'\}$. By **Negation**, $r^\perp \preceq q^\perp$. By **de-Finetti's axiom**, $r' \preceq q'$. \square

Claim 2. Let $u_1, u_2 \in S^2$ be orthogonal vectors such that $u_1 \sim u_2 \sim m$. If $u \in \text{span}\{u_1, u_2\}$, then $u \sim m$.

Proof. Note that u_2 is the orthogonal complement of u_1 in $\text{span}\{u_1, u_2\}$. Let u' be the orthogonal complement of u in $\text{span}\{u_1, u_2\}$. Then $u_1 \sim m \preceq u'$. By Claim 1, $u \preceq u_2 \sim m$. Since $m \preceq u$, it follows that $u \sim m$. \square

Claim 3. Assume that there exists an orthogonal triple u_1, u_2, u_3 such that $m \sim u_1 \sim u_2 \sim u_3$. Then, \preceq is uniform.

Proof. Let $v \in S^2$. Then there exists $u \in \text{span}\{u_1, u_2\}$ such that $v \in \text{span}\{u, u_3\}$. By Claim 2, $u \sim m$. But $u \perp u_3$ and therefore again by Claim 2, $v \sim m$. \square

For $q \in S^2$ we denote by $\theta(p, q)$ the angle between p and q . Thus $0 \leq \theta(p, q) \leq \pi$ and $\cos \theta(p, q) = \langle p, q \rangle$. Let $N_p = \{q \in S^2 \mid 0 < \theta(p, q) < \pi/2\}$ be the *northern hemisphere* relative to p , and $E_p = \{q \in S^2 \mid \theta(p, q) = \pi/2\} = E \cap S^2$ be the *equator* relative to p . Let $q \in N_p$. Among the great circles which pass through q there is a unique one that intersects E_p in vector x orthogonal to q . We follow Gleason (1957) and denote this circle by $\text{EW}(q)$. Note that q is the northern most point in $\text{EW}(q)$ and that $\text{EW}(q)$ is tangent to the latitude circle of q . We will need the following lemma, that appears in ([12]) (see also [4]).

Piron's Lemma. Let $q, r \in N_p$ such that $\theta(p, q) < \theta(p, r)$; then there exists a finite sequence $q = q_0, q_1, \dots, q_n = r$ of points in N_p such that $q_{i+1} \in \text{EW}(q_i)$.

Claim 4. Under the assumption of Theorem 2, if $q, r \in N_p$ and $\mu(r) < \mu(q)$, then $r \prec q$.

Proof. Note that $\mu(q) > \mu(r)$ iff $\theta(p, q) < \theta(p, r)$.

Let $q \in N_p$ and $q_1 \in \text{EW}(q)$. Let $q'_1 \in \text{EW}(q)$ be the orthogonal complement of q_1 in the plane of $\text{EW}(q)$, and $q' \in E_p$ be the orthogonal complement of q in $\text{EW}(q)$. Since $q' \sim m \preceq q'_1$, it follows from Claim 1 that $q_1 \preceq q$. Moreover, $q_1 \sim q$ only if $q'_1 \sim m$. By induction it follows from Piron's Lemma that $r \preceq q$. Furthermore, $r \sim q$ only if there exists $z \in N_p$ such that $z \sim m$. We prove that in this case \preceq is uniform, which is excluded by assumption. This will complete the proof.

Note that for every y such that $\theta(p, z) < \theta(p, y) < \pi - \theta(p, z)$, $m \preceq y \preceq z \sim m$. Thus, all the vectors in the band below z are equivalent to m . We now show that one can find another point p' , such that $x' \sim m$ for every $x' \in E_{p'}$ and $\theta(p', z) = \frac{1}{2}\theta(p, z)$, and thus obtaining a wider band. By iterating this argument one can get wider and wider bands until one obtains a band that is wide enough to contain three orthogonal vectors. By Claim 3, it would imply that \preceq is uniform.

Let p' be point in $\text{span}\{p, z\}$ for which $\theta(p, p') = \theta(p', z) = \frac{1}{2}\theta(p, z)$. It follows that, for every $x' \in E_{p'}$, $\frac{3}{2}\theta(p, z) < \theta(p, x') < \pi - \frac{3}{2}\theta(p, z)$. Thus, $E_{p'}$ is entirely contained in the band defined by p and z and therefore $E_{p'} \sim m$. \square

Claim 5. If $q, r \in N_p$ and $\mu(q) = \mu(r)$, then $q \sim r$.

Proof. We know from Claim 4 that for $q, r \in N_p$ such that $\mu(q) < \mu(r)$, $r \prec q$. Now suppose that there exist, for some α , $0 < \alpha < \frac{\pi}{2}$, vectors $q_0, r_0 \in N_p$ such that $q_0 \prec r_0$ and $\mu(q_0) = \mu(r_0) = \alpha$. Let $Q = \{q \in S^2; \mu(q) = \alpha, q \prec r_0\}$ and $R = \{r \in S^2; \mu(r) = \alpha, q_0 \prec r\}$. Then at least one of the sets Q, R must be uncountable. Assume w.l.o.g. that Q is uncountable. For every $q \in Q$, let q', r' be the orthogonal complements of q, r_0 resp. in $\text{span}\{q, r_0\}$. It follows from Claim 1 that $r' \prec q'$. Notice moreover, that $\mu(q') = \mu(r') = 1 - \alpha - \mu(n(q))$ where $n(q) \in S^2$ is orthogonal to $\text{span}\{q, r_0\}$. Since $\theta(p, n(q)) = \cos \sqrt{\mu(n(q))}$ increases as q approaches r_0 along the latitude circle of r_0 , we get uncountable set of pairs (r', q') such that $\mu(r') = \mu(q')$, but $r' \prec q'$ with different values of μ for different pairs. This, together with Claim 4 contradicts separability. \square

From Claims 4 and 5 it follows that $\mu(q) \leq \mu(r)$ iff $q \preceq r$, and therefore the proof of Proposition 2 is complete.

Back to the proof of Theorem 2. By assumption, \preceq is non-trivial. Therefore, there exists a subspace A which is strictly more likely than $\{0\}$. Suppose that A is spanned by the orthogonal vectors u_1, \dots, u_k .

Claim 6. At least one u_i is strictly more likely than $\{0\}$.

Proof. Otherwise, $u_i \sim \{0\}$ for every $i = 1, \dots, k$. By **de-Finetti's axiom** $\text{span}\{u_1, u_2\} \sim \text{span}\{u_2\} \sim \{0\}$. By successively adding the u_i 's and by using **de-Finetti's axiom** one obtains that $A \sim \{0\}$, in contradiction with the assumption. \square

By Claim 6 we can assume that there is a vector $x \in S^2$ such that $x \succ \{0\}$. Let $y \in \text{span}\{x, p\} \cap E$. Since $y \in E$, $y \sim \{0\}$. Furthermore, $y \perp p$. Let x' be the orthogonal complement of x in $\text{span}\{x, p\}$. Since $x' \succ y$, by Claim 1 $x \preceq p$. As \preceq is an order, $\{0\} \prec x \preceq p$, and thus, $\{0\} \prec p$. This implies that \preceq , when restricted to \mathcal{H}_{uv} , is not uniform, as assumed by Proposition 2. This enables us to use this proposition in order to complete the proof of Theorem 2. \square

Remark 1. No sort of continuity is assumed in Theorem 2. Nevertheless, \preceq is represented by a measure and is therefore continuous.

5. THE UNIFORM MEASURE

The only quantum probability measures over a finite dimensional Hilbert space \mathcal{H} which receives discrete values is given by the *uniform measure*, $\mu(A) = \frac{\dim(A)}{\dim(\mathcal{H})}$. It turns out that this is the case characterized by the property that all one-dimensional subspaces are equally likely. Formally,

Proposition 3. *Let \preceq be a weak order over subspaces of a finite dimensional Hilbert space that satisfies **de-Finetti's axiom**. If all one-dimensional subspaces are equivalent, then either \preceq is trivial (i.e. $\{0\} \sim A$ for every subspace A of \mathcal{H}) or \preceq is represented by the uniform measure.*

Proof. Assume that every one-dimensional subspace is equivalent to some one-dimensional subspace, say, m . If A_1, A_2 are two-dimensional such that $A_1 \cap A_2$ is one-dimensional, we can assume that $A_1 = \text{span}\{a_0, a_1\}$ and $A_2 = \text{span}\{a_0, a_2\}$ where $a_0 \perp a_1$ and $a_0 \perp a_2$. Since $a_1 \sim a_2 \sim m$ we get, by **de-Finetti's axiom**, that $A_1 \sim A_2$. If $A_1 \cap A_2 = \{0\}$, we can find a two-dimensional subspace A' such that $A_1 \cap A'$ and $A_2 \cap A'$ are one-dimensional. Therefore $A_1 \sim A' \sim A_2$. Thus every two-dimensional subspaces are equivalent. By a similar argument, two subspaces of the same dimension are equivalent.

Finally, if $\{0\} \sim m$, it follows by **de-Finetti's axiom** that $\{0\} \sim \mathcal{H}$. If $0 \prec m$, then again by **de-Finetti's axiom**, if $A \perp m$ and $A' = A + m$ then $A \prec A'$. Using the equivalence of two subspaces with the same dimension, it follows that if $\dim(A') = \dim(A) + 1$, then $A \prec A'$ and therefore \preceq is represented by $\mu(A) = \frac{\dim(A)}{\dim(\mathcal{H})}$. \square

The following nontrivial fact about quantum probabilities follows from Gleason's Theorem:

Proposition 4. *Let \mathcal{H} be a finite-dimensional Hilbert space and μ be a quantum probability over \mathcal{H} . Assume that there exist one-dimensional subspaces (not necessarily orthogonal) u_1, \dots, u_n of \mathcal{H} such that $\mathcal{H} = u_1 + \dots + u_n$ and $\mu(u_1) = \dots = \mu(u_n) \leq \mu(x)$ for every one-dimensional subspace x of \mathcal{H} . Then, μ is the uniform measure.*

We show that this proposition is a consequence of the standard assumptions, with the additional assumption that \preceq is continuous over one-dimensional subspaces.

Definition 4. The likelihood order \preceq is *continuous over one-dimensional subspaces* if for every unit vector v the sets $\{u; u \prec v\}$ and $\{u; v \prec u\}$ are open.

We note that if \preceq is continuous over one-dimensional subspaces then its restriction to one-dimensional subspaces is also separable. Indeed, let $D \subseteq S^2$ be a countable dense set w.r.t. the Euclidean topology of S^2 . For every $u, v \in S^2$ such that $u \prec v$, let $U = \{u' \in S^2 | u \prec u'\}$ and $V = \{v' \in S^2 | v' \prec v\}$. Since $S^2 = U \cup V$ and S^2 is connected, $U \cap V \neq \emptyset$. As D is dense, there exists $d \in D$ such that $d \in U \cap V$. Thus, $u \prec d \prec v$.

We state the result in \mathbb{R}^3 . It can easily be extended to every finite-dimensional Hilbert space.

Theorem 3. *Let \preceq be a weak order over \mathbb{R}^3 that satisfies the standard assumptions. Assume that \preceq is continuous over one-dimensional subspaces. If u_1, u_2, u_3 is a basis (not necessarily orthogonal) that satisfies $u_1 \sim u_2 \sim u_3 \sim m$, where m is a minimum of \preceq , then $x \sim m$ for every $x \in S^2$.*

The theorem is proved in a few steps. Denote by M a maximum of \preceq .

Claim 7. Let $u, v \in S^2$ such that $u \sim m$ and $v \sim M$. Let u', v' be the orthogonal complements of u, v in $\text{span}\{u, v\}$, respectively. Then, $u' \sim M$ and $v' \sim m$.

Proof. Since $m \sim u \preceq v'$ it follows from Claim 1 that $v \preceq u'$. But $v \sim M$ and M is a maximum. Therefore $u' \sim M$. By a similar argument $v' \sim m$. \square

Claim 8. If $u_1 \perp u_2 \in S^2$ and $m \sim u_1 \sim u_2$, then either all one-dimensional subspaces are equivalent or \preceq is represented by a pure state.

Proof. Let $E = \text{span}\{u_1, u_2\}$. By Claim 2, $u \sim m$ for every $u \in E$. By Proposition 2, either \preceq is trivial or \preceq is represented by a pure state. \square

Claim 9. If $u_1 \neq \pm u_2 \in S^2$, $M \sim u_1 \sim u_2$ and $p \perp u_1, u_2$, then $p \sim m$.

Proof. Let $x \in N_p$ be such that $x \sim m$ and $\theta(p, x)$ is minimal. If $x \neq p$ then x cannot be orthogonal to both u_1 and u_2 . Assume therefore w.l.o.g. that $\langle x, u_1 \rangle \neq 0$. Let u'_1 be the orthogonal complement of u_1 in the plane $\text{span}\{x, u_1\}$. By Claim 7, $u'_1 \sim m$. Moreover $\theta(p, u'_1) < \theta(p, x)$ since $x \in EW(u'_1)$. This contradicts the choice of x . It therefore follows that $x = p$, meaning that $p \sim m$. \square

Claim 10. If m and M are any minimal and maximal elements in S^2 and $m \prec M$ (i.e., \preceq is not trivial), then $m \perp M$.

Proof. Assume the contrary. Let a be the orthogonal complement of m in $\text{span}(m, M)$. By Claim 7, $a \sim M$. Let p satisfy $p \perp M$ and $p \perp a$. By Claim 9, $p \sim m$. However, since $m \in \text{span}(M, a)$, $p \perp m$. Therefore it follows from Claim 8 that \preceq is represented by a pure state, in which case the claim holds. \square

We now turn to the proof of the Theorem 3. Let M be a maximal element. If \preceq is not trivial then from the last claim it follows that $M \perp u_i$ for every i . This is impossible since since u_1, u_2, u_3 are linear independent, and the proof is complete.

Remark 2. We do not know whether Theorem 3 holds true without the assumption that \preceq is continuous over one-dimensional spaces. The proof hinges on this assumption in two ways. First, in that \preceq attains a minimum and a maximum. Second, in Claim 9 x is chosen so that among all $x \sim m$, $\theta(p, x)$ is minimal. While we could explicitly assume that \preceq attains a minimum and a maximum, we could not dispose of the continuity assumption in the proof of Claim 9.

6. A COUNTEREXAMPLE

In this section we present an example of a separable (though not continuous) weak order over subspaces of \mathbb{R}^3 that satisfies the standard assumptions but does not admit a representation via a quantum measure. We need the following two lemmas.

Lemma 1. *Let \preceq be a weak order over one-dimensional subspaces of \mathbb{R}^3 such that for every two-dimensional subspace U of \mathbb{R}^3 and every one-dimensional subspaces u, v of U one has $u \preceq v \iff v' \preceq u'$ where u', v' are the orthogonal complements of u, v resp. in U , then \preceq can be extended to a weak order over \mathbb{R}^3 that satisfies the standard assumptions.*

Proof. We define \preceq as follows. Let U, V be two subspaces of \mathbb{R}^3 . If $\dim(U) < \dim(V)$ then $U \prec V$. If $\dim(U) = \dim(V) = 2$ then $U \preceq V$ iff $V^\perp \preceq U^\perp$. **Negation** is obviously satisfied. As for **de-Finetti's axiom**, let u, v be two different one-dimensional subspaces and x be the one-dimensional subspaces such that $x \perp u, v$. Let u', v' be the orthogonal complements of u, v in $u + v$. Then $(x + u)^\perp = u'$ and $(x + v)^\perp = v'$. Since, by the assumption of the lemma $v' \preceq u'$, it follows by definition of \preceq that $x + u \preceq x + v$. \square

The second lemma states that if \preceq is represented by a probability measure, then the order over one-dimensional subspaces of a fixed two-dimensional subspace U has a very specific form. As usual we identify one-dimensional subspaces with unit vectors. If S^1 is the unit circle of U , the lemma essentially says that either all elements of S^1 are equivalent, or there is a single maximal element $x \in S^1$ that satisfies $y_1 \preceq y_2$ iff y_2 is closer than y_1 to x .

Lemma 2. *Let μ be a probability measure over \mathbb{R}^3 and \preceq the corresponding weak order over subspaces. Let U be a two-dimensional subspace of \mathbb{R}^3 . Then, either all one-dimensional subspaces of U are equivalent, or there exists unit vector $x \in U$ such that for every pair y_1, y_2 of unit vectors $y_1 \preceq y_2$ iff $|\langle x, y_1 \rangle| \leq |\langle x, y_2 \rangle|$.*

Proof. By Gleason's Theorem, there exists a positive semidefinite operator T such that $\mu(A) = \text{tr}(\Pi_A T)$. Consider the operator $\Pi_U T \Pi_U$. This is a positive semidefinite operator. Its spectral decomposition is of the form

$$\Pi_U T \Pi_U = \alpha \Pi_x + \beta \Pi_{x'},$$

where x, x' are orthogonal eigenvectors in U with corresponding eigenvalues α, β such that $\alpha + \beta = 1$. We assume that $\alpha \geq \beta$. It follows that for every unit vector y in U ,

$$\begin{aligned} \mu(y) &= \text{tr}(\Pi_y T) = \text{tr}(\pi_y \Pi_U T \Pi_U) = \\ &= \alpha |\langle y, x \rangle|^2 + \beta |\langle y, x' \rangle|^2 = \beta + (\alpha - \beta) |\langle y, x \rangle|^2. \end{aligned}$$

Thus, if $\alpha = \beta$ then all $y \in S^2 \cap U$ are equivalent. If $\alpha > \beta$ then $\mu(y)$ is a monotonic function of $|\langle y, x \rangle|$. \square

Example 3. Let \succeq' be a weak order on one-dimensional subspaces of \mathbb{R}^2 that satisfies the condition of Lemma 1 but not the condition of Lemma 2. Define \preceq on one-dimensional subspaces of \mathbb{R}^3 as follows: Let p be the northern pole of the unit sphere in \mathbb{R}^3 . Let u and v be unit vectors, then $u \preceq v$ either when $u \notin E$ and $\mu(u) \geq \mu(v)$ or when $u, v \in E$ and $u \succeq' v$. By Lemma 1 \prec can be extended to a weak order over \mathbb{R}^3 that satisfies the standard assumptions. However, since the condition of Lemma 2 is not satisfied, \preceq cannot be represented via a quantum measure.

7. FINAL COMMENTS AND OPEN PROBLEMS

7.1. Representation and continuity. In Gleason's Theorem ([9]) continuity is not assumed and is a consequence of the existence of a frame function. When the primitive of the model is a likelihood order, matters are different. The likelihood order in Example 3 satisfies **de-Finetti's axiom**, **Negation**, **Monotonicity** and **Separability** and cannot be represented by a quantum measure. This order, which is not continuous, suggests that continuity must be explicitly assumed and cannot be derived from more plausible assumptions.

The question whether every continuous likelihood order which satisfies **de-Finetti's axiom**, **Negation**, **Monotonicity** and **Separability** can be represented by a quantum measure is still open.

7.2. Partial representation.

Definition 5. We say that μ *partially represents* \preceq if $A \preceq B \longrightarrow \mu(A) \leq \mu(B)$ for every two subspaces A, B of \mathcal{H} .

It turns out (we state without a proof) that if \preceq satisfies the cancellation condition, then there exists a quantum probability measure that partially represents \preceq . Also, from the proof of Theorem 2 it follows that, if there exists a one-dimensional subspace p , such that $p \sim \mathcal{H}$, then (without assuming separability) \preceq admits a partial representation by a pure state.

7.3. Qualitative additivity and discrete orders. Gleason's Theorem implies that the only quantum probability measure which obtains a discrete set of values is the uniform measure. The question arises whether the same is true for likelihood orders. We say that \preceq is *discrete* if the restriction of \sim to one-dimensional subspaces has only finitely many equivalence classes. For instance, if \preceq is represented by the uniform probability, then its restriction to one-dimensional subspaces has only one equivalence class.

Kochen-Specker's Theorem ([11]) actually refers to likelihood orders whose restriction to one-dimensional subspaces have precisely two equivalence classes. In order to prove this result using likelihood orders terms only, one needs to strengthen **de-Finetti's axiom** and **Negation**. The following axiom is a consequence of **de-Finetti's axiom** in the classical case, but not in the quantum set-up.

Qualitative additivity: Let A_1, A_2, B_1, B_2 be linear subspaces of \mathcal{H} such that $A_1 \perp A_2$ and $B_1 \perp B_2$. If $A_i \preceq B_i$, $i = 1, 2$ then $A_1 \oplus A_2 \preceq B_1 \oplus B_2$. Furthermore, one strict likelihood on the former inequalities implies strict likelihood in the later inequality.

Suppose that A_1 and A_2 are orthogonal and the same for the B_i 's. **Qualitative additivity** states that, if the A_i 's are at least as likely as B_i 's, then the subspace spanned by the A_i 's is at least as likely as that spanned by B_i 's. That is, adding a more likely subspace to a subspace which is already more likely, cannot make the outcome less likely.

Suppose that \preceq is defined over \mathbb{R}^3 and there are only two equivalence classes of one-dimensional subspaces, say, green and red. If \preceq satisfies **Qualitative additivity**, then in any orthogonal triple there is the same number of green representatives, and moreover, a two dimensional subspace spanned by uni-colored vectors contains only vectors of the same color. These are precisely the terms of Kochen-Specker's Theorem ([11]). It states that there exists no likelihood order that satisfies **Qualitative additivity** and has precisely two equivalence classes of one-dimensional spaces.

This result suggests that the only discrete likelihood order that satisfies **Qualitative additivity** is that induced by the uniform measure.

REFERENCES

- [1] Barnum H., C.M. Caves, J. Finkelstein, C.A. Fuchs and R. Schack (2000) Quantum probability from decision theory? *Proc. R. Soc. Lond. A* 456, 1175-1182.
- [2] Caves C.M., C.A. Fuch and R. Schack (2002) Quantum probabilities as Bayesian probabilities. *Physical Review A*. 65, 022305.
- [3] Chevalier G., A. Dvurecenkij and K. Svozil (2000) Piron's and Bell's Geometric Lemmas and Gleason's Theorem. *Foundations of Physics* 30, 1737-1755.
- [4] Cooke, R., M. Keane, and W. Moran (1985) An Elementary proof of Gleason's Theorem. *Proc. Cambr. Phil. Soc* 98, 117-128.
- [5] Debreu, G. (1954) Representation of a preference ordering by a numerical function. In *Decision Processes*, Thrall, R.M., Coombs, C.H. and Davis, R.L., Editors, Wiley, New York, 159-165.
- [6] de-Finetti, B. (1974-5) *Theory of Probabilty*, (translation of 1970 book) 2 volumes, New York: Wiley.
- [7] Deutsch, D. (1999) Quantum theory of probability and decisions. *Proc. R. Soc. Lond. A* 455, 3129-3137.
- [8] Fishburn, P. (1999) Preference structures and their numerical representations. *Theoret. Comput. Sci.* 217, 359-383.
- [9] Gleason, A.M. (1957) Measures on the closed subspaces of a Hilbert Space. *Journal of Mathematics and Mechanics* 6, 885-893.
- [10] Gyntelberg, J. and F. Hansen (2004) Expected utility theory with "small worlds", <http://www.econ.ku.dk/wpa/pink/2004/0420.pdf>
- [11] Kochen, S. and E. P. Specker (1967) The problem of hidden variables in quantum mechanics. *Journal of Mathematics and Mechanics* , 17, 59-87.
- [12] Piron C. (1976), *Foundations of Quantum Physics*, Addison-Wesley, Mass.
- [13] Pitowsky, I. (2003) Betting on the outcomes of measurements. *Studies In History and Philosophy of Modern Physics* 34, 395-414.
- [14] Rockefeller, R.T., (1970) *Convex Analysis*, Princeton University Press, Princeton.
- [15] Savage, L.J. (1954) *The Foundations of Statistics* London: John Wiley and Sons.
- [16] von Neumann, J. (1955) *Mathematical Foundations of Quantum Mechanics*. Princeton: Princeton University Press. (German Edition 1932).
- [17] von Neumann, J. and O. Morgenstern (1944) *Theory of games and economic behavior*, Princeton University Press.
- [18] Wallace, D. (2003) *Studies In History and Philosophy of Modern Physics* 34, 415-439.