

Allocation in Multi-Agenda Disputes: a Set-Valued Games Approach*

Ehud Lehrer[†] and Roei Teper[‡]

January 9, 2019

Abstract

We study allocation problems when agents negotiate across different agendas. Unlike existing papers on multi-agenda disputes, we consider environments in which resources are constrained and investing (say, time or effort) in one agenda reduces the ability to invest in other agendas. In order to analyze such a scenario, we introduce a class of cooperative games, referred to as set-valued – SVG. The value of each coalition in an SVG is a subset of payoff vectors. Each vector is associated with a specific distribution of the resources that the coalition may allocate across the different agendas. In this environment we introduce and analyze the notion of the core. We show that the core of an SVG allows for more cooperation opportunities and exchanging favors than existing cooperative multi-agenda models. Interestingly, proving this relies on a general notion of a comparative advantage. It is shown that the classical core characterization, resorting to duality, does not hold in the current setup. We then extend the balancedness condition that guarantees a non-empty core in a wide range of games.

*The authors wish to thank Gabi Gayer, Michel Grabisch, Dotan Persitz, two anonymous referees and an editor for helpful comments.

[†]School of Mathematical Sciences, Tel Aviv University, Tel Aviv 69978, Israel and INSEAD, Bd. de Constance, 77305 Fontainebleau Cedex, France. e-mail: lehrer@post.tau.ac.il. This research was supported in part by ISF through Grant #963/15.

[‡]Department of Economics, University of Pittsburgh, 230 South Bouquet St., Pittsburgh, 15260, USA. email: rteper@gmail.com.

Keywords: Set-valued games, core, balancedness, logrolling.

JEL Classification: C71, D80, D81, D84

1 Introduction

Typically a (cooperative) *game* measures the ‘worth’ of any subgroup of individuals. This value can be interpreted as the productivity of the subgroup under consideration when investing time and effort in a particular enterprise, its social or political power, etc. Often times though there are multiple enterprises that could be invested in while the resources, say time, are constrained. In such scenarios there are many different time allocations to each of the different projects, each of which resulting in a different outcome. It seems that if one wishes to model such tradeoffs without pre-committing to a specific time allocation, and thereby to a specific level of production in each project, one needs to consider a more general notion of a game than the classical ones.

In this paper we introduce and study the concept of *set-valued games*: each subgroup of individuals is associated with a *set of real valued vectors*. The set of vectors associated with a subgroup is *all* production possibilities across the different enterprises. Note that this approach does not take a stand on the aggregation of (or preferences over) payoffs across the different agendas as in existing work on multi-agenda disputes. We seek to study the primitive, as opposed to the reduced form, and wish to consider a model that is robust to the aggregation process. As in the classical theory, we address the issue of allocation. The appropriate notion of the core of a set-valued game is defined and analyzed.

Vector games have been studied ever since Lind [7]. Set-valued games and their cores were studied as non-transferable utility games (see, e.g., Predtetchinski and Herings [9]), and more generally as in the current manuscript (see Voorneveld and van den Nouweland [8] and Fernandez et al. [4]). The current manuscript introduces a new model and interpretation to allocation in multi-agenda disputes and studies its relation and distinction to existing economic theories on this topic. We show that our approach to multi-agenda disputes raises interesting results relative to the standard ones in cooperative games.¹ First, the notion of the core presented here provides more

¹Work on the topic of multi-dimensional bargaining exists also in the literature of noncooperative

opportunities for cooperation, and exchange of “power” across the different agendas, among the individuals. Second, we show that the classical Bondareva-Shapley result of core non-emptiness [10, 2] does not hold in our model. The well-known balancedness condition is sufficient for core non-emptiness, but is not necessary.

For the sake of clarity of our results, consider a grand group of N individuals that have to invest in k enterprises. A set-valued game (henceforth SVG) is a function \mathbf{v} that associates a subset $\mathbf{v}(S) \subseteq \mathbb{R}_+^k$ for every subgroup (or coalition) S of individuals out of N . Under the classical notion of a game, $k = 1$ and $\mathbf{v}(S)$ is a singleton. The interpretation behind an SVG \mathbf{v} is the following. Every $x \in \mathbb{R}_+^k$ is a vector of production levels for each of the k projects. A coalition can produce the bundle $x = (x_1, \dots, x_k)$ if by distributing their limited resources (say one unit of time) across the different projects, they can accomplish x_ℓ of project ℓ , for every $\ell = 1, \dots, k$. If $x \in \mathbb{R}_+^k$ is indeed produceable by coalition S (in one time unit), then $x \in \mathbf{v}(S)$. A member of the *core* of an SVG is a payoff for each player for each of the k enterprises, that is feasible (that is, in $\mathbf{v}(N)$), and such that no coalition can deviate and do better by itself in every one of the k enterprises.²

We start by paying specific attention to a special class of SVGs called *multi-game based*. An SVG in this class is a convex span of k (classical) games. Formally, letting v_ℓ be the ℓ th game in the base of the SVG \mathbf{v} , the value of a coalition S is all vectors for which the ℓ th production component is $\alpha_\ell v_\ell(S)$ (where the α_ℓ 's are non-negative and sum to 1). This corresponds to production output being linear in effort. For this class of games, it is interesting to see the relation between our solution concept and the cores of the individual games (which are the base of the SVG), and between our notion of a core and existing approaches to allocation in multi-agenda disputes, in which the payoffs across agendas are aggregated uniformly (in particular, Bloch and de Clippel [3] and Gayer and Persitz [5]).

We provide the following results. First, we show that it is possible that while the cores of all base games, and of the sum of these games, are all empty, and thus the solution concepts in both Bloch and de Clippel and Gayer and Persitz are empty, the core of the SVG itself is not empty. We further show that if the cores of all base games

games. See [3] for a review.

²Note that this is consistent with the approach that no particular aggregation of the payoffs across projects is considered.

are non empty, then the convex span of these cores are a subset of the core of the SVG. In addition, this containment could be strict. In particular, we prove that whenever the cores of the base games all consist of a single allocation, then there is always a possibility of logrolling among several individuals that is not possible when resorting to previous approaches.³ These results indicate that the approach and solution concept presented here are conceptually different than the existing approaches, and allow for more cooperation.

We then provide a non Bondareva-Shapley result: an appropriate definition of balancedness, that serves as a characterization of core non-emptiness in the classical cooperative games setup,⁴ is a sufficient condition for the core of an SVG being non-empty, but it is not necessary. An example is provided in which an SVG has a non-empty core while it is not balanced. We point to the fact that a characterization in the current setup cannot rely on straightforward convexity and duality considerations as appearing in Bondareva-Shapley. Finally, we generalize the Bondareva-Shapley condition in a proper way, and show that it 1. coincides with the classical notion of balancedness in case of classical cooperative games, and 2. is necessary and sufficient for core non-emptiness in a large class of SVGs.

The paper is organized as follows. Section 2 introduces the notion of set-valued games and related notions that are important for the analysis of such games. Section 3 presents the notion of a core. Section 4 discusses the relation of the core of an SVG to solution concepts in other approaches to multi-agenda disputes. Section 5 points to that a straightforward Bondareva-Shapley balancedness generalized condition cannot serve as a characterization of the core non-emptiness in the context of SVGs. Section 6 provides a proper generalization of the notion of balancedness, showing that it coincides with the classical one for standard cooperative games. Then, Section 7 shows that the proposed generalization characterizes the core non-emptiness for SVGs that satisfy some convexity conditions. Lastly, all proofs that do not appear in the main text appear in the Appendix.

³Logrolling is the trading of favors such as vote trading by legislative members.

⁴The characterization also holds in many other setups, such as those discussed in Gayer and Persitz [5] and Assa, Elliston and Lehrer [1].

2 Set-Valued Games

2.1 Cooperative Games

We begin by recalling a few classical definitions in cooperative games. Let $N = \{1, \dots, n\}$ be a finite set of players. A *coalition* is any subset of N , $S \subseteq N$. According to the classical definition, a (*cooperative*) *game* v associates a nonnegative real valued number, $v(S)$, to each coalition $S \subseteq N$. One of the leading solution concepts for games is the *core*: the core of a game v is defined as

$$\text{Core}(v) = \left\{ (x^i)_{i=1, \dots, N} : \sum_{i \in S} x^i \geq v(S), \forall S \subseteq N, \sum_{i \in N} x^i = v(N) \right\}. \quad (1)$$

The *cover* of a game v at the grand coalition N is defined as⁵

$$\hat{v}(N) = \max \left\{ \sum_{j=1}^{\ell} \alpha_j v(A_j); \sum_{j=1}^{\ell} \alpha_j \mathbf{1}_{A_j} = \mathbf{1}_N, \alpha_j \geq 0, A_j \subseteq N, j = 1, \dots, \ell \right\}.$$

Note that $v(N) \leq \hat{v}(N)$ for every game v . Bondareva and Shapley [2, 10] provided a necessary and sufficient condition for core non-emptiness: they showed that the core of a game v is not empty if and only if v is *balanced*, that is $v(N) = \hat{v}(N)$.

2.2 The definition

A *set-valued game* is a generalization of the notion of a game. For some $k \in \mathbb{N}$, a set-valued game associates to each coalition a *subset of vectors in \mathbb{R}_+^k* .

Definition 1. A set-valued game (SVG) over a collection N of players, is a function $\mathbf{v} : 2^N \rightarrow 2^{\mathbb{R}_+^k}$ defined over all subsets of N , which satisfies:

1. $\mathbf{v}(\emptyset) = \{0\}$;
2. Closedness. For every $S \subseteq N$, $\mathbf{v}(S)$ is a closed set;
3. Comprehensiveness. If $x \in \mathbf{v}(S)$ and $y \in \mathbb{R}_+^k$ is such that⁶ $y \leq x$, then $y \in \mathbf{v}(S)$.

⁵For any coalition $S \subseteq N$, the indicator $\mathbf{1}_S$ stands for the characteristic function of S , where $\mathbf{1}_S(i) = 1$ if $i \in S$ and 0 otherwise.

⁶ $y \leq x$ means that $y_j \leq x_j$ for every $j = 1, \dots, k$.

We will explicitly refer to a classical cooperative game as *a game*, as opposed to the more general set-valued game that will be referred to as *SVG*. In the remainder of this section we discuss possible properties of an SVG and its relation to games. The role of the definitions and examples we introduce is to motivate the notion of an SVG, and to familiarize the reader with this notion and related definitions pertinent to the analysis in subsequent sections.

Note, the definition of an SVG relies on transferable utilities within each game and non-transferable utilities across games. One can generalize the definition, and extend the analysis below appropriately, while relying solely on non-transferable utilities: an environment in which each game is a non-transferable utility one, and across games utility is non-transferable as well.⁷

2.3 Set-valued and classical games

The values that an SVG takes are subsets of \mathbb{R}_+^k . A (classical) game is defined like an SVG on subsets of N , but opportunities are modeled with a single dimension, and thus each coalition $S \subseteq N$ is associated with a single number reflecting its (maximum) payoff.

2.4 A multi-game based SVG

Consider a family of $k \in \mathbb{N}$ games v_1, \dots, v_k , and define a SVG \mathbf{v} by

$$\mathbf{v}(S) := \left\{ (\alpha_1 v_1(S), \dots, \alpha_k v_k(S)) \in \mathbb{R}_+^k; \alpha_j \geq 0 \forall j, \sum_{j=1}^k \alpha_j \leq 1 \right\}. \quad (2)$$

This definition is interpreted as follows. Consider $v_j(S)$ to be the amount of project j coalition S can complete in one unit of time. When the coalition invests $\alpha_j(S)$ in project j , where $\sum_{j=1}^k \alpha_j(S) \leq 1$, a vector $(\alpha_1 v_1(S), \dots, \alpha_k v_k(S))$ can be produced by coalition S in one time unit. An SVG of this kind is referred to as *multi-game based*.

The base games (v_j 's) capture the stand-alone productivity (or, value) for each project. The SVG defined through these games, \mathbf{v} , reflects the natural time constraint

⁷While not explicitly described as such, Voorneveld and van den Nouweland [8] study a model of multiple non-transferable utility set-valued games in that spirit and characterize different notions of the core.

and tradeoffs between investing in the different projects. Clearly, not every SVG is multi-game based, but the interpretation remains: $\mathbf{v}(S)$ captures the productivity of coalition S and reflects the time constraints faced by the coalition when contemplating how much time (or, more generally, resources) to invest in each project.

The following example illustrates the structure of a multi-game based SVG. It will play an important role in our subsequent analysis.

Example 1. Let $N = \{1, 2, 3\}$, $k = 2$, and define the following SVG: $\mathbf{v}(N) = \{(w_1, w_2); w_1 + w_2 \leq 3\}$, $\mathbf{v}(1, 2) = \{(w_1, w_2); 2w_1 + w_2 \leq 3\}$, $\mathbf{v}(2, 3) = \{(w_1, w_2); w_1 + 2w_2 \leq 3\}$, $\mathbf{v}(1, 3) = \{(w_1, w_2); w_1 + w_2 \leq 2\}$ and $\mathbf{v}(i) = \{(w_1, w_2); w_1 + w_2 \leq 1\}$, $i = 1, 2, 3$.

This SVG is multi-game based. Indeed, define two games v_1, v_2 in the following manner. $v_1(N) = 3$, $v_1(1, 2) = 1.5$, $v_1(2, 3) = 3$, $v_1(1, 3) = 2$, and $v_1(i) = 1$ for every player $i \in N$. Similarly, $v_2(N) = 3$, $v_2(1, 2) = 3$, $v_2(2, 3) = 1.5$, $v_2(1, 3) = 2$, and $v_2(i) = 1$ for every player $i \in N$. It is easy to verify that \mathbf{v} is a multi-game based on v_1, v_2 .

2.5 Additive SVGs

An SVG \mathbf{v} is *additive* if for every two disjoint coalitions, $S, T \subseteq N$,

$$\mathbf{v}(S) + \mathbf{v}(T) = \mathbf{v}(S \cup T), \quad (3)$$

where on the left hand side of the equality we mean the Minkowski sum. It is clear that \mathbf{v} is additive if and only if for every $S \subseteq N$, $\mathbf{v}(S) = \sum_{i \in S} \mathbf{v}(i)$. In terms of productivity, each worker's productivity is independent of the group she is working with: her contribution to a group is always the set of vectors she can produce alone.

2.6 The induced game

The following definition will be helpful in analyzing the core of an SVG in subsequent sections. Let $\lambda \neq 0$ be a non-negative vector in \mathbb{R}^k (i.e., $\lambda \in \mathbb{R}_+^k$) and \mathbf{v} be an SVG over N .

Definition 2. The λ -induced-game of \mathbf{v} is the game v_λ defined by⁸

$$v_\lambda(S) = \max_{y \in \mathbf{v}(S)} y \cdot \lambda \quad (4)$$

for every $S \subseteq N$.

Remark 1. Note that for every $S \subseteq N$, $\mathbf{v}(S) \subseteq \{y \in \mathbb{R}_+^k; y \cdot \lambda \leq v_\lambda(S)\}$.

3 The Core of an SVG

This section provides the definition of a central concept in the theories of cooperative games and capacities: the *core*.

Definition 3. The core of an SVG \mathbf{v} is defined as

$$\begin{aligned} \text{CORE}(\mathbf{v}) := & \left\{ (x^i)_{i \in N}; \text{ (a) for every } i \in N, x^i \in \mathbb{R}_+^k; \right. \\ & \text{(b) } \sum_{i \in N} x^i \in \mathbf{v}(N); \text{ and} \\ & \left. \text{(c) } \forall S \subseteq N, \text{ if } y \in \mathbf{v}(S) \text{ and } y \geq \sum_{i \in S} x^i, \text{ then } y = \sum_{i \in S} x^i \right\}. \end{aligned} \quad (5)$$

When $(x^i)_{i \in N}$ is in the core of \mathbf{v} , member i of N would get the share x^i , which is a vector in \mathbb{R}_+^k . That is, a core member is a “payoff” to each player for each of the k agendas. The total share of all the members of N is a feasible vector, namely in $\mathbf{v}(N)$. Finally, it maintains stability in the sense that there is no coalition S that could find a better $y \in \mathbf{v}(S)$. That is, there is no $y \in \mathbf{v}(S)$ that dominates (Pareto) the total share of the S -members, $\sum_{i \in S} x^i$. Note that when $k = 1$ the definition of the core of an SVG as in Eq. (5) coincides with the classical definition of the core (as in Eq. (1)).

Example 1 (cont.) Recall the SVG in Example 1. Consider $x^1 = (0, 1)$, $x^2 = (1, 0)$, and $x^3 = (0, 1)$. The sum, $x^1 + x^2 + x^3 = (1, 2)$ is on the Pareto frontier of $\mathbf{v}(N)$. Furthermore, $x^1 + x^2 = (1, 1)$, $x^1 + x^3 = (0, 2)$, $x^2 + x^3 = (1, 1)$, x^1, x^2 , and x^3 , are on the Pareto frontiers of $\mathbf{v}(1, 2)$, $\mathbf{v}(1, 3)$, $\mathbf{v}(2, 3)$, $\mathbf{v}(1)$, $\mathbf{v}(2)$ and $\mathbf{v}(3)$, respectively. Thus, (x^1, x^2, x^3) is in the core of \mathbf{v} .

⁸ $y \cdot \lambda$ denotes the inner product of y and λ .

4 Multi-Game Based SVGs and Logrolling

SVGs that are multi-game based were introduced in Section 2.4. For an SVG in this family it is natural to study the relation between the cores of the base games and the core of the SVG itself.

Bloch and de Clippel [3] and Gayer and Persitz [5] also study multi-agenda disputes, but, as explained in the Introduction, take a different approach. Both papers are conceptually different than the current study, and consider the sum of the base games and study notions of the core in this set up. The core of the sum and the core of an SVG live in different spaces, however the relation between these approaches can be studied. We point to the relation and distinction between the approaches.

4.1 Base games, cores, and their sums

We formally introduce the ideas in Bloch and de Clippel [3] and Gayer and Persitz [5]. Let v_1, \dots, v_k be a family of (base) games, and $v = v_1 + \dots + v_k$ be the sum of these games. Bloch and de Clippel [3] study the relation between the cores of the base games and that of v . Gayer and Persitz [5] then introduce the notion of a multi-core of v_1, \dots, v_k . Conceptually, a multi-core element is a payoff to each individual for participating in the game v , where: 1. for each individual there may be a (different) subjective assessment of how these individual payoffs are generated from the payoffs for the base games; and 2. given these subjective assessments, no individual thinks that she can jointly deviate with some of the other agents and increase their payoffs. That is, a payoff $(w^i)_{i \in N}$ is in the multi-core of v if for each agent $p \in N$ there is a feasible payoff vector $y(p) \in \mathbb{R}^{n \times k}$, where $y(p)_j^i$ is agent p 's subjective assessment of i 's payoff in game j , such that:

- $\sum_{j=1}^k y(p)_j^i = w_i$ for every $p, i \in N$;
- $\sum_{i=1}^N y(p)_j^i = v_j(N)$ for every $p \in N$ and $j \in \{1, \dots, k\}$;
- For every coalition $S \subseteq N$ and $p \in S$, $\sum_{i \in S} y(p)_j^i \geq v_j(S)$ for every $j \in \{1, \dots, k\}$.

A result of interest is Theorem 3 in Gayer and Persitz [5] stating that given any collection of base games, the sum of their cores is a subset of the multi-core, which is a subset of the core of the sum of the base games.

To begin understanding the relation of all of these concepts to the core of an SVG, consider the following example. It turns out that it is possible that the cores of the v_j 's, the sum of the cores, the core of the sum, and the multi-core of the v_j 's, are all empty, while the core of the multi-game SVG based on v_1, \dots, v_k is not.

Indeed, consider the SVG presented in Example 1. We saw in Section 3 that the core of the SVG is not empty. Now, both v_1 and v_2 are not balanced. For example,

$$\hat{v}_1(N) \geq v_1(2, 3) + v_1(1) = 4 > 3 = v_1(1, 2, 3).$$

A similar calculation holds for the game v_2 . Thus, the cores of v_1 and v_2 are empty, and we obtain that the sum of the cores is empty. In addition, the core of the sum $v_1 + v_2$ is empty as well. Letting $v = v_1 + v_2$, we have that

$$\hat{v}(N) \geq v(2, 3) + v(1) = 6.5 > 6 = v(1, 2, 3),$$

implying v is not balanced, and thus has an empty core. By Theorem 3 in Gayer and Persitz [5], the multi-core of v_1, v_2 is empty as well.

Now consider the agendas, v_1, \dots, v_k . We show that if the core of each v_j is not empty, then every ‘‘convex combination’’ of elements (across the k cores), is in the core of the multi-game SVG based on v_1, \dots, v_k .

Proposition 1. *Let \mathbf{v} be a multi-game SVG over N based on v_1, \dots, v_k , and assume x_j is in the core of v_j for every j . Let $z^i = (\alpha_j x_j^i)_{j=1}^k$ (where $\alpha_j \geq 0$ for every j and $\sum_{j=1}^k \alpha_j = 1$). Then $(z^i)_{i \in N}$ is in the core of \mathbf{v} .*

Proof. For each j , let x_j be a core member of v_j . For every player $i \in N$, let $z^i = (\alpha_j x_j^i)_{j=1}^k$. We show that $(z^i)_{i \in N}$ satisfies the three conditions in Eq. (5). Indeed, (a) is clearly satisfied. As for (b), since x_j is in the core of v_j , we have that $\sum_{i \in N} z^i = \alpha_j v_j(N)$, and thus $\sum_{i \in N} z^i \in \left\{ (\beta_1 v_1(N), \dots, \beta_k v_k(N)) \in \mathbb{R}_+^k; \beta_j \geq 0 \forall j, \sum_{j=1}^k \beta_j \leq 1 \right\} = \mathbf{v}(N)$. And lastly, to see that condition (c) holds, fix a coalition $S \subseteq N$ and consider a vector $(\beta_1 v_1(S), \dots, \beta_k v_k(S)) \in \mathbf{v}(S)$ such that $(\beta_1 v_1(S), \dots, \beta_k v_k(S)) \geq \sum_{i \in S} z^i = (\alpha_1 \sum_{i \in S} x_1^i, \dots, \alpha_k \sum_{i \in S} x_k^i)$. Since x_j is in the core of the respective v_j , we have that $\sum_{i \in S} x_j^i \geq v_j(S)$. Thus, (c) is satisfied and the proof is completed. \square

A similar result can be stated regarding the multi-core of the sum of the base games. The following proposition states that every subjective assessment of a member in the

multi-core is also a member in the core of the SVG. The proof is identical to that of 1 and is therefore omitted.

Proposition 2. *Let \mathbf{v} be a multi-game based SVG over N based on v_1, \dots, v_k , and assume $(y(p))_{p \in N}$ is a subjective justification vector (for some member of the multi-core of (v_1, \dots, v_k)). For every player $p \in N$ and game j , let $x(p)_j = (y(p)_j)_{i \in N}$ and $z(p)^i = (\alpha_j x(p)_j)_{j=1}^k$ (where $\alpha_j \geq 0$ for every j and $\sum_{j=1}^k \alpha_j = 1$). Then $(z(p)^i)_{i \in N}$ is in the core of \mathbf{v} , for every $p \in N$.*

The converse statements of Propositions 1 and 2 do not hold. We saw in Example 1 that it is possible that the base games have an empty core, while the core of the multi-game based SVG is not empty. The following example shows that even if the base games have non-empty cores, they do not necessarily span the core of the multi-game based SVG.

Example 2. *Consider a variation of the game described in Example 1. Let $N = \{1, 2, 3\}$, and define $u_1(N) = 3, u_1(1, 2) = u_1(1, 3) = 1.5, u_1(2, 3) = 3$, and $u_1(i) = 0$ for all $i \in N$. Similarly, let $u_2(N) = 3, u_2(1, 3) = u_2(2, 3) = 1.5, u_2(1, 2) = 3$, and $u_2(i) = 0$ for all $i \in N$.*

The core of u_1 consists only of the allocation $(0, 1.5, 1.5)$, and the core of u_2 consists only of the allocation $(1.5, 1.5, 0)$. It is simple to verify that $x^1 = (0, 1), x^2 = (1, 0)$, and $x^3 = (0, 1)$ is in the core of the multi-game SVG \mathbf{u} based on u_1 and u_2 . However, (x^1, x^2, x^3) is not spanned (in the sense of Proposition 1) by the cores of u_1 and u_2 .

In addition, defining $u = u_1 + u_2$, we obtain that $u(N) = 6, u(1, 2) = u(2, 3) = 4.5, u(1, 3) = 3$, and $u(i) = 0$ for all $i \in N$. We see that the only core member of u is $(1.5, 3, 1.5)$, which is also the sum of the cores of u_1 and u_2 . Again, by Theorem 3 in Gayer and Persitz [5], we obtain that the multi-core comprises only of $(1.5, 3, 1.5)$.

4.2 Logrolling

For the remaining of this section we show that for a class of multi-game based SVGs, the core of the SVG is not a convex span of the cores of the base games. This class consists of base games with a unique core element. An example for such games is the family of additive games: a game v is additive if $v(S) = \sum_{i \in S} v(i)$ for every coalition $S \subseteq N$. In this case the unique core element, x , of v is given by $x^i = v(i)$ for every

player $i \in N$. Note also that the sum of additive games is additive, implying that the core of the sum is the sum of the cores of the base games. In particular, for such games the multi-core coincides with the sum of the cores, and consists of the unique element only.

Example 3. *Consider two additive two-player base games, u, v , where $v(i) = u(i) = 1$ for every player i . The core element of each of these games assigns payoff 1 for each of the players.*

From the propositions above, the core of the SVG based on these two games contains the convex span of the cores of the base games. In particular, (x^1, x^2) , where $x^1 = x^2 = (0.5, 0.5)$, is in the core of the SVG. However, notice that (\hat{x}^1, \hat{x}^2) , where $\hat{x}^1 = (0, 1), \hat{x}^2 = (1, 0)$ is also in the core of the SVG, and is not a convex span of the cores of the two base games.

The example shows that by reallocating payoffs between players the SVG provides a space for advantageous cooperation beyond what is given by the separate base games, or their sum. The next example shows that not every reallocation is possible if one wishes to maintain stability.

Example 4. *Consider two additive two-player base games, u, v , where $v(1) = u(3) = 1, v(2) = u(2) = 2$, and $v(3) = u(1) = 3$. Say that players tried to reallocate their payoffs as follows: player 3 provides 1 payoff unit in game v for player 1 in favor of 1 unit payoff in game u , and considering the 0.5 – 0.5 mixture between the two games. The resulting payoffs are $x^1 = (0, 2)$ (where the left coordinate corresponds to the payoff in agenda v), $x^2 = (1, 1)$, and $x^3 = (2, 0)$. We show that this allocation is not in the core of the SVG based on v and u .*

In this case, the coalition comprising of players 1 and 2 has an incentive to deviate and play on their own. Indeed, if they consider a $p, 1 - p$ mixture of the base games, their joint payoff as a function of p will satisfy $3p \geq 1$ and $5(1 - p) \geq 3$ for any $0.4 \geq p \geq 0.333\dots$. That is, for any p strictly in between these values, players 1 and 2 have an incentive to deviate as a coalition. We obtain that $(x^i)_i$ as suggested above is not a core member.

It is possible, however, to find another payoff reallocation between players 1 and 3 that would give rise to a new core element that is not in the convex span of the cores of the base games. Indeed, $x^1 = (0.75, 1.25)$, $x^2 = (1, 1)$, and $x^3 = (1.25, 0.75)$ is such

a reallocation.⁹ The coalition comprising of players 1 and 2 would deviate if $3p \geq 1.75$ and $5(1 - p) \geq 2.25$, implying that $0.55 \geq p \geq 0.58333\dots$ —such a p does not exist. Similarly, The coalition comprising of players 3 and 2 would deviate if $5p \geq 2.25$ and $3(1 - p) \geq 1.75$, which is satisfied only for $0.41666\dots \geq p \geq 0.45$. Again, there is no such p .

The next Theorem states that we can always find a core member of an SVG based on additive games (with strictly positive individual payoffs) that cannot be obtained by the convex span of the base games. The proof relies on a simple economic intuition behind the notion of comparative advantage adapted to the current setup (that is, where there are more than two agents). It shows that we can always reallocate some “value” between two of the players—the one with the lowest comparative advantage in one agenda and the one with the highest comparative advantage in that same agenda—and construct a new core member that is not in the span of the cores of the base games.

Theorem 1. *Let \mathbf{v} be an SVG based on the additive games v_1, \dots, v_k , not all are equivalent¹⁰ and $v_j(i) > 0$ for every $j = 1, \dots, k$ and $i \in N$. Then, there exists a core element of \mathbf{v} , x^1, \dots, x^n , such that there are no $\alpha_j \geq 0 \forall j, \sum_{j=1}^k \alpha_j = 1$ where $x^i = (\alpha_1 v_1(i), \alpha_2 v_2(i), \dots, \alpha_n v_n(i)), i \in N$.*

Proof. We first assume that $k = 2$. Let $r = (v_1(N), -v_2(N)) \in \mathbb{R}^2$, and for $i \in N$ let $\beta_i = v_2(i) - \frac{v_2(N)}{v_1(N)} v_1(i)$ and $y^i = (\frac{1}{2} v_1(i), \frac{1}{2} v_2(i))$. For every $S \subseteq N$ let $y(S) = \sum_{i \in S} y^i$. Note that $y(S)$ is on the Pareto efficient frontier of $\mathbf{v}(S)$ and is strictly positive in both coordinates. Thus, $y(S) + \gamma \sum_{i \in S} \beta_i r$ is also strictly positive in both coordinates for sufficiently small $\gamma > 0$.

For $i \in N$ define $x^i = y^i + \gamma \beta_i r$, and for $S \subseteq N$ let $x(S) = \sum_{i \in S} x^i$. By assumption not all of v_1, \dots, v_k , are equivalent. This implies that $\beta_i \neq 0$ and thus $x^i \neq y^i$ for at least one i . We show that x^1, \dots, x^n is a core element of \mathbf{v} .

Note that $\sum_{i \in N} \beta_i = v_2(N) - \frac{v_2(N)}{v_1(N)} v_1(N) = 0$. Thus, $x(N) = y(N)$ and therefore $x(N) \in \mathbf{v}(N)$. Now fix $S \subseteq N$. We show that $x(S)$ is not dominated by any point in

⁹These payoffs correspond to reallocation of half a payoff unit between players 1 and 3 across the two base games (player 3 provides 0.5 in game v for player 1 in favor of 0.5 in game u), and considering the 0.5 – 0.5 mixture between the two games.

¹⁰Two additive games v and u are equivalent if $\frac{v(i)}{u(i)}$ is constant across i .

$\mathbf{v}(S)$. To this end we show that

$$\frac{v_2(S)}{v_1(S)} \geq \frac{v_2(N)}{v_1(N)} \text{ if and only if } \sum_{i \in S} \beta_i \geq 0.$$

This equivalence means that the frontier of $\mathbf{v}(S)$ is steeper than that of $\mathbf{v}(N)$ if and only if $x(S) - y(S) = \sum_{i \in S} \gamma \beta_i r$. That is, $x(S) - y(S)$ is a vector r multiplied by a nonnegative coefficient $(\gamma(\sum_{i \in S} \beta_i)r)$.¹¹ Since $y(S)$ is on the frontier of $\mathbf{v}(S)$, this implies that $x(S)$ lies beyond this frontier.

Indeed,

$$\sum_{i \in S} \beta_i = \sum_{i \in S} v_2(i) - \frac{v_2(N)}{v_1(N)} v_1(i) = v_2(S) - \frac{v_2(N)}{v_1(N)} v_1(S) \geq 0 \text{ if and only if } \frac{v_2(S)}{v_1(S)} \geq \frac{v_2(N)}{v_1(N)},$$

as desired.

As for a general k , one can restrict attention to two non-equivalent games, say w.l.o.g. v_1 and v_2 , and consider $y^i = (\frac{1}{2}v_1(i), \frac{1}{2}v_2(i), 0, \dots, 0), i \in N$ as the original core member of \mathbf{v} . Then define $r = (v_1(N), -v_2(N), 0, \dots, 0) \in \mathbb{R}^n$ to be the vector by which y^i is shifted. The proof proceeds along the same line as in the case of $k = 2$. \square

In case there are only two players Ricardo defined the notion of comparative advantage. In our context there are many players. The advantage of player i is measured in comparison with the grand coalition. We say that player i has a comparative advantage in project 2 if $\frac{v_2(i)}{v_1(i)} \geq \frac{v_2(N)}{v_1(N)}$.

The proof of Theorem 1 is based on the idea that when player i has a comparative advantage in project 2, he can give up some value related to project 2 and get in return some value related to project 1. The rate at which this logrolling/trade-off is carried out is determined by the grand coalition, and formally by the vector r . Player i would receive (in addition to y^i) the vector $\gamma \beta_i r$.

The trade-off of player i depends not only on the project at which player i has a comparative advantage, but also on two additional factors: the magnitude of the comparative advantage (i.e., $\frac{v_2(i)}{v_1(i)} - \frac{v_2(N)}{v_1(N)}$) and the “power” of player i (measured as $v_1(i)$). This is the reason why player i ’s trade-off across all projects, represented by $x^i - y^i$, is proportional to $v_1(i) \left(\frac{v_2(i)}{v_1(i)} - \frac{v_2(N)}{v_1(N)} \right) r$.

¹¹In other words, it is vector in the “south-western” orthant whose slope (like that of r) is smaller than that of the frontier of $\mathbf{v}(S)$.

Remark 2. The vector $r = (r_1, r_2)$ has been chosen so that $|r_2|/r_1 = |v_2(N)|/v_1(N)$. Notice, the same proof would hold if r is such that

$$\frac{|r_2|}{r_1} \leq \min \left\{ \frac{v_2(S)}{v_1(S)}; \frac{v_2(N)}{v_1(N)} < \frac{v_2(S)}{v_1(S)}, S \neq \emptyset \right\},$$

and

$$\frac{|r_2|}{r_1} \geq \max \left\{ \frac{v_2(S)}{v_1(S)}; \frac{v_2(N)}{v_1(N)} > \frac{v_2(S)}{v_1(S)}, S \neq \emptyset \right\}.$$

It implies that the set of possible logrolling opportunities has a full dimension (i.e., two).

5 Non Bondareva-Shapley and SVGs

5.1 The cores of \mathbf{v} and v_λ

In this section we point to the relation between the core of \mathbf{v} and that of the induced game v_λ . This relation will be significant in terms of finding conditions for the non-emptiness of the core of \mathbf{v} . Prior to this we need to introduce the notion of a cover of an SVG.

Definition 4. Let \mathbf{v} be an SVG. The cover of \mathbf{v} at the coalition N is

$$\hat{\mathbf{v}}(N) = \left\{ \sum_{j=1}^{\ell} \alpha_j y_j; \sum_{j=1}^{\ell} \alpha_j \mathbf{1}_{A_j} = \mathbf{1}_N, y_j \in \mathbf{v}(A_j), \alpha_j \geq 0, A_j \subseteq N, j = 1, \dots, \ell \right\}. \quad (6)$$

Notice that $\hat{\mathbf{v}}(N)$ is a convex set.

Proposition 3. Fix $\lambda \in \mathbb{R}_+^k$, $\lambda \neq 0$. The core of v_λ is not empty if and only if there exists $x \in \mathbf{v}(N)$ such that

$$x \in \operatorname{argmax} \left\{ y \cdot \lambda; y \in \hat{\mathbf{v}}(N) \right\}. \quad (7)$$

Proof. Assume first that there is $x \in \mathbf{v}(N)$ such that $x \in \operatorname{argmax}_{y \in \hat{\mathbf{v}}(N)} y \cdot \lambda$. The sets $\hat{\mathbf{v}}(N)$ and $C := \{y; y \geq x\}$ (the latter being the positive orthant relative to x) are convex. Due to Eq. (7), $\lambda \in \mathbb{R}_+^k$ is a separating vector:

$$y \cdot \lambda \leq z \cdot \lambda \text{ for every } y \in \hat{\mathbf{v}}(N) \text{ and } z \in C. \quad (8)$$

We show that the core of v_λ is non-empty. Otherwise, by the Bondareva-Shapley Theorem, $\hat{v}_\lambda(N) > v_\lambda(N)$. It implies that there is a decomposition of $\mathbf{1}_N$, $\mathbf{1}_N = \sum_{j=1}^{\ell} \alpha_j \mathbf{1}_{A_j}$, with $\alpha_j \geq 0, j = 1, \dots, \ell$, such that $\sum_{j=1}^{\ell} \alpha_j v_\lambda(A_j) > v_\lambda(N)$. Thus, for every $j = 1, \dots, \ell$ there is $x_j \in \mathbf{v}(A_j)$ such that

$$\sum_{j=1}^{\ell} \alpha_j x_j \cdot \lambda > \max_{w \in \mathbf{v}(N)} w \cdot \lambda \geq x \cdot \lambda. \quad (9)$$

Since $\sum_{j=1}^{\ell} \alpha_j x_j \in \hat{v}(N)$, Eq. (9) contradicts Eq. (8). We conclude that the core of v_λ is not empty.

Now assume that no $x \in \mathbf{v}(N)$ satisfies Eq. (7). Since $\mathbf{v}(N) \subseteq \hat{\mathbf{v}}(N)$, it implies that $\max_{y \in \hat{v}(N)} y \cdot \lambda > \max_{y \in \mathbf{v}(N)} y \cdot \lambda$. Therefore, $\hat{v}_\lambda(N) > v_\lambda(N)$, which by the Bondareva-Shapley Theorem implies that the core of v_λ is empty. \square

The following proposition will be useful in the subsequent section. This result also appears in Fernandez et al. [4]. The proof is provided for completeness.

Proposition 4. *Suppose that there exists $\lambda \in \mathbb{R}_+^k$ such that the core of the induced game v_λ is not empty. Then, $\text{CORE}(\mathbf{v}) \neq \emptyset$.*

Proof. Let $(w_i)_{i \in N}$ be in the core of v_λ , in particular, $\sum_{i \in N} w_i = v_\lambda(N)$. Denote $x := \operatorname{argmax}_{z \in \mathbf{v}(N)} z \cdot \lambda$. By definition, $v_\lambda(N) = x \cdot \lambda$. Also denote, $w := v_\lambda(N)$ and $x^i := \frac{w_i}{w} x$ (recall, $x \in \mathbb{R}_+^k$, $\frac{w_i}{w} \in \mathbb{R}$ and $\sum_i \frac{w_i}{w} = 1$).

We obtain: (a) $x^i \in \mathbb{R}_+^k$; (b) $\sum_{i \in N} x^i = \sum_{i \in N} \frac{w_i}{w} x = x \in \mathbf{v}(N)$; and (c) for every $S \subseteq N$, $\sum_{i \in S} w_i \geq v_\lambda(S)$, implying

$$\left(\sum_{i \in S} x^i \right) \cdot \lambda = \left(\sum_{i \in S} \frac{w_i}{w} x \right) \cdot \lambda = \left(\sum_{i \in S} \frac{w_i}{w} \right) v_\lambda(N) = \sum_{i \in S} w_i \geq v_\lambda(S) = \max_{y \in \mathbf{v}(S)} y \cdot \lambda.$$

This implies that if $y \in \mathbf{v}(S)$ and $y \geq \sum_{i \in S} x^i$, then $y = \sum_{i \in S} x^i$. \square

5.2 A sufficient condition for the non-emptiness of the core

The next theorem provides a sufficient condition for the non-emptiness of the core. The following notion is useful.

Definition 5. *Let $D \subseteq \mathbb{R}^k$ be compact. We say that $x \in D$ is D -efficient if $y \in D$ and $y \geq x$ imply $y = x$.*

Theorem 2. *Let \mathbf{v} be an SVG. If there exists $x \in \mathbf{v}(N)$ which is $\hat{\mathbf{v}}(N)$ -efficient, then the core of \mathbf{v} is not empty.*

Before presenting the proof, note that the condition in the theorem is a generalization of the classical notion of balancedness of a game. Indeed, a game v is said to be *balanced* if $\hat{v}(N) = v(N)$.

Proof. Let $x \in \mathbf{v}(N)$ be $\hat{\mathbf{v}}(N)$ -efficient. Define, $C := \{z; z \geq x\}$. The sets $\hat{\mathbf{v}}(N)$ and C are two convex sets. Due to the fact that x is $\hat{\mathbf{v}}(N)$ -efficient, the intersection of these sets contains only x . The set C is of full dimension and therefore there is a non-zero vector $\lambda \in \mathbb{R}_+^k$ separating $\hat{\mathbf{v}}(N)$ and C . That is, $y \cdot \lambda \leq z \cdot \lambda$ for every $y \in \hat{\mathbf{v}}(N)$ and $z \in C$. We have that x satisfies Eq. (7) with respect to λ . Due to Proposition 3, the core of v_λ is not empty which implies, by Proposition 4, that the core of \mathbf{v} is not empty. \square

The condition of Theorem 2 has the same spirit as that of the Bondareva-Shapley Theorem. Indeed, when $k = 1$, this condition is equivalent to that of the Bondareva-Shapley Theorem and is therefore not only sufficient, but also necessary. The following example shows that the core of \mathbf{v} can be non-empty, while the core of v_λ is empty for every $\lambda \in \mathbb{R}_+^k$. It implies that the condition of Theorem 2 is not necessary. This example requires only two agendas.

5.3 The sufficient condition is not necessary

In Section 3 we saw that the core of the SVG in Example 1 is not empty. On the other hand, we show now that for every $\lambda \in \mathbb{R}_+^k$, the core of v_λ is empty. For this end, it is sufficient, by Proposition 3, to show that the efficient frontier of $\mathbf{v}(N)$ is strictly dominated by a point in $\hat{\mathbf{v}}(N)$. That is, for every $x \in \mathbf{v}(N)$ there is $y \in \hat{\mathbf{v}}(N)$ such that $y_i > x_i, i = 1, 2$. However, since the efficient frontier of $\mathbf{v}(N)$ is a straight line, since $\hat{\mathbf{v}}(N)$ is convex, it is sufficient to show that this statement is true for the two $\mathbf{v}(N)$ -efficient points that lie on the axes. The $\mathbf{v}(N)$ -efficient point on the vertical axes is $(0, 3)$. However, consider the decomposition $\mathbf{1}_N = \frac{1}{2}(\mathbf{1}_{\{1,2\}} + \mathbf{1}_{\{2,3\}} + \mathbf{1}_{\{1,3\}})$ and the most upper points in $\mathbf{v}(1, 2), \mathbf{v}(2, 3), \mathbf{v}(1, 3)$ that lie on the vertical axes. These points are, respectively, $(0, 3), (0, 1.5)$ and $(0, 2)$. Thus, the point $\frac{1}{2}((0, 3) + (0, 1.5) + (0, 2)) = (0, 3.25) \in \hat{\mathbf{v}}(N)$. This point is strictly above $(0, 3)$. A similar argument applies to

the horizontal axes. We therefore conclude that the two $\mathbf{v}(N)$ -efficient points that lie on the vertical axes are strictly dominated by points in $\hat{\mathbf{v}}(N)$ and the argument is complete.

To summarize, this example shows that the core of \mathbf{v} is not empty despite the facts that the core of any v_λ is empty and that any $\mathbf{v}(N)$ -efficient point is strictly dominated by a point in $\hat{\mathbf{v}}(N)$.

6 Extended Bondareva-Shapley

In this section we provide a necessary and sufficient condition for the non-emptiness of the core of an SVG in the case where the set $\mathbf{v}(N)$ is convex and, moreover, the set of points in \mathbb{R}_+^k that dominate $\mathbf{v}(T)$ is convex for every $T \subsetneq N$.

6.1 The one-dimensional case

We start out with the one-dimensional case which is tightly connected with classical cooperative games and the Bondareva-Shapley characterization of core non-emptiness. Section 6.3 below, and Theorem 3 therein, generalizes these results to multi-dimensions and SVGs.

Let $x_1, \dots, x_l \in \mathbb{R}^n$ and $A_1, \dots, A_l \subseteq \mathbb{R}$ be non-empty, convex and closed. Assume that each A_i is either bounded or polyhedral. Define X to be the $l \times n$ dimensional matrix whose i -th row is x_i . In addition, define the following two sets: $A = A_1 \times \dots \times A_l$ and $B = \{Xc; c \in \mathbb{R}^n\}$. Both sets are closed and convex.

Proposition 5. *There exists $c \in \mathbb{R}^n$ such that $x_i \cdot c \in A_i$ for every $i = 1, \dots, l$ if and only if for every vector $s = (s_1, \dots, s_l) \in \mathbb{R}^l$,*

$$\sum_i s_i x_i = 0 \text{ implies } 0 \in \sum_i s_i A_i. \quad (10)$$

Note that the condition in Eq. (10) is equivalent to the following condition:

$$sX = 0 \text{ implies that there exists } a \in A \text{ such that } s \cdot a = 0. \quad (11)$$

6.2 Implication: the core of a game

The problem related to the non-emptiness of the core of a game v with n players is defined by the following parameters. The vectors $x_S = \mathbf{1}_S, S \subseteq N$ are in \mathbb{R}^n , and the sets $A_S = [v(S), \infty)$ when $S \subsetneq N$ and $A_N = \{v(N)\}$. The existence of $c \in \mathbb{R}^n$ such that $x_S \cdot c \in A_S$ for every $S \subseteq N$ is equivalent to the existence of $c \in \mathbb{R}^n$ such that

$$\mathbf{1}_S \cdot c \geq v(S) \text{ for every } S \subsetneq N, \text{ and } \mathbf{1}_N \cdot c = v(N). \quad (12)$$

Such a vector c is a core member of v . Proposition 5 therefore states that the core of v is non-empty if and only if the condition of Eq. (10) is satisfied.

To see that the condition Proposition 5 is equivalent to the Bondareva-Shapley consider the equation $sX = 0$, where $s = (s_T)_{T \subseteq N}$. In case there are two proper coalitions $R, T \subsetneq N$ such that $s_R s_T < 0$, then $s_R A_R + s_T A_T = \mathbb{R}$. This implies in particular that $0 \in \sum_T s_T A_T$. In other words, when the product $s_R s_T$ is negative, the proposition's condition is fulfilled automatically.

We can therefore assume that $s_T \geq 0$ for every proper coalition T . It must be then that $s_N < 0$. Therefore $sX = 0$ implies $\sum_{T \neq N} s_T x_T = |s_N| \mathbf{1}_N$. If $s_N \neq 0$, it means that we have a *balanced collection*, as referred to in the theory of cooperative games, with $s_T/|s_N|$ being the balancing coefficients. The Bondareva-Shapely condition dictates that a balanced collection must satisfy $\sum_{T \neq N} (s_T/|s_N|)v(T) \leq v(N)$. The latter inequality is equivalent to $0 \in \sum_{T \neq N} s_T A_T + s_N A_N$, which is what stated in Eq. (11).

6.3 The multi-dimensional case

Let $x_1, \dots, x_l \in \mathbb{R}^n$ and $A_1, \dots, A_l \subseteq \mathbb{R}^k$ be non-empty, convex and closed. As above X is defined as the $l \times n$ dimensional matrix whose i -th row is x_i . Let A is the set of all $l \times k$ matrices whose i -th row is a vector in A_i .

Definition 6. We say that $x_1, \dots, x_l; A_1, \dots, A_l$ are balanced if for every $k \times l$ -dimensional matrix S ,

$$SX = \mathbf{0} \text{ implies that there exists } a \in A \text{ such that } Sa = \mathbf{0}. \quad (13)$$

The term ‘balanced’ is inspired by the similarity between Eqs. (11) and (13) and, given this, the following is an extended Bondareva-Shapley theorem.

Theorem 3. *Assume that each A_i is a union of a bounded convex and closed set and a polyhedron.¹² There exists an $n \times k$ dimensional matrix C such that*

$$x_i C \in A_i \text{ for every } i = 1, \dots, l$$

if and only if $x_1, \dots, x_l; A_1, \dots, A_l$ are balanced.

Proof. The ‘only if’ direction is simple and is therefore omitted. For the inverse direction define $B = \{XC; C \text{ is an } n \times k \text{ matrix}\}$. Note that both sets A and B are closed and convex.

In case $A \cap B \neq \emptyset$, the proof is complete. Otherwise, A and B are disjoint.

Lemma 1. *There exists a separating $k \times l$ -dimensional matrix $S \neq 0$, such that¹³*

$$\text{tr}(Sb) > \text{tr}(Sa) \text{ for every } b \in B \text{ and } a \in A. \quad (14)$$

The proof is postponed to the Appendix.

Lemma 1 states that there exists a matrix S such that $\text{tr}(SXC) > \text{tr}(Sa)$ for every $n \times k$ -dimensional matrix C and $a \in A$. We claim that $SX = \mathbf{0}$ ($\mathbf{0}$ being the all-0 matrix). Otherwise, one can find a matrix C such that $\text{tr}(SXC) < \text{tr}(Sa)$ for some $a \in A$. Indeed, fix one $a \in A$. If $SX \neq 0$, then there is an entry, say (i, j) , of the matrix SX which is not 0, namely $(SX)_{ij} \neq 0$. Now consider the matrix C whose (j, i) entry is equal to $\frac{-2|\text{tr}(Sa)|}{(SX)_{ij}}$, and all other entries are 0. We obtain that $\text{tr}(SXC) = -2|\text{tr}(Sa)|$, which is strictly smaller than $\text{tr}(Sa)$.

As $SX = \mathbf{0}$, we obtain that for every $n \times k$ dimensional matrix C , $SXC = \mathbf{0}C = \mathbf{0}$ implying

$$\text{tr}(SXC) = 0. \quad (15)$$

On the other hand, balanceness condition (i.e., Eq. (13)) guarantees that due to the fact that $SX = \mathbf{0}$, there is $a \in A$ such that $Sa = \mathbf{0}$ and therefore

$$\text{tr}(Sa) = 0. \quad (16)$$

¹²A polyhedron is a set defined by a finite number of weak linear inequalities. Such a set might be unbounded.

¹³ tr is the trace operator: $\text{tr}(E)$ is the sum of the diagonal entries in the matrix E .

Eqs. (15) and (16) violate Eq. (14). We conclude that $A \cap B \neq \emptyset$ which completes the proof. \square

7 Non-emptiness of the Core in the Convex Case

In this section we discuss the non-emptiness of the core of an SVG under some convexity conditions, and its characterization by the extended Bondareva-Shapley condition presented in Theorem 3.

7.1 The problem behind core non-emptiness

In order to explain the difficulty behind the non-emptiness of the core, define (compare with Eq. (5))

$$C(\mathbf{v}, T) = \left\{ (x^i)_{i \in N}; \begin{array}{l} \text{(a) for every } i \in N, x^i \in \mathbb{R}_+^k; \\ \text{(b) } \sum_{i \in N} x^i \in \mathbf{v}(N); \text{ and} \\ \text{(c) if } y \in \mathbf{v}(T) \text{ and } y \geq \sum_{i \in T} x^i, \text{ then } y = \sum_{i \in S} x^i. \end{array} \right\} \quad (17)$$

Note that

$$CORE(\mathbf{v}) = \bigcap_{T \subseteq N} C(\mathbf{v}, T). \quad (18)$$

Thus, in order for the core of \mathbf{v} to be non-void, the intersection of all $C(\mathbf{v}, T), T \subseteq N$, must be non-empty. Condition (c) in Eq. (17) can be also expressed as

$$\min_{y \in \mathbf{v}(T)} \max_{j=1, \dots, k} \sum_{i \in T} x_j^i - y_j \geq 0,$$

which is equivalent to¹⁴

$$\forall S \subseteq N, \sum_{i \in T} x^i \in \text{cl}(\mathbb{R}_+^k \setminus \mathbf{v}(T)).$$

¹⁴ $\text{cl}()$ is the closure operator.

While the first two conditions of Eq. (17) are convex (when $\mathbf{v}(N)$ is convex), typically $\mathbb{R}_+^k \setminus \mathbf{v}(T)$ is not convex, hence the last condition is not. Condition (c) makes it clear why the set defined in Eq. (17) is not convex. It therefore seems that finding a necessary and sufficient condition for the non-emptiness of the intersection of sets that are not convex (see Eq. 18), goes beyond straightforward duality.

In what follows we address this issue and provide a characterization of when the core of a SVG is not empty in cases where $\mathbf{v}(N)$ is convex, and $\mathbb{R}_+^k \setminus \mathbf{v}(T), T \subsetneq N$ are convex, or a finite union of convex sets. The question remains open in more general cases.

7.2 Characterization of the core of a SVG: the convex case

Theorem 3 implies the following.

Theorem 4. *Let \mathbf{v} be a set-valued game defined on the set N . Define $x_T = \mathbf{1}_T$ for every $T \subseteq N$. Also, for every $T \subsetneq N$ define $A_T = \text{cl}(\mathbb{R}_+^k \setminus \mathbf{v}(T))$ and $A_N = \mathbf{v}(N)$.¹⁵ Suppose that A_T is convex for every $T \subseteq N$. Then, the core of \mathbf{v} is not empty if and only if $(x_T, A_T)_{T \subseteq N}$ is balanced.*

Proof. For every $T \subseteq N$ the set $\mathbf{v}(T)$ is bounded. Thus, there is a constant $m > 0$ such that set $M := \{x \in \mathbb{R}_+^k; x \leq (m, \dots, m)\}$ contains $\mathbf{v}(T)$ for every $T \subseteq N$: $\mathbf{v}(T) \subseteq M$. Denote $D_T^1 := \text{cl}(M \cap A_T)$ and $D_T^2 := \text{cl}(\mathbb{R}_+^k \setminus D_T^1)$. By assumption, for every $T \subsetneq N$, A_T is convex, hence D_T^1 is compact and convex. Finally, D_T^2 is a polyhedron and $A_T = D_T^1 \cup D_T^2$. This feature of A_T is required in order to employ Theorem 3.

By Theorem 3, there exists an $n \times k$ dimensional matrix C such that

$$x_T C \in A_T \text{ for every } T \subseteq N \quad (19)$$

if and if $(x_T, A_T)_{T \subseteq N}$ is balanced. To complete the proof note that C is a core member if and only if it satisfies Eq. 19. \square

In the current context, the interpretation of the equation $SX = 0$ is the following. As we have seen above, in the one-dimensional case $sX = 0$ means that s is a vector of balanced coefficients. Here $SX = 0$ is interpreted as a multiple balanced coefficients: every row of the matrix S is a vector of balanced coefficients.

¹⁵Note that l in consideration equals to $2^n - 1$, the number of coalitions in the game.

Remark 3. In the case where for every $T \subsetneq N$ the set of the points in \mathbb{R}_+^k that dominate $\mathbf{v}(T)$ is a union of finitely many convex sets, one can list a finite number of problems, namely, $A_T^j \subseteq \mathbb{R}^k, T \subseteq N, j = 1, \dots, L$, such that the core of \mathbf{v} is not empty if and only if there is at least one j and a matrix C such that $x_T C \in A_T^j$ for every $T \subsetneq N$ and $x_T C \in \mathbf{v}(N)$. An example of such a case is where $\mathbf{v}(N)$ is convex and each $\mathbf{v}(T), T \subsetneq N$ is a polytope (i.e., bounded polyhedron).

We conclude that Theorem 4 provides a characterization of the non-emptiness of the core also in cases where each set $\mathbb{R}_+^k \setminus \mathbf{v}(T), T \subsetneq N$ and $\mathbf{v}(N)$ is a finite union of convex sets.

References

- [1] Assa, H., Elliston, S. and E. Lehrer (2016), Joint Games and Compatibility, *Economic Theory*, 61, 91-113.
- [2] Bondareva, Olga N. (1963), Some Applications of Linear Programming Methods to the Theory of Cooperative Games, *Problemy Kybernetiki*, 10, 119-139. (In Russian)
- [3] Bloch, F., and G. de Clippel (2010), Cores of combined games, *Journal of Economic Theory*, 145, 2424-2434.
- [4] Fernandez, F. R., Hinojosa, M. A., and J. Puerto (2004), Set-Valued TU-games, *European Journal of Operational Research*, 159, 181-195.
- [5] Gayer, G. and D. Persitz, Negotiation Across Multiple Issues (2016), *Theoretical Economics*, 11, 937-969.
- [6] Jünger, M., T.M. Lieblich, and D. Naddef (2010) *50 Years of Integer Programming 1958-2008: From the Early Years to the State-of-the-Art*, Dordrecht: Springer.
- [7] Lind, M. (1996) Cooperative Game Theory and Multiple Criteria Decision Making, University of Aarhus, working paper.
- [8] Voorneveld, M. and A. van den Nouweland (1998), Cooperative Multicriteria Games with Public and Private Criteria, working paper.

- [9] Predtetchinski, A. and P. J. J. Herings (2004), A Necessary and Sufficient Condition for Non-Emptiness of the Core of a Non-Transferable Utility Game, *Journal of Economic Theory*, 116, 84-92.
- [10] Shapley, L. S. (1967), On Balanced Sets and Cores, *Naval Research Logistics Quarterly*, 14, 453-460.

8 Appendix

For any two convex sets E, F in the same Euclidean space, define $d(E, F) := \inf\{\|e - f\|; e \in E, f \in F\}$.

Proof of Lemma 1. A and B are disjoint. Thus, in order to prove the lemma it is sufficient to show that $d(A, B) > 0$. If otherwise, $d(A, B) = 0$, we obtain that for every $i = 1, \dots, l$,

$$d(A_i, B_i) = 0, \tag{20}$$

where $B_i = \{x_i C; C \text{ is an } n \times k \text{ matrix}\}$. We show that Eq. 20 implies $A_i \cap B_i \neq \emptyset$. This would imply that A and B are not disjoint, which is a contradiction.

B_i is a subspace while (by assumption) A_i is a union of a convex and compact set, say D_i , and a polyhedron, say P_i . If $d(D_i, B_i) = 0$, then due to the compactness of D_i , $D_i \cap B_i \neq \emptyset$ and the proof is done. It must be then that $d(D_i, B_i) > 0$, while $d(P_i, B_i) = 0$.

By Minkowski-Weyl Theorem (see [6]) P_i , as a polyhedron, is a Minkowski sum of a convex hull of finitely many points,

$$C = \text{conv}\{c_1, \dots, c_s\} = \left\{ \sum_{i=1}^s \alpha_i c_i; \alpha_i \geq 0, \sum_{i=1}^s \alpha_i = 1 \right\},$$

and a finitely generated cone,

$$V = \text{cone}\{v_1, \dots, v_t\} = \left\{ \sum_{i=1}^t \beta_i v_i; \beta_i \geq 0 \right\}.$$

In other words, $P_i = \{c + v; c \in C, v \in V\}$. Since $d(P_i, B_i) = d(C + V, B_i) = 0$, we obtain $d(C, B_i - V) = 0$. However, C is convex and compact and $B_i - V$ is convex and closed, hence $C \cap (B_i - V) \neq \emptyset$. This, in turn, implies that $P_i \cap B_i = (C + V) \cap B_i \neq \emptyset$ which completes the proof. \square