

The de Bruijn-Erdős Theorem for Hypergraphs

Noga Alon^{*} Keith E. Mellinger[†] Dhruv Mubayi[‡] Jacques Verstraëte[§]

May 17, 2011

Abstract

Fix integers $n \geq r \geq 2$. A clique partition of $\binom{[n]}{r}$ is a collection of proper subsets $A_1, A_2, \dots, A_t \subset \binom{[n]}{r}$ such that $\bigcup_i A_i$ is a partition of $\binom{[n]}{r}$.

Let $\text{cp}(n, r)$ denote the minimum size of a clique partition of $\binom{[n]}{r}$. A classical theorem of de Bruijn and Erdős states that $\text{cp}(n, 2) = n$. In this paper we study $\text{cp}(n, r)$, and show in general that for each fixed $r \geq 3$,

$$\text{cp}(n, r) \geq (1 + o(1))n^{r/2} \quad \text{as } n \rightarrow \infty.$$

We conjecture $\text{cp}(n, r) = (1 + o(1))n^{r/2}$. This conjecture has already been verified (in a very strong sense) for $r = 3$ by Hartman-Mullin-Stinson. We give further evidence of this conjecture by constructing, for each $r \geq 4$, a family of $(1 + o(1))n^{r/2}$ subsets of $[n]$ with the following property: no two r -sets of $[n]$ are covered more than once and all but $o(n^r)$ of the r -sets of $[n]$ are covered.

We also give an absolute lower bound $\text{cp}(n, r) \geq \binom{n}{r} / \binom{q+r-1}{r}$ when $n = q^2 + q + r - 1$, and for each r characterize the finitely many configurations achieving equality with the lower bound. Finally we note the connection of $\text{cp}(n, r)$ to extremal graph theory, and determine some new asymptotically sharp bounds for the Zarankiewicz problem.

1 Introduction

A classical theorem of de Bruijn and Erdős [8] states that the minimum number of proper complete subgraphs (henceforth cliques) of the complete graph K_n that are needed to partition its edge set is n . Equality holds only for $n - 1$ copies of K_2 on a vertex together with a clique of order $n - 1$ (a near-pencil) or the cliques whose vertex sets are the $q^2 + q + 1$ point sets of lines in a projective plane of order q when $n = q^2 + q + 1$ and a projective plane exists.

^{*}Schools of Mathematics and Computer Science, Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv 69978, Israel. Email: nogaa@tau.ac.il Research supported in part by an ERC Advanced grant and by a USA-Israeli BSF grant

[†]Department of Mathematics, University of Mary Washington, Fredericksburg, VA 22401. Email: kmelling@umw.edu

[‡]Department of Mathematics, Statistics, and Computer Science, University of Illinois, Chicago, IL 60607. Research supported in part by NSF grants DMS-0653946 and DMS-0969092. Email: mubayi@math.uic.edu

[§]Department of Mathematics, University of California, San Diego, La Jolla CA 92093-0112. Email: jacques@ucsd.edu Research supported in part by NSF grant DMS-0800704 and a Hellman Fellowship

In this paper, we study the analog of the de Bruijn-Erdős theorem for hypergraphs. Write $\binom{X}{r}$ for the collection of r -elements subsets of a set X ; this is a clique. Throughout this paper, we write $[n] := \{1, 2, \dots, n\}$. We will associate a hypergraph with its edge set.

Definition. Fix integers $n \geq r \geq 2$. A *clique partition* of a hypergraph H is a collection of proper subcliques that partition H . Let $\text{cp}(n, r)$ be the minimum size of a clique partition of $\binom{[n]}{r}$. In other words, $\text{cp}(n, r)$ is the minimum t for which there are proper subsets $A_1, A_2, \dots, A_t \subset [n]$ such that $\bigcup_i \binom{A_i}{r}$ is a partition of $\binom{[n]}{r}$.

The de Bruijn-Erdős Theorem can now be restated as $\text{cp}(n, 2) = n$ together with the characterization of equality. In this paper, we consider $\text{cp}(n, r)$ for $r > 2$. A minimum clique partition of $\binom{[n]}{r}$ shall be referred to as an *optimal clique partition*.

As noted above, there are essentially two types of configurations achieving equality in the de Bruijn-Erdős theorem: near pencils and projective planes. For $r = 3$ and $n = q^2 + 1$ there is only one type of configuration that achieves equality, and this is called an *inversive plane*.

An *inversive plane* is a pair (V, \mathcal{C}) where V is a set of points and \mathcal{C} is a set of subsets of points called circles satisfying the following axioms:

- (1) any three points of V must lie in exactly one circle in $\mathcal{C} \in \mathcal{C}$,
- (2) four points in V must exist that are not contained in a common circle and
- (3) for any circle $C \in \mathcal{C}$ and points $p \in C$ and $q \in V \setminus C$, there is exactly one circle $D \in \mathcal{C}$ such that $p, q \in D$ and $C \cap D = \{p\}$.

It is known that if an inversive plane on n points exists, then n is necessarily of the form $q^2 + 1$ for some integer $q \geq 2$, and the number q is called the *order* of the inversive plane.

If the cliques in a clique partition form an inversive plane, then we identify the partition with the inversive plane. The problem of determining $\text{cp}(n, 3)$ is quite well understood due to the following theorem.

Theorem 1 (Hartman-Mullin-Stinson [17]) *Let $q \geq 3$ be an integer and $n = q^2 + 1$. Then*

$$\text{cp}(n, 3) \geq n\sqrt{n-1}.$$

If q is a prime power, then

- $\text{cp}(n, 3) = qn = n\sqrt{n-1}$, and
- if P is a clique partition of $\binom{[n]}{3}$, $|P| = qn$, then P is an inversive plane of order q .

Consequently, as $n \rightarrow \infty$

$$\text{cp}(n, 3) \sim n^{3/2}.$$

There are many models for finite inversive planes, and the reader is referred to the classic book by Dembowski [9], the very complete book by Benz [5], or the more recent survey by Wilker [29] for details and examples.

In [17], $\text{cp}(n, 3)$ is determined when n is close to $q^2 + 1$ and q is a prime power, due to the existence of inversive planes and classification [26] of linear spaces with n points and m blocks such that $(m - n)^2 \leq n$. In general, however, it is likely to be challenging to determine $\text{cp}(n, 3)$ exactly for all n , in contrast to the de Bruijn-Erdős Theorem.

1.1 Clique partitions of $\binom{[n]}{r}$

The problem of determining $\text{cp}(n, r)$ for $r > 3$ appears to be difficult. First we present a lower bound for $\text{cp}(n, r)$ in terms of $\text{cp}(n - 1, r - 1)$:

Theorem 2 *Let $r > 2$ and $n > r$ be integers. If $\text{cp}(n - 1, r - 1) = \delta \text{cp}(n, r)$ then*

$$\text{cp}(n, r) \binom{\delta n}{r} \leq \binom{n}{r}. \quad (1)$$

Consequently, for each fixed $r \geq 2$, and $n \rightarrow \infty$,

$$\text{cp}(n, r) \geq (1 - o(1))n^{r/2}.$$

Via constructions we shall give evidence that this lower bound may be the true asymptotic behavior of $\text{cp}(n, r)$ for every fixed $r \geq 2$.

Theorem 3 *For every fixed $r \geq 2$ there is a family $H \subseteq \binom{[n]}{r}$ of r -element sets such that $|H| = \binom{n}{r} - o(n^r)$ and H has a clique partition with $(1 + o(1))n^{r/2}$ cliques.*

It appears to be difficult to extend the constructions for Theorem 3 to full partitions of $\binom{[n]}{r}$ without adding many more subcliques. We nevertheless conjecture that in general, the above lower bound is asymptotically sharp:

Conjecture 4 *For every fixed $r \geq 2$ we have $\text{cp}(n, r) = (1 + o(1))n^{r/2}$ as $n \rightarrow \infty$.*

1.2 Characterization of clique partitions

Equality holds in the de Bruijn-Erdős Theorem for projective planes and near-pencils. The nature of clique partition numbers $\text{cp}(n, r)$ for $r > 2$ surely depends on number theoretic properties of n , so unlike in the de Bruijn-Erdős Theorem, a characterization of optimal partitions for all n is likely to be more difficult.

Towards this goal, we prove the following theorem, which considers values of n that are parameterized in a way that includes the de Bruijn-Erdős Theorem as a special case. As is customary, we define the binomial coefficient $\binom{x}{r} = x(x - 1) \cdots (x - r + 1)/r!$ for any positive integer r and real number x . Also, recall that a *Steiner (n, k, t) -system* is a collection of k -element sets of $[n]$ such that every t -element subset of $[n]$ lies in precisely one of the k -element sets.

Theorem 4 *Let $r \geq 2$ and let n be a positive integer. Define the positive real number q by the equation $n = q^2 + q + r - 1$. Then*

$$\text{cp}(n, r) \geq \frac{\binom{n}{r}}{\binom{q+r-1}{r}}. \quad (2)$$

Equality holds if and only if one of the following holds:

- $n = r + 1$ and the partition is $\binom{[r+1]}{r}$
- $n > r + 1$ and $r = 2$ and the partition is a projective plane of order q or a near pencil
- the partition is a Steiner (n, k, r) -system where $(n, k, r) \in \{(8, 4, 3), (22, 6, 3), (23, 7, 4), (24, 8, 5)\}$.

Note that in the third case above such Steiner systems are known to exist for $r \in \{3, 4, 5\}$, and so for those values of r we have a complete characterization of equality in Theorem 4. The case $r = 2$ is the de Bruijn-Erdős Theorem. We note that not a single construction of a Steiner (n, m, r) -system is known for any $n > m > r > 5$. This is considered to be one of the major open problems in design theory [28].

1.3 Zarankiewicz problem

There is a tight connection between $\text{cp}(n, r)$ and the Zarankiewicz problem from extremal graph theory. The *Zarankiewicz number* $z(m, n, s, t)$ denotes the maximum possible number of 1s in an $m \times n$ matrix containing no $s \times t$ minor consisting entirely of 1s. This can be rephrased in terms of the maximum number of edges in an $m \times n$ bipartite graph containing no complete bipartite subgraph with s vertices in the part of size m and t vertices in the part of size n . Our results for the clique partition number $\text{cp}(n, r)$ imply the following new asymptotically sharp results for Zarankiewicz numbers.

Theorem 5 *Fix $r \geq 3$. If $m = (1 + o(1))n^{r/2}$, then $z(m, n, 2, r) = (1 + o(1))\sqrt{n}m$ as $n \rightarrow \infty$.*

The special case $r = 3$ had earlier been shown by Alon-Rónyai-Szabó [2], as part of a more general result motivated by a question in discrepancy theory posed by Matoušek. However, our constructions here are different.

Theorem 5 is in contrast to a result of Füredi [14] which shows that for each fixed $r \geq 2$, we have $z(m, n, 2, r) = (1 + o(1))\sqrt{(r-1)n}m$ whenever $m = (1 + o(1))n$. The problem of determining the asymptotic behavior of $\text{cp}(n, r)$ for fixed $r \geq 2$ seems more challenging than that of determining the Zarankiewicz numbers $z(m, n, 2, r)$ for $m = (1 + o(1))n^{r/2}$.

1.4 Organization

In Section 2, we discuss $\text{cp}(n, r)$ for $r > 3$, starting in Section 2.1 with the proof of Theorem 2, proceeding in Section 2.2 with the proof of Theorem 3, and mentioning an alternative construction

for $r \in \{4, 5\}$ in Section 2.3. In Section 3, we prove Theorem 4, and in Section 4 we point out the connection to the Zarankiewicz problem and prove Theorem 5.

2 Clique partitions of $\binom{[n]}{r}$

In this section, we prove Theorems 2 and 3.

2.1 Proof of Theorem 2

Let P be an optimal clique partition of $\binom{[n]}{r}$ which has vertex set $[n]$. Then $P_v^* = \{C \setminus \{v\} : v \in C \in P\}$ is a clique partition of $\binom{[n-1]}{r-1}$ so $|P_v^*| \geq \text{cp}(n-1, r-1)$. Noting the identity

$$\sum_{C \in P} |C| = \sum_{v \in [n]} |P_v^*|,$$

we have

$$\sum_{C \in P} |C| \geq n \text{cp}(n-1, r-1). \quad (3)$$

On the other hand since P is a clique partition of $\binom{[n]}{r}$,

$$\sum_{C \in P} \binom{|C|}{r} = \binom{n}{r}. \quad (4)$$

By convexity of binomial coefficients, the right hand side is a minimum when the $|C|$ all equal their average value, which by (3) is at least $n \text{cp}(n-1, r-1) / \text{cp}(n, r) = \delta n$ by definition of δ . Inserting this in (4) gives (1). \square

2.2 Proof of Theorem 3

By the known results about the distribution of primes it suffices to prove the result for $n = q^2$, where q is a prime power. For other values of n we can then take the construction for the smallest $n' > n$ satisfying $n' = q'^2$, with q' being a prime power, and consider the induced constructions on n vertices among those n' .

Suppose, thus, that $n = q^2$, and let $F = GF(q)$ denote the finite field of size q . Let V be the set of all ordered pairs (x, y) with $x, y \in F$. For every polynomial $p(x)$ of degree at most $r-1$ over F , let C_p denote the subset $C_p = \{(x, p(x)) : x \in F\}$ of V . The collection \mathcal{C} of all sets C_p is a collection of $q^r = n^{r/2}$ subsets of V , each of size $q = \sqrt{n}$. Clearly, no two members of \mathcal{C} share more than $r-1$ elements, as two distinct polynomials of degree at most $r-1$ can have the same value in at most $r-1$ points. Moreover, every set $\{(x_i, y_i) : 1 \leq i \leq r\}$ of r points of V in which the x_i -s are pairwise distinct is contained in a unique set C_p , as there is a unique polynomial p of degree at most $r-1$ satisfying $p(x_i) = y_i$ for all i . Let H be the set of all r -subsets of V contained in a member of \mathcal{C} . Then H has a clique partition with the $n^{r/2}$ cliques C_p , and the only r -sets

in V that do not belong to H are those that have at least two points (x, y) with the same first coordinate. The number of these r -sets is at most

$$(1 + o(1))q \binom{q}{2} \binom{q-1}{r-2} q^{r-2} = (1 + o(1)) \frac{1}{2(r-2)!} q^{2r-1} = O(n^{r-1/2}) = o(n^r),$$

completing the proof. \square

2.3 An alternative construction for $r \in \{4, 5\}$

In this subsection we present an alternative construction that proves the assertion of Theorem 3 for $r \in \{4, 5\}$. Although this is less general than the previous construction we believe it is interesting and may provide some extra insight.

We work in the classical finite projective plane of order q , denoted by $\pi = PG(2, q)$. The plane π contains $q^2 + q + 1$ points that can be represented by homogeneous coordinates (x, y, z) . A non-degenerate conic of π is a collection of points whose homogeneous coordinates satisfy some non-degenerate quadratic form, and it is well-known that there is exactly one such conic in π , up to isomorphism. A typical example is the set of points satisfying the form $y^2 = xz$ which contains the points $\{(0, 0, 1)\} \cup \{(1, x, x^2) : x \in \mathbb{F}_q\}$. An arc is a set of points, no three collinear, and it is straightforward to show that conics form arcs. Moreover, a classical result of Segre [23] says that when q is odd, every set of $q + 1$ points, no three collinear, is in fact a conic.

For an overview of results on conics and arcs in general, the reader is referred to Chapters 7 and 8 of [18]. It is well-known that there are precisely $q^5 - q^2$ conics in $PG(2, q)$ and that five points in general position (that is, no three of which are collinear) determine a unique conic. We can use these facts to cover 4-sets and 5-sets of π .

The case $r = 4$. Distinguish a special point P of π and consider the set of all non-degenerate conics passing through P . Our points will be the points of $\pi \setminus \{P\}$. Hence, $n = q^2 + q$. Let X be the number of non-degenerate conics passing through an arbitrary point Q of π . By the transitive properties of $Aut(\pi)$, it follows that X is independent of Q . We count pairs (Q, \mathcal{C}) with Q a point of the conic \mathcal{C} , by first counting Q , and then counting \mathcal{C} . This give us

$$(q^2 + q + 1)X = (q^5 - q^2)(q + 1) = q^2(q - 1)(q^2 + q + 1)(q + 1).$$

It follows that there are $X = q^4 - q^2$ conics through a point of π . The number of 4-sets in the set $\pi \setminus \{P\}$ is $\binom{q^2+q}{4} = (\frac{1}{24} + o(1))q^8$. The number of 4-sets of $\pi \setminus \{P\}$ covered by the conics through P is $(q^4 - q^2)\binom{q}{4} = (\frac{1}{24} + o(1))q^8$ (the difference is asymptotic to $\frac{10}{24}q^7$ which is of a lower order as $q \rightarrow \infty$). Hence, we have shown that for $n = q^2 + q$, there is a collection of $(1 - o(1))q^4 = (1 - o(1))n^2$ proper edge-disjoint subcliques of $\binom{[n]}{4}$ that cover all but $o(q^8) = o(n^4)$ edges of $\binom{[n]}{4}$.

The case $r = 5$. We can repeat our construction, using all points and conics of π , to obtain a similar result for $r = 5$. The construction here is in fact simpler than the one for $r = 4$. The total number of 5-sets of points in π is $\binom{q^2+q+1}{5} = (\frac{1}{120} + o(1))q^{10}$. The number of 5-sets covered by conics is $(q^5 - q^2) \times \binom{q+1}{5} = (\frac{1}{120} + o(1))q^{10}$ (the difference is asymptotically $\frac{1}{12}q^9$ which is of lower order).

Hence, we have shown that for $n = q^2 + q + 1$, there is a collection of $(1 - o(1))q^5 = (1 - o(1))n^{5/2}$ proper edge-disjoint subcliques of $\binom{[n]}{5}$ that cover all but $o(q^{10}) = o(n^5)$ edges of $\binom{[n]}{5}$. This completes the (second) proof of Theorem 3 for $r \in \{4, 5\}$. \square

3 Proof of Theorem 4

We first address the lower bound in Theorem 4. Before beginning the proof we need the following simple lemma, known as Chebyshev Sum Inequality (c.f., e.g., [16].)

Lemma 6 (Chebyshev Sum Inequality) *Suppose that $p > 1$ is an integer and $f, g : [p] \rightarrow \mathbb{Z}^+$ are non-decreasing functions. Then*

$$\sum_{i=1}^p f(i)g(i) \geq \frac{1}{p} \sum_{i=1}^p f(i) \sum_{i=1}^p g(i).$$

There are several simple proofs of this inequality. A short one is to observe that the right hand side is the expectation of the sum $\sum_{i=1}^p f(i)g(\sigma(i))$, where σ is a random uniformly chosen permutation of $\{1, 2, \dots, p\}$, and that the maximum value of this sum is obtained when σ is the identity, since if $\sigma(i) > \sigma(j)$ for some $i < j$, then the permutation σ' obtained from σ by swapping the values of $\sigma(i)$ and $\sigma(j)$ satisfies $\sum_{i=1}^p f(i)g(\sigma'(i)) \geq \sum_{i=1}^p f(i)g(\sigma(i))$.

3.1 Proof of the lower bound

In this section we show the lower bound in Theorem 4. Define $(n)_r := n(n-1)\dots(n-r+1)$ and $\phi(n, r) = \binom{n}{r} / \binom{q+r-1}{r}$. Let P be a clique partition of $\binom{[n]}{r}$ where $n = q^2 + q + r - 1$ and q is a positive real number. We aim to show $|P| \geq \phi(n, r)$, and we proceed by induction on r . For $r = 2$, this follows from the de Bruijn-Erdős Theorem since $\phi(n, 2) = q^2 + q + 1 = n$. Since $\phi(r+1, r) = r+1$, we may assume $n > r+1$. Suppose $r > 2$ and let C_1, C_2, \dots, C_p be the cliques in P , of respective sizes $n_1 \leq n_2 \leq \dots \leq n_p$. Note as in (3),

$$\sum_{i=1}^p n_i \geq n\phi(n-1, r-1). \quad (5)$$

For a set $S \subset [n]$, let $P_S^* = \{C \setminus S : C \in P\}$. Then

$$\sum_{i=1}^p \binom{n_i}{r-2} = \sum_{S \in \binom{[n]}{r-2}} |P_S^*|.$$

By the de Bruijn-Erdős Theorem, since P_S^* is a clique partition of $\binom{[n-r+2]}{2}$, $|P_S^*| \geq n-r+2$ and therefore

$$\sum_{i=1}^p \binom{n_i}{r-2} \geq \binom{n}{r-2} (n-r+2) \quad (6)$$

with equality if and only if there is a projective plane on $n - r + 2$ points, q is an integer, and every clique in P has size $q + r - 1$. Note here that we are ruling out the possibility of a near pencil in every P_S^* , otherwise P would have a clique of size $n - 1$ and then $|P| \geq 1 + \binom{n-1}{r-1} > \phi(n, r)$ since $n > r + 1$. On the other hand since P is a clique partition of $\binom{[n]}{r}$,

$$\sum_{i=1}^p \binom{n_i}{r} = \binom{n}{r}. \quad (7)$$

Since $n_i \geq r$ for all i , we may apply Lemma 6 with $f(i) = (n_i)_{r-2}$ and $g(i) = (n_i - r + 2)(n_i - r + 1)$ to obtain

$$\sum_{i=1}^p (n_i)_r \geq \frac{1}{p} \sum_{i=1}^p (n_i)_{r-2} \cdot \sum_{i=1}^p (n_i - r + 2)(n_i - r + 1).$$

By (6) and (7), we have

$$p \geq \frac{1}{n - r + 1} \sum_{i=1}^p (n_i - r + 2)(n_i - r + 1).$$

The function $h(x) = (x - r + 2)(x - r + 1)$ is convex for $x \geq r$. By Jensen's Inequality which asserts that the average value of a convex function over a set of points is at least the value of the function in the average, we conclude that

$$\sum_{i=1}^p (n_i - r + 2)(n_i - r + 1) \geq p(c - r + 2)(c - r + 1)$$

where c is the average size of a clique in P . Consequently,

$$n - r + 1 \geq (c - r + 2)(c - r + 1)$$

and since $n = q^2 + q + r - 1$, we find $c \leq q + r - 1$. Together with (5) we get

$$p(q + r - 1) \geq n\phi(n - 1, r - 1).$$

Using the identity $\phi(n, r)(q + r - 1) = n\phi(n - 1, r - 1)$, this gives $p \geq \phi(n, r)$ and completes the proof of the lower bound in Theorem 4.

3.2 Equality in Theorem 4

For the constructions giving equality in Theorem 4, we note that for $r = 2$ this is exactly the de Bruijn-Erdős Theorem. Moving on to $r \geq 3$, we have $n = q^2 + q + r - 1$, and equality holds in the above arguments if and only if equality holds in (6), which means that every $(r - 2)$ -set S has the property that P_S^* is an optimal partition $\binom{[n] \setminus S}{2}$. Therefore every r -set is covered at most (in fact exactly) once and P is a Steiner $(n, q + r - 1, r)$ -system. It is well known (see [28]) that the following divisibility requirements are necessary for the existence of such Steiner systems:

$$\binom{q + r - 1 - i}{r - i} \mid \binom{n - i}{r - i} \quad \text{for } i = 0, 1, \dots, r - 1. \quad (8)$$

Equivalently, since $n = q^2 + q + r - 1$, (8) is equivalent to:

$$\prod_{j=1}^{r-i} (q + r - i - j) \mid \prod_{j=1}^{r-i} (q^2 + q + r - i - j) \quad \text{for } i = 0, 1, \dots, r - 1. \quad (9)$$

For each r , let Q_r be the set of values of q permitted by (9). Then the Q_r form a decreasing chain $Q_3 \supseteq Q_4 \supseteq \dots$ and $1 \in Q_r$ for all $r \geq 3$. Now Q_3 is the set of q such that

$$\begin{aligned} q & \mid (q^2 + q) \\ q(q+1) & \mid (q^2 + q + 1)(q^2 + q) \\ q(q+1)(q+2) & \mid (q^2 + q + 2)(q^2 + q + 1)(q^2 + q). \end{aligned}$$

The first two conditions are trivially satisfied, and the last is

$$(q+2) \mid (q^2 + q + 2)(q^2 + q + 1).$$

This is equivalent to $(q+2) \mid 12$. It follows that $q \in \{1, 2, 4, 10\}$, $Q_3 = \{1, 2, 4, 10\}$ and then $n \in \{4, 8, 22, 112\}$. Since Steiner $(n, q+2, 3)$ -systems exist for $(n, q) \in \{(4, 1), (8, 2), (22, 4)\}$ (see van Lint and Wilson [28]), Theorem 2 is tight in those cases and

$$\text{cp}(4, 3) = 4 \quad \text{cp}(8, 3) = 14 \quad \text{cp}(22, 3) = 77.$$

The case $q = 10$ may be ruled out, since it is accepted (see [21]) that a projective plane of order ten does not exist, hence it is impossible that P_S^* is a partition of $\binom{[n] \setminus S}{2}$ for $|S| = 1$. For $r = 4$, we have the same divisibility requirements as $r = 3$ together with

$$q(q+1)(q+2)(q+3) \mid (q^2 + q + 3)(q^2 + q + 2)(q^2 + q + 1)(q^2 + q).$$

Equivalently, we are looking for $q \in Q_3$ satisfying

$$(q+2)(q+3) \mid (q^2 + q + 3)(q^2 + q + 2)(q^2 + q + 1).$$

This is not satisfied for $q \in \{2, 10\}$ and hence the only values $q \in Q_3 = \{1, 2, 4, 10\}$ which satisfy these requirements are $q \in \{1, 4\}$ and $Q_4 = \{1, 4\}$. Since a Steiner $(23, 7, 4)$ -system exists which corresponds to $q = 4$, we have

$$\text{cp}(5, 4) = 5 \quad \text{cp}(23, 4) = 253.$$

For $r = 5$, we are seeking Steiner $(q^2 + q + 4, q + 4, 5)$ -systems, which implies $q \in \{1, 4\} = Q_5$ and $n \in \{6, 24\}$. Since a Steiner $(24, 8, 5)$ -system exists, we have the complete solution

$$\text{cp}(6, 5) = 6 \quad \text{cp}(24, 5) = 759.$$

Finally we show $Q_r = \{1\}$ for $r \geq 6$ to complete the proof. First we have $Q_5 = \{1, 4\}$ as just seen, and since $Q_r \subseteq Q_5$ for all $r \geq 5$, we only have to show $Q_6 = \{1\}$. If $4 \in Q_6$, then the divisibility requirement (9) with $i = 0$ and $q = 4$ is

$$9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \mid 25 \cdot 24 \cdot 23 \cdot 22 \cdot 21 \cdot 20.$$

This is false, since 3 is a prime factor with multiplicity three on the left, and only two on the right. Therefore $Q_r = \{1\}$ for $r \geq 6$, and the only possible clique partition achieving equality in Theorem 4 has $q = 1$ and $n = r + 1$, and therefore it must be $\binom{[r+1]}{r}$. \square

4 Zarankiewicz Problem and Theorem 5

In this section we point out the connection between Zarankiewicz numbers and clique partitions. As a byproduct, we prove Theorem 5. Recall that the Zarankiewicz number $z(m, n, s, t)$ is the maximum number of edges in an $m \times n$ bipartite graph containing no complete bipartite subgraph with s vertices in the part of size m and t vertices in the part of size n . The clique partition number $\text{cp}(n, r)$ is related to the Zarankiewicz numbers in the following sense:

Lemma 7 *Let $n \geq r$ be a positive integer and $m = \binom{n}{r} / \binom{k}{r}$. Then there is a partition of $\binom{[n]}{r}$ into m cliques of size k if and only if $z(m, n, 2, r) = km$.*

Proof: We follow the classical argument of [20]. For any $m \times n$ bipartite graph with parts A and B , not containing a $2 \times r$ complete bipartite subgraph (with the two vertex set lying in A),

$$\sum_{a \in A} \binom{d(a)}{r} \leq \binom{|B|}{r}. \quad (10)$$

By convexity, if e is the number of edges in the graph we obtain

$$m \binom{e/m}{r} \leq \binom{n}{r}. \quad (11)$$

Suppose $e = z(m, n, 2, r)$ and $e > km$. Then the above formula becomes

$$m \binom{k}{r} < \binom{n}{r}.$$

However this contradicts the identity relating m, n and k in the lemma. It follows that $z(m, n, 2, r) \leq km$.

Now consider the incidence graph of points and sets of size k in the clique partition of $\binom{[n]}{r}$. This is an $m \times n$ bipartite graph, and since we have a clique partition, it does not contain a $2 \times r$ complete bipartite subgraph. Therefore $z(m, n, 2, r) \geq km$.

If $z(m, n, 2, r) = km$, then equality holds in (10) and (11), which means that $d(a) = k$ for every $a \in A$ and every r -set of B is covered exactly once. This gives us a partition of $\binom{[n]}{r}$ into m cliques of size k . \square

The proof of Lemma 7 can be easily rewritten to prove the following: Fix $r \geq 2$ and let $H \subset \binom{[n]}{r}$ satisfy $|H| \geq \binom{n}{r} - o(n^r)$. Suppose that there is a partition of H into $m = (1 + o(1)) \binom{n}{r} / \binom{k}{r}$ cliques of size k . Then $z(m, n, 2, r) = (1 + o(1))km$.

Using the preceding results and constructions in Sections 2 and 4, we obtain the following theorem on Zarankiewicz numbers which proves Theorem 5.

Theorem 8 *Let $n = q^2 + 1$ and $m = qn$ where q is a prime power. Then $z(m, n, 2, 3) = (q^2 + q)n$. Furthermore, if $r \geq 4$, n is an integer and $m = (1 + o(1))n^{r/2}$, then $z(m, n, 2, r) = (1 + o(1))\sqrt{n}m$ as $n \rightarrow \infty$.*

Proof: In the first case we apply Lemma 7 with $k = q + 1$ and $r = 3$. Such a clique partition exists by the construction of inversive planes. Then we see that $m = qn = \binom{n}{3} / \binom{q+1}{3}$ and so we immediately get $z(m, n, 2, 3) = km = (q + 1)qn = (q^2 + q)n$.

For the cases $r \geq 4$ we use the asymptotic version of Lemma 7 above and the results in Section 3.2. If $n = q^2$ with q a prime power, the desired result follows from the construction in the proof of Theorem 3. For other values of n we take the smallest prime power q so that $q^2 \geq n$, consider the construction for that which is a bipartite graph with vertex classes A and B , and take the expected number of edges in an induced subgraph on randomly chosen sets of sizes n and m in A and B , respectively. \square

5 Concluding remarks

• Clique partitions and extensions of Fisher's Inequality

The de Bruijn-Erdős Theorem is an extension of a well known inequality of Fisher [12] that asserts that any nontrivial clique partition of $\binom{[n]}{2}$ in which all cliques have the same cardinality contains at least n cliques. Fisher's Inequality has been extended in several ways. One such extension is due to Ray-Chaudhuri and Wilson [22] who proved that for even r , any clique partition of $\binom{[n]}{r}$ in which all cliques have the same cardinality contains at least $\binom{n}{r/2}$ cliques. Our bound here is stronger (by a factor of $(1 + o(1))(r/2)!$ for even r), and applies to odd values of r as well without having to assume that all cliques have the same cardinality. On the other hand, the result in [22], whose proof is algebraic, holds even if every r -set is covered exactly λ times for some $\lambda > 0$.

• Partitions into complete r -partite r -graphs

A well known result of Graham and Pollak [15] asserts that the complete graph on n vertices can be edge partitioned into $n - 1$, but not less, complete bipartite graphs. This can be viewed as a bipartite analogue of the de Bruijn-Erdős Theorem. In the bipartite case, unlike the one dealing with clique-partitions, there are many extremal configurations, and the only known proofs of the lower bound are algebraic (though recently Vishwanathan [27] presented a counting argument that replaces the linear algebraic part of one of these proofs).

The r -partite version of the problem considered here is to determine the minimum possible number of complete r -partite r -graphs in a decomposition of the edges of the complete r -graph on n vertices. More formally, this is the minimum $p = p(n, r)$ so that there are p collections of the form $\{A_1^{(i)}, A_2^{(i)}, \dots, A_r^{(i)}\}$, ($1 \leq i \leq p$), satisfying $A_j^{(i)} \cap A_s^{(i)} = \emptyset$ for all $1 \leq i \leq p$ and all $1 \leq j < s \leq r$, $A_j^{(i)} \subset [n]$ for all admissible i and j , and for every r -subset R of n there is a unique i , $1 \leq i \leq p$ so that $|R \cap A_j^{(i)}| = 1$ for all $1 \leq j \leq r$. Thus, the Graham Pollak result asserts that $p(n, 2) = n - 1$. In [1] it is proved that $p(n, 3) = n - 2$ and that for every fixed r there are two positive constants $c_1(r), c_2(r)$ so that $c_1(r)n^{\lfloor r/2 \rfloor} \leq p(n, r) \leq c_2(r)n^{\lfloor r/2 \rfloor}$ for all n (see also [6] for slight improvements). Note that for even r the exponent of n is the same as the one appearing in our bounds for the clique partition problem (that is, the function $\text{cp}(n, r)$ discussed here), but for odd values of r , and in particular for $r = 3$ where both functions are well understood, the exponents differ.

• **Geometric description of inversive planes.**

Inversive planes are more widely modeled using the techniques of finite projective geometry. For completeness, we briefly describe the known models here since these planes provide optimal configurations. Inversive planes can be constructed from ovoids. These are sometimes called egg-like inversive planes (or Miquelian planes when the ovoid is an elliptic quadric as we describe below). Thas [24] showed that for odd q , if P is an optimal clique partition of $\binom{[n]}{3}$ and $n = q^2 + 1$, and for some $v \in [n]$ the vertex sets $V(C) \setminus \{v\} : C \in P$, with $v \in C$, constitute the point sets of the lines of the affine plane $AG(2, q)$, then the vertex sets of the cliques in P are the circles of the Miquelian inversive plane.

To describe the ovoidal construction, the setting is finite projective 3-space $PG(3, q)$. An elliptic quadric of $PG(3, q)$ forms a set of $q^2 + 1$ points, no three of which are collinear; in general, such a set of points in $PG(3, q)$ is called an *ovoid*. An exception occurs when $q = 2$. In this case, an ovoid of $PG(3, 2)$ is a set of five points, no four on a common plane. Now consider the points of an ovoid \mathcal{O} of $PG(3, q)$ as the vertices of $\binom{[q^2+1]}{3}$ together with the non-tangent planar cross sections of \mathcal{O} . As three points uniquely determine a plane, each set of three points of \mathcal{O} (each edge of the hypergraph) is covered by one of these planar cross sections. Moreover, it is well-known that each non-tangent plane meets \mathcal{O} in $q + 1$ points (see, e.g., [19]), forming a planar arc. When \mathcal{O} is an elliptic quadric, these planar cross sections are conics, objects we described in Section 2.3. It can be shown that the set of points of \mathcal{O} together with the non-tangent planar cross-sections of \mathcal{O} forms an inversive plane.

It is an important open question in projective geometry as to whether for any q , an inversive plane of order q must be constructed from an ovoid as above. Barlotti [3] showed that if q is odd, then any ovoid is projectively equivalent to an elliptic quadric – the set of points satisfying a quadratic form $xy + \phi(w, z)$ (here ϕ is an irreducible quadratic form over \mathbb{F}_q). In this case, there is an alternative description: consider the ground set to be $\mathbb{F}_{q^2} \cup \{\infty\}$ and consider the images of the set $\mathbb{F}_q \cup \{\infty\}$ under the group of permutations

$$PGL(2, q^2) = \left\{ z \mapsto \frac{az + b}{cz + d} : ad - bc \neq 0 \right\}.$$

The $q^3 + q$ images are taken to be the circles, each with $q + 1$ points, and it can be shown that the elliptic quadric construction is equivalent to the linear fractional group construction. Note that this model can also be viewed as the points of $PG(1, q^2)$ with blocks corresponding to the Baer sublines (i.e., copies of $PG(1, q)$) therein.

For q even there exist ovoids that are not elliptic quadrics, the so-called Tits ovoids [25]. However, Dembowski's Theorem [9] shows that if q is even, then any inversive plane is derived from an ovoid as above, and so in this sense, optimal clique partitions of $\binom{[n]}{3}$ when $n = q^2 + 1$ and $q = 2^h$ for $h \geq 2$ are unique up to the choice of the ovoid. It follows from the previously mentioned result of Thas [24], translated into the language of clique partitions, that for odd q , if P is an optimal clique partition of $\binom{[n]}{3}$ and $n = q^2 + 1$, and some P_v^* is the classical affine plane $AG(2, q)$, then P itself is constructed as above from an elliptic quadric. In particular, the optimal clique partitions are unique. It appears difficult, however, to claim that any of the clique partitions P_v^* are isomorphic to the classical affine plane. Much more on finite projective 3-space, including results on ovoids and quadrics, can be found in the book by Hirschfeld [19].

• **Classification of linear spaces.** Theorem 2 gives a general lower bound on $\text{cp}(n, r)$ in terms of $\text{cp}(n-1, r-1)$. We concentrate on the case $r = 3$. A necessary condition for tightness in Theorem 4 for $r = 3$ when $n = q^2 + q + 2$ and q is a positive integer is the existence of a projective plane of order q on $n-1$ points. If n is an integer such that no projective plane on $n-1$ points exists, this raises the question as to the minimum linear spaces which are not projective planes and not near pencils on $n-1$ points. A complete analysis of linear spaces on v points with b blocks such that $(b-v)^2 \leq v$ was carried out by Totten [24] (1976) (see also [13]). In particular, those linear spaces are one of the following:

- (a) near pencils
- (b) an affine plane of order q with a linear space on at most $q+1$ new points at infinity (add a point to every line in a parallel class, and then amongst the new points create a linear space of lines – this in particular contains the projective planes when $q+1$ points are added and a single line through them is added)
- (c) embeddable in a projective plane (delete at most $v+1$ points from a projective plane, deleting any line which becomes a singleton)
- (d) an exceptional configuration with $v = 6$ and $b = 8$.

We point out that Theorem 2 can be tight. Suppose $n = 21$. By the classification of linear spaces, $|P_v^*|$ is at least the number of lines in an affine plane of order $q = 4$ together with the trivial linear space on q points at infinity, so $\text{cp}(n-1, 2) \geq q^2 + q + 1$ i.e. $\text{cp}(20, 2) \geq 21$. Using this in Theorem 2, if $x = \text{cp}(21, 3)$ we obtain

$$x \cdot \binom{(21)^2/x}{3} \leq \binom{21}{3}$$

which gives

$$x \geq \frac{441}{676}(\sqrt{32361} - 63) > 76.$$

Since x is an integer, $x \geq 77$. However (see Theorem 4) $\text{cp}(22, 3) = 77$ due to the existence of the Steiner $(22, 6, 3)$ -system, so we conclude $x \leq \text{cp}(22, 3) = 77$. It follows that $\text{cp}(21, 3) = 77$. While it is possible to generalize the argument we just used for $n = 21$ by using the classification of linear spaces, it is the lack of constructive upper bounds where more work is needed, and in general there is a gap between the upper and lower bounds for $\text{cp}(n, 3)$. An extensive survey of the existence problem for Steiner systems may be found in [7].

6 Acknowledgments

We thank the referees for their comments that helped improve the presentation.

References

- [1] N. Alon, Decomposition of the complete r -graph into complete r -partite r -graphs, *Graphs and Combinatorics* 2 (1986), 95–100.

- [2] N. Alon, L. Rónyai, T. Szabó, Norm-graphs: variations and applications. *J. Combin. Theory Ser. B* 76 (1999) no. 2, 280–290.
- [3] A. Barlotti, Un' estensione del teorema di Segre-Kustaanheimo, *Boll. U.M.I.*, 10 (1955), 498-506.
- [4] W. Benz, Ueber Möbiusebenen. Ein Bericht, *Jahresb. D.M.V.*, 63 (1960) 1–27.
- [5] W. Benz, Vorlesungen über Geometrie der Algebren, Geometrien von Möbius, Laguerre-Lie, Minkowski in einheitlicher und grundlagengeometrischer Behandlung. Die Grundlehren der mathematischen Wissenschaften, Band 197. Springer-Verlag, Berlin-New York (1973).
- [6] S. Cioaba, A. Kündgen, J. Verstraëte, On decompositions of complete hypergraphs, *J. Combin. Theory Series A*, 116 (2009), 1232–1234.
- [7] C. Colbourn, R. Mathon, Steiner systems: in *Handbook of Combinatorial Designs*, second edition, (2007) 102–110.
- [8] N. G. de Bruijn, P. Erdős, On a combinatorial problem. *Nederl. Akad. Wetensch., Proc.* 51, (1948) 1277–1279 = *Indagationes Math.* 10, (1948) 421–423.
- [9] P. Dembowski, Inversive planes of even order. *Bull. Amer. Math. Soc.* 69 (1963) 850–854.
- [10] P. Dembowski, D. R. Hughes, On finite inversive planes. *J. London Math. Soc.* 40 (1965) 171–182.
- [11] P. Erdős, R. C. Mullin, V. T. Sós, D. Stinson: Finite linear spaces and projective planes, *Discrete Math.* 47 (1983) no. 1, 49–62.
- [12] R. A. Fisher, An examination of the different possible solutions of a problem in incomplete blocks, *Annals of Eugenics* 10 (1940), 52–75.
- [13] J. Fowler, A short proof of Totten's classification of restricted linear spaces, *Geometriae Dedicata*, 15 (1984) no. 4, 413–422.
- [14] Z. Füredi, New asymptotics for bipartite Turán numbers. *J. Combin. Theory Ser. A* 75 (1996) no. 1, 141–144.
- [15] R.L. Graham and H.O. Pollak, On the addressing problem for loop switching, *Bell System Tech. J.* 50 (1971), 2495–2519.
- [16] G. H. Hardy, J. E. Littlewood and G. Polya, *Inequalities*, 2nd edition, Cambridge University Press (1988), Cambridge, England, pp. 43–44.
- [17] A. Hartman, R. C. Mullin, D. Stinson, Exact covering configurations and Steiner Systems, *J. London. Math. Soc* (2) 25, (1982) 193–200.
- [18] J. Hirschfeld, *Projective geometries over finite fields*, Second edition. Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York (1998).
- [19] J. Hirschfeld, *Finite projective spaces of three dimensions*, Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, New York (1985).

- [20] T. Kővári, V. T. Sós and P. Turán, On a problem of K. Zarankiewicz, *Colloquium Math.*, 3, (1954), 50-57.
- [21] C. Lam, C. The search for a finite projective plane of order 10. *Amer. Math. Monthly* 98 (1991), no. 4, 305–318.
- [22] D. K. Ray-Chaudhuri and R. M. Wilson, On t -designs, *Osaka J. Math.* 12 (1975), 737–744.
- [23] B. Segre, Ovals in a finite projective plane, *Canad. J. Math.* 7 (1955) 414–416.
- [24] J. Thas, The affine plane $AG(2,q)$, q odd, has a unique one point extension, *Invent. math.* 118 (1994) 133–139.
- [25] J. Tits, Ovoïdes et groupes de Suzuki, *Arch. Math.* 13 (1962) 187–198.
- [26] J. Totten, Classification of restricted linear spaces, *Canad. J. Math.* 28 (1976), 321–333.
- [27] S. Vishwanathan, A counting proof of the Graham-Pollak Theorem, preprint
- [28] R. van Lint, R. Wilson, *Designs, Graphs, Codes, and Their Links*, Cambridge University Press (1991).
- [29] J.B. Wilker, Inversive geometry. *The geometric vein*, Springer, New York-Berlin (1981) 379–442.
- [30] E. Witt, Über Steinersche Systeme, *Abh. Math. Sem. Univ. Hamburg* 12 (1938) 265–275.