

# Asymptotically optimal induced universal graphs

Noga Alon <sup>\*†</sup>

## Abstract

We prove that the minimum number of vertices of a graph that contains every graph on  $k$  vertices as an induced subgraph is  $(1 + o(1))2^{(k-1)/2}$ . This improves earlier estimates of Moon, of Bollobás and Thomason, of Brightwell and Kohayakawa and of Alstrup, Kaplan, Thorup and Zwick. The method supplies similarly sharp estimates for the analogous problems for directed graphs, tournaments, bipartite graphs, oriented graphs and more. We also show that if  $\binom{n}{k}2^{-\binom{k}{2}} = \lambda$  then the probability that the random graph  $G(n, 0.5)$  contains every graph on  $k$  vertices as an induced subgraph is  $(1 - e^{-\lambda})^2 + o(1)$ .

The proofs combine combinatorial and probabilistic arguments with tools from group theory.

## 1 Introduction

Let  $\mathcal{F}$  be a finite family of graphs. A graph  $G$  is *induced universal* for  $\mathcal{F}$  if every member  $F$  of  $\mathcal{F}$  is an induced subgraph of  $G$ . The definition naturally extends to digraphs, tournaments or oriented graphs. This notion was introduced by Rado [28]. There is a vast literature about the question of determining or estimating the minimum possible number of vertices of an induced universal graph for given families of  $k$ -vertex graphs or digraphs. See [25], [26], [11], [15], [20], [23], [19], [17], [14], [5], [4] and the references therein. In some of these papers the minimum investigated is determined up to a constant factor, but there is no known nontrivial example in which the correct constant is known.

The most basic question, and the one with the longest history deals with the family of all undirected graphs on a given number of vertices. Let  $\mathcal{F}(k)$  denote the family of all  $k$ -vertex undirected graphs, and let  $f(k)$  denote the smallest possible number of vertices of

---

<sup>\*</sup>Sackler School of Mathematics and Blavatnik School of Computer Science, Tel Aviv University, Tel Aviv 69978, Israel. [nogaa@tau.ac.il](mailto:nogaa@tau.ac.il). Research supported in part by a USA-Israeli BSF grant 2012/107, by an ISF grant 620/13 and by the Israeli I-Core program.

<sup>†</sup>*Mathematics Subject Classification:* 05C35, 05C80

an induced universal graph for  $\mathcal{F}(k)$ . Moon [25] observed that a simple counting argument gives  $f(k) \geq 2^{(k-1)/2}$  and proved that  $f(k) \leq O(k2^{k/2})$ . Alstrup, Kaplan, Thorup and Zwick [4] determined  $f(k)$  up to a constant factor, showing that  $f(k) \leq 16 \cdot 2^{k/2}$ . Bollobás and Thomason [11] proved that the random graph  $G(n, 0.5)$  on  $n = k^2 2^{k/2}$  vertices is induced universal for  $\mathcal{F}(k)$  with high probability, that is, with probability that tends to 1 as  $k$  tends to infinity. This was improved to  $n = O(k2^{k/2})$  by Brightwell and Kohayakawa in [13]. The question of finding tighter bounds for  $f(k)$ , suggested by the work of Moon, is mentioned by Vizing in [29] and by Alstrup et. al (whose work determines it up to a constant factor of  $16\sqrt{2}$ ) in [4]. Here we show that the lower bound is tight, up to a lower order additive term.

**Theorem 1.1**

$$f(k) = (1 + o(1))2^{(k-1)/2}.$$

The proof combines probabilistic and combinatorial arguments with some group theoretical facts about graphs with large automorphism groups. Similar arguments supply asymptotically tight estimates for the analogous questions for directed graphs, oriented graphs, tournaments, bipartite graphs or complete graphs with colored edges, improving results in [26], [23], [4]. Since the proofs in all cases, besides possibly that of bipartite graphs, are similar, we focus here on the undirected case and merely include the statements and sketches of proofs of these variants, with more details about the family of bipartite graphs.

As a byproduct of (a variant of) the first part of the proof we show that the minimum number of vertices  $n$  so that the random graph on  $n$  vertices is induced universal for  $\mathcal{F}(k)$  with high probability is  $(1 + o(1))\frac{k}{e}2^{(k-1)/2}$ , improving the estimate in [11]. This was harder to improve in 1981, when [11] was written, but is simpler now, using some of the more recently developed high deviation inequalities. The bound above also improves the estimate in [13] by a constant factor. Combining our argument with some group theoretic tools and the Stein-Chen method we prove a more precise statement, as follows.

**Theorem 1.2** *Let  $n > k > 1$ , let  $G = G(n, 0.5)$  be the binomial random graph, and put*

$$\lambda = \binom{n}{k} 2^{-\binom{k}{2}}.$$

*Then the probability that  $G$  is induced universal for  $\mathcal{F}(k)$  is  $(1 - e^{-\lambda})^2 + o(1)$ , where the  $o(1)$  term tends to 0 (uniformly in  $k = k(n)$ ) as  $n$  tends to infinity.*

The rest of this paper is organized as follows. Theorem 1.1 is proved in Section 2 and Section 3 deals with the minimum possible  $n$  so that the random graph  $G(n, 0.5)$  is induced universal for  $\mathcal{F}(k)$  with high probability. The proof of Theorem 1.2 appears in the second part of this section. In Section 4 we describe several variants of Theorem 1.1 that can be proved using a similar approach, and the final Section 5 contains some concluding remarks and open problems.

In order to simplify the presentation we omit all floor and ceiling signs whenever these are not essential, and assume, whenever needed, that  $k$  is sufficiently large. Throughout the proof we make no attempt to optimize the absolute constants whenever these are not crucial. All logarithms are in base 2, unless otherwise specified.

## 2 The main proof

In this section we prove Theorem 1.1. It is convenient to split the proof into two main parts.

### 2.1 Asymmetric graphs

Call a graph on  $k$  vertices *asymmetric* if every induced subgraph of it has at most  $k^{4m}$  automorphisms, where  $m = 2\sqrt{k \log k}$ . Note that, in particular, the number of automorphisms of any such graph is at most  $k^{4m}$ . Let  $\mathcal{H}(k)$  denote the family of all asymmetric graphs on  $k$  vertices.

In this subsection we prove that there are small induced universal graphs for  $\mathcal{H}(k)$ : in fact, the random graph with the appropriate number of vertices is, with high probability, such a universal graph.

Let  $n$  be the smallest integer that satisfies the following inequality

$$\binom{n}{k} \frac{k!}{k^{8m}} 2^{-\binom{k}{2}} \geq 1. \quad (1)$$

Since the ratio between  $\binom{n+1}{k}$  and  $\binom{n}{k}$  is  $\frac{n+1}{n-k+1}$  which is very close to 1 for the relevant parameters, the left-hand-side of (1) for this smallest  $n$  is  $1 + o(1)$  and thus one can solve for  $n$  and see that it satisfies

$$n = 2^{(k-1)/2} \left(1 + O\left(\frac{\log^{3/2} k}{\sqrt{k}}\right)\right).$$

In particular,  $n = (1 + o(1))2^{(k-1)/2}$ .

**Theorem 2.1** *Let  $n$  be as above and let  $G = (V, E) = G(n, 0.5)$  be the random graph on a set  $V$  of  $n$  vertices obtained by picking, randomly and independently, each pair of vertices to be an edge with probability  $1/2$ . Then, with high probability (that is, with probability that tends to 1 as  $k$ , and hence  $n$ , tend to infinity),  $G$  is an induced universal graph for  $\mathcal{H}(k)$ .*

In the proof of the above theorem we apply (a known consequence of) Talagrand's Inequality, described, for example, in [3], Theorem 7.7.1. The statement follows.

**Theorem 2.2 (Talagrand's Inequality)** *Let  $\Omega = \prod_{i=1}^p \Omega_i$ , where each  $\Omega_i$  is a probability space and  $\Omega$  has the product measure, and let  $h : \Omega \rightarrow \mathbb{R}$  be a function. Assume that  $h$  is Lipschitz, that is,  $|h(x) - h(y)| \leq 1$  whenever  $x, y$  differ in at most one coordinate. For a function  $f : \mathbb{N} \rightarrow \mathbb{N}$ ,  $h$  is  $f$ -certifiable if whenever  $h(x) \geq s$  there exists  $I \subseteq \{1, \dots, p\}$  with  $|I| \leq f(s)$  so that for every  $y \in \Omega$  that agrees with  $x$  on the coordinates  $I$  we have  $h(y) \geq s$ . Suppose that  $h$  is  $f$ -certifiable and let  $Y$  be the random variable given by  $Y(x) = h(x)$  for  $x \in \Omega$ . Then for every  $b$  and  $t$*

$$\text{Prob}[Y \leq b - t\sqrt{f(b)}] \cdot \text{Prob}[Y \geq b] \leq e^{-t^2/4}.$$

We also need the following simple lemma.

**Lemma 2.3** *Let  $H \in \mathcal{H}(k)$ . Let  $K$  and  $K'$  be two sets of labelled vertices, each of size  $k$ , where  $|K \cap K'| = k - i$ . Then the number of ways to choose the edges and nonedges among all  $2\binom{k}{2} - \binom{k}{k-i}$  pairs that lie in  $K$  or in  $K'$  so that the induced subgraph on  $K$  is isomorphic to  $H$  and the induced subgraph on  $K'$  is also isomorphic to  $H$  is at most*

$$\frac{k!}{|\text{Aut}(H)|} k^i k^{4m}.$$

**Proof:** There are exactly  $\frac{k!}{|\text{Aut}(H)|}$  copies of  $H$  on  $K$ . For each such fixed copy, we bound the number of embeddings of  $H$  in  $K'$ . There are at most  $k(k-1) \cdots (k-i+1) < k^i$  ways to choose the vertices of  $H$  mapped to the vertices of  $K' - K$ . Fix a set  $T$  of these  $i$  vertices and their embedding. In order to complete the embedding, the induced subgraph of the copy of  $H$  placed in  $K$  on the set of vertices  $K \cap K'$  has to be isomorphic to the induced subgraph of  $H$  on  $V(H) - T$ . If so, then the number of ways to embed these  $k-i$  vertices is the number of automorphisms of this induced subgraph of  $H$ , which is, by the definition of  $\mathcal{H}$ , at most  $k^{4m}$ .  $\square$

**Proof of Theorem 2.1:** Let  $H$  be a fixed member of  $\mathcal{H}(k)$  and let  $s = |\text{Aut}(H)|$  be the size of its automorphism group. Then

$$s \leq k^{4m} = k^{8\sqrt{k \log k}}.$$

Let  $G = (V, E)$  be the random graph  $G(n, 0.5)$ , where  $n$  is as chosen in (1). For every subset  $K \subset V$  of size  $|K| = k$  let  $X_K$  denote the indicator random variable whose value is 1 if the induced subgraph of  $G$  on  $K$  is isomorphic to  $H$  and let  $X = \sum_K X_K$ , where the summation is over all subsets  $K \subset V$  of cardinality  $k$ . Thus  $X$  is the number of copies of  $H$  in  $G$ . The expectation of each  $X_K$  is clearly

$$E(X_K) = \frac{k!}{s} 2^{-\binom{k}{2}}.$$

Thus, by linearity of expectation,

$$E(X) = \binom{n}{k} \frac{k!}{s} 2^{-\binom{k}{2}} \geq k^{4m}$$

where in the last inequality we used (1) and the fact that  $s \leq k^{4m}$ . Note also that since

$$\binom{n}{k} \frac{k!}{k^{8m}} 2^{-\binom{k}{2}} = 1 + o(1)$$

it follows that the expectation of  $X$  is at most  $(1 + o(1))k^{8m} < n^{0.01}$  (even if  $s = 1$ ).

We say that two copies of  $H$  in  $G$  have a *nontrivial intersection* if they share at least two vertices. Put  $\mu = E(X)$ . Let  $Z$  denote the random variable  $Z = \sum_{K, K'} X_K X_{K'}$  where the summation is over all (ordered) pairs of  $k$ -subsets  $K, K'$  of  $V$  that satisfy  $2 \leq |K \cap K'| \leq k - 1$ . Thus,  $Z$  is the number of pairs of copies of  $H$  in  $G$  that have a nontrivial intersection. We next compute the expectation of  $Z$  and show that it is much smaller than  $\mu$  (and hence also much smaller than  $\mu^2$ ). Put  $\Delta = E(Z)$  and note that  $\Delta = \sum_{j=2}^{k-1} \Delta_j$  where  $\Delta_j$  is the expected number of pairs  $K, K'$  with  $X_K = X_{K'} = 1$  and  $|K \cap K'| = j$ .

**Claim:** For each  $2 \leq j \leq k - 1$

$$\Delta_j \leq \mu \frac{1}{n^{0.48}} \tag{2}$$

**Proof of Claim:** Consider two possible cases, as follows.

**Case 1:**  $2 \leq j \leq 3k/4$ .

In this case

$$\Delta_j \leq \binom{n}{k} \frac{k!}{s} 2^{-\binom{k}{2}} \binom{k}{j} \binom{n-k}{k-j} \frac{k!}{s} 2^{-\binom{k}{2} + \binom{j}{2}}.$$

Indeed, there are  $\binom{n}{k}$  ways to choose the set  $K$ , and  $\binom{k}{j} \binom{n-k}{k-j}$  ways to choose  $K'$  with  $|K \cap K'| = j$ . There are  $\frac{k!}{s}$  ways to place a copy of  $H$  in  $K$  and  $\frac{k!}{s}$  ways to place a copy of  $H$  in  $K'$  (this is an overcount, as these two copies have to agree on the edges in their common part). This determines all the edges and nonedges in the induced graph on  $K$  and on  $K'$ , and the probability that  $G$  indeed has exactly these edges is

$$2^{-\binom{k}{2}} \cdot 2^{-\binom{k}{2} + \binom{j}{2}}$$

Therefore,

$$\frac{\Delta_j}{\mu^2} \leq \frac{\binom{k}{j} \binom{n-k}{k-j} 2^{\binom{j}{2}}}{\binom{n}{k}} \leq \left( \frac{k^2 2^{(j-1)/2}}{n} \right)^j \leq \left( \frac{1}{n^{1/4-0.005}} \right)^j \leq \frac{1}{n^{0.49}}.$$

Here we used the fact that  $k = (2 + o(1)) \log_2 n$  and that since  $j \leq 3k/4$  it follows that  $2^{(j-1)/2} \leq n^{3/4+o(1)}$ . Recall that  $\mu \leq n^{0.01}$  and thus

$$\frac{\Delta_j}{\mu} \leq \mu \frac{\Delta_j}{\mu^2} \leq \frac{1}{n^{0.48}}$$

as claimed.

**Case 2:**  $j = k - i$ ,  $i \leq k/4$ .

In this case we have, by Lemma 2.3

$$\Delta_j \leq \binom{n}{k} \binom{k}{j} \binom{n-k}{k-j} \frac{k!}{s} k^i k^{4m} 2^{-2\binom{k}{2} + \binom{j}{2}}.$$

Indeed, there are

$$\binom{n}{k} \binom{k}{j} \binom{n-k}{k-j}$$

ways to choose the sets  $K, K'$  having intersection  $j = k - i$ , and by Lemma 2.3 for each such choice there are at most  $\frac{k!}{s} k^i k^{4m}$  ways to place copies of  $H$  in each of them. The probability that this coincides with all edges and nonedges of the induced subgraph of  $G$  on  $K$  and on  $K'$  is  $2^{-2\binom{k}{2} + \binom{j}{2}}$ .

Since  $\mu \geq k^{4m} > 1$  this implies that

$$\frac{\Delta_j}{\mu^2} < \frac{\Delta_j}{\mu} \leq \binom{k}{j} \binom{n-k}{k-j} k^i k^{4m} 2^{-\binom{k}{2} + \binom{j}{2}}$$

$$\leq \binom{k}{i} \binom{n-k}{i} k^i k^{4m} 2^{-i(k-i)} \leq (k^2 n 2^{-(k-i)})^i k^{4m} \leq \frac{1}{n^{0.5-o(1)}} \leq \frac{1}{n^{0.48}}.$$

This completes the proof of the claim.

Returning to the proof of the theorem, note that the variance of the random variable  $X$  satisfies

$$\text{Var}(X) \leq E(X) + \sum_{K, K'} \text{Cov}(X_K, X_{K'})$$

where the summation is over all ordered pairs  $K, K'$  where  $2 \leq |K \cap K'| \leq k-1$  (since for all other pairs the covariance is zero). Since

$$\text{Cov}(X_K, X_{K'}) = E(X_K X_{K'}) - E(X_K)E(X_{K'}) \leq E(X_K X_{K'})$$

it follows from the claim above that

$$\text{Var}(X) \leq \mu + \Delta \leq (1 + o(1))\mu.$$

Therefore, by Chebyshev's Inequality (and the fact that  $\mu$  is large), the probability that  $X \geq 3\mu/4$  is (much) bigger than  $3/4$ . By Markov's inequality, the probability that  $\Delta \leq \mu/4$  is also (much) bigger than  $3/4$  and hence with probability larger than  $1/2$ , both events happen simultaneously, that is, the number of copies of  $H$  in  $G$  is at least  $3\mu/4$  and the number of pairs of copies of  $H$  with nontrivial intersection is smaller than  $\mu/4$ . By removing one copy of  $H$  from each pair with a nontrivial intersection we conclude that if this is the case, then  $G$  contains a family of at least  $\mu/2$  copies of  $H$  with no two having a nontrivial intersection.

Let  $h(G)$  be the maximum cardinality of a family of copies of  $H$  in  $G$  in which no two members have a nontrivial intersection, and let  $Y$  be the random variable  $Y = h(G)$ . Our objective is to apply Talagrand's Inequality (Theorem 2.2) to deduce that the probability that  $Y$  is zero is tiny.

By the above discussion, the probability that  $Y$  is at least  $\mu/2$  exceeds  $1/2$ . It is also clear that the value of  $Y = h(G)$  can change by at most 1 if we add or delete one edge to  $G$ , and that  $h$  is  $f$ -certifiable where  $f(s) = s \binom{k}{2}$ . Therefore, by Theorem 2.2 with  $b = \mu/2$  and  $t = \sqrt{\mu}/k$  we conclude that the probability that  $Y = 0$  is smaller than

$$e^{-\mu/4k^2}$$

which is much much smaller than  $2^{-k^2}$ . As  $Y = 0$  if and only if there is no copy of  $H$  in  $G$ , and as the total number of graphs in  $\mathcal{H}(k)$  is smaller than  $2^{\binom{k}{2}}$ , we conclude that  $G$  is induced universal for  $\mathcal{H}(k)$  with high probability. This completes the proof of Theorem 2.1.  $\square$

## 2.2 Symmetric graphs

Recall that we called a graph on  $k$  vertices asymmetric if no induced subgraph of it has more than  $k^{4m}$  automorphisms, where  $m = 2\sqrt{k \log k}$ . Call a graph on  $k$  vertices *symmetric* if it is not asymmetric. Let  $\mathbf{T}(k) = \mathcal{F}(k) - \mathcal{H}(k)$  denote the set of all symmetric graphs on  $k$  vertices

In this subsection we construct a small induced universal graph for the family  $\mathbf{T}(k)$ . To this end it is desirable to obtain some useful structural properties of graphs with a large automorphism group by utilizing known results about large subgroups of the symmetric group. There is a rich literature about automorphism groups of graphs, see, for example [7] and its many references. It seems, however, hopeless to try to characterize all graphs with at most  $k$  vertices and at least  $k^{8\sqrt{k \log k}}$  automorphisms. Fortunately, for our purpose here it suffices to prove and apply some partial information about their structure, as described in what follows.

The *minimal degree* of a permutation group is the size of the minimum support of a nontrivial element in it. There are several results stating that large permutation groups have nontrivial elements with small supports (or equivalently, many fixed points), see, for example, [21] and the references therein. For our purpose here the following simple fact suffices.

**Lemma 2.4** *For any  $p > 1$  and  $t$ , any subgroup  $S$  of size at least  $p^{4t}$  of the symmetric group  $S_p$  contains a permutation with at least  $t$  and at most  $p - 3t$  fixed points.*

**Proof:** Consider the subgroup as a group of permutations of  $[p] = \{1, 2, \dots, p\}$ . By the pigeonhole principle there is a subset  $A = \{a_1, a_2, \dots, a_t\}$  of  $t$  elements of  $[p]$  so that there are at least  $\frac{|S|}{p(p-1)\dots(p-t+1)} > p^{3t}$  permutations  $\sigma$  in  $S$  satisfying  $\sigma(i) = a_i$  for all  $i \in [t] = \{1, 2, \dots, t\}$ . For any two such permutations  $\sigma_1, \sigma_2$ , the product  $\sigma_1\sigma_2^{-1}$  fixes all points of  $A$ . Let  $S'$  be the subgroup of  $S$  that fixes all points of  $A$ . Then  $|S'| > p^{3t}$ . The number of permutations in  $S'$  that fixes all points but at most  $i$  is clearly at most  $\binom{p-t}{i} i! < p^i$ , and since  $p^{3t-1} < p^{3t}$  there is an element of  $S'$  that fixes at most  $p - 3t$  points.  $\square$

**Corollary 2.5** *Let  $T = (V, E)$  be a graph in  $\mathbf{T}(k)$ . Then there are three pairwise disjoint sets of vertices  $A, B, C$  of  $T$ , each of cardinality  $m$ , so that the following holds. There is a numbering of the elements of  $A, B, C$ :  $A = \{a_1, a_2, \dots, a_m\}$ ,  $B = \{b_1, b_2, \dots, b_m\}$  and  $C = \{c_1, c_2, \dots, c_m\}$  such that for any  $1 \leq i, j \leq m$ ,  $a_i b_j$  is an edge of  $T$  if and only if  $a_i c_j$  is an edge of  $T$ . That is, for each  $1 \leq j \leq m$ ,  $b_j$  and  $c_j$  have exactly the same neighbors in the set  $A$ .*



**Proof:** By the definition of  $\mathbf{T}(k)$  there is an induced subgraph  $T'$  of  $T$  on  $p \leq k$  vertices whose group of automorphisms  $S$  is of size at least  $k^{4m} \geq p^{4m}$ . By Lemma 2.4 this group contains a permutation  $\sigma$  with at least  $m$  and at most  $p - 3m$  fixed points. Let  $A = \{a_1, a_2, \dots, a_m\}$  be  $m$  of these fixed points and consider the expression of  $\sigma$  as a product of (nontrivial) cycles. The total length of these cycles is at least  $3m$ . From each cycle  $(w_1, w_2, \dots, w_r)$  of length  $r$ , define  $\lfloor r/2 \rfloor$  disjoint pairs

$$(w_1, w_2), (w_3, w_4), \dots, (w_{2\lfloor r/2 \rfloor - 1}, w_{2\lfloor r/2 \rfloor}).$$

Altogether we get at least  $m$  such pairs, let  $(b_i, c_i)$ ,  $(1 \leq i \leq m)$  be  $m$  of them. Observe, now, that for every  $1 \leq j \leq m$ ,  $\sigma$  maps  $b_j$  to  $c_j$  and fixes all elements  $a_i$ . As  $\sigma$  is an automorphism of  $T$  this means that for every  $i$ ,  $a_i b_j$  is an edge of  $T$  if and only if so is  $a_i c_j$ .  $\square$

**Lemma 2.6** *Let  $T'$  be the graph obtained from a complete graph on  $k$  vertices by removing all edges of a complete bipartite graph  $K_{m,m}$  in it, where, as before,  $m = 2\sqrt{k \log k}$ . Let  $[k] = \{1, 2, \dots, k\}$  be the set of vertices of  $T'$  and suppose  $A' = \{1, 2, \dots, m\}$ ,  $B' = \{m+1, \dots, 2m\}$ ,  $C' = \{2m+1, 2m+2, \dots, 3m\}$  and  $D' = \{3m+1, 3m+2, \dots, k\}$ , where there are no edges between  $A'$  and  $C'$  and all other pairs of vertices of  $T'$  are adjacent. Then there is an orientation  $T''$  of the edges of  $T'$  in which all edges between  $A'$  and  $B'$  are oriented from  $A'$  to  $B'$  and the maximum outdegree of a vertex in  $T''$  is at most  $\frac{k}{2} - \frac{m^2}{k} + O(1) < \frac{k}{2} - 2 \log k$ .*

**Proof:** We need two simple facts, as follows.

**Fact 1:** For any integer  $g > 1$  the edges of the complete graph  $K_g$  on  $g$  vertices can be oriented so that every outdegree is at most  $g/2$ .

Indeed, if  $g$  is odd then  $K_g$  has an Eulerian orientation in which every outdegree is exactly  $(g-1)/2$ , and if  $g$  is even, omit a vertex from an Eulerian orientation of  $K_{g+1}$  to get an orientation as needed.

**Fact 2:** For any positive integers  $p, q$  and  $r \leq q$  there is a bipartite graph with classes of vertices  $P$  and  $Q$  of sizes  $p$  and  $q$ , respectively, so that every vertex of  $P$  has degree exactly  $r$  and every vertex of  $Q$  has degree either  $\lfloor pr/q \rfloor$  or  $\lceil pr/q \rceil$ .

One simple way to prove this fact is to number the vertices of  $Q$ :  $u_1, u_2, \dots, u_q$  and to connect, for each  $i$ , vertex number  $i$  of  $P$  to the vertices

$$u_{(i-1)r+1}, u_{(i-1)r+2}, \dots, u_{ir}$$

where the indices are reduced modulo  $q$ .

Using these two facts construct an orientation  $T''$  of  $T'$  as follows. Orient the edges of the complete graph on  $A'$ , using Fact 1, so that each outdegree is at most  $m/2$ . Similarly, orient the edges of the complete graph on  $B' \cup C'$  so that every outdegree is at most  $m$ , and orient the edges of the complete graph on  $D'$  so that every outdegree is at most  $(k - 3m)/2$ . All edges between  $A'$  and  $B'$  are oriented from  $A'$  to  $B'$ . By Fact 2, for any real  $x \in (0, 1)$  the edges of the complete bipartite graph with vertex classes  $A'$  and  $D'$  can be oriented so that the outdegree of each vertex of  $A'$  is at most  $x|D'| + 1 = x(k - 3m) + 1$  and the outdegree of each vertex of  $D'$  is at most  $(1 - x)m + O(1)$ . Similarly, for any real  $y \in (0, 1)$  the edges of the complete bipartite graph with vertex classes  $B' \cup C'$  and  $D'$  can be oriented so that the outdegree of each vertex of  $B' \cup C'$  is at most  $y(k - 3m) + 1$  and the outdegree of each vertex of  $D'$  is at most  $(1 - y)2m + O(1)$ . In the resulting orientation the outdegrees of the vertices of  $A'$ ,  $B' \cup C'$  and  $D'$  are bounded, up to absolutely bounded additive terms, by

$$\frac{3}{2}m + x(k - 3m), \quad m + y(k - 3m) \quad \text{and} \quad \frac{k - 3m}{2} + (1 - x)m + (1 - y)2m,$$

respectively. Since  $m = o(k)$  it is not difficult to check that there are  $x, y \in (0, 1)$  (which are both  $1/2 + o(1)$ ) so that these 3 quantities are equal. Although the precise expressions for  $x$  and  $y$  are not needed, we note here that these are:

$$x = \frac{1}{2} - \frac{m^2}{k(k - 3m)}, \quad y = \frac{1}{2} - \frac{m^2}{k(k - 3m)} + \frac{m}{2(k - 3m)}.$$

With these  $x$  and  $y$  all three quantities above are exactly  $\frac{k}{2} - \frac{m^2}{k}$ .

It follows that there is an orientation  $T''$  in which all outdegrees are equal, up to an  $O(1)$  additive error, and as the total number of edges is  $\binom{k}{2} - m^2$  this (as well as the explicit computation above) implies that every outdegree in  $T''$  is at most  $\frac{k}{2} - \frac{m^2}{k} + O(1) < \frac{k}{2} - 2 \log k$ , completing the proof of the lemma.  $\square$

**Theorem 2.7** *There is an induced universal graph for  $\mathbf{T}(k)$  whose number of vertices is at most  $2^{k/2 - \log k}$ .*

**Proof:** An *adjacency labeling scheme* for a family of graphs is a way of assigning labels to the vertices of each graph in the family such that given the labels of two vertices in the graph it is possible to determine whether or not they are adjacent in the graph, without using any additional information besides the labels. It is easy and well known that a family of graphs has a labeling scheme in which every label contains  $L$  bits if and only if there is an induced universal graph for the family with at most  $2^L$  bits. This

is implicit in the work of Moon [25] and is mentioned explicitly in lots of subsequent papers starting with [20]. Indeed, for a given labeling scheme for a family, the graph whose vertices are all possible labels in which two vertices are adjacent if and only if their labels correspond to adjacent vertices is an induced universal graph for the family (and the converse is equally simple). It thus suffices to describe a labeling scheme for the family  $\mathbf{T}(k)$  in which each label consists of at most  $k/2 - \log k$  bits. Given a graph  $T \in \mathbf{T}(k)$  we describe the labels of its vertices. By Corollary 2.5  $T$  contains three disjoint subsets of vertices  $A = \{a_1, a_2, \dots, a_m\}$ ,  $B = \{b_1, b_2, \dots, b_m\}$  and  $C = \{c_1, c_2, \dots, c_m\}$  satisfying the assertion of the lemma. Number the vertices of  $T$  by the integers  $1, 2, \dots, k$  so that  $a_i$  gets the number  $i$ ,  $b_i$  the number  $m + i$  and  $c_i$  the number  $2m + i$ , where the rest of the numbering is arbitrary. Let  $T''$  be the graph constructed in Lemma 2.6. The label of vertex number  $i$  of  $T$  is the number  $i$  followed by one bit for each outneighbor  $j$  of  $i$  in  $T''$ , in order. This bit is 0 if  $i$  and  $j$  are not adjacent in  $T$ , and is 1 if they are adjacent. Note that the length of each label is at most  $\log k + k/2 - 2 \log k \leq k/2 - \log k$ , where the first  $\log k$  bits are used to present the number of the vertex.

It is not difficult to check that this is a valid labeling scheme. Given the labels of two vertices, if it is not the case that one of them lies in  $A$  and the other in  $C$ , one of the labels contains the information about the adjacency between the two vertices, and the labels (together with the graph  $T''$  which is known as part of the labeling scheme) determine this information. The only exceptional case is when one of the two vertices is  $a_i$  and the other is  $c_j$ . But in this case the label of  $a_i$  determines whether or not  $a_i$  is adjacent to  $b_j$ , and this determines the information about the adjacency between  $a_i$  and  $c_j$  as well, by Corollary 2.5. This completes the proof of the theorem.  $\square$

### 2.3 The proof of Theorem 1.1

The assertion of Theorem 1.1 clearly follows from that of Theorem 2.1 together with that of Theorem 2.7. The required induced universal graph is simply the vertex disjoint union of the graph in Theorem 2.1 and the one in Theorem 2.7. Note that the size of the second graph is negligible compared to that of the first one, and thus the proof gives that the minimum possible number of vertices of an induced universal graph for  $\mathcal{F}(k)$ , namely, the function  $f(k)$ , satisfies

$$f(k) \leq 2^{(k-1)/2} \left( 1 + O\left(\frac{\log^{3/2} k}{\sqrt{k}}\right) \right).$$

$\square$

### 3 Random graphs

In this section we establish a sharp estimate for the minimum integer  $n = n(k)$  so that the random graph  $G(n, 0.5)$  is induced universal for  $\mathcal{F}(k)$  with high probability. We first establish this estimate in subsection 3.1, and then prove a stronger hitting time result in subsection 3.2. The basic argument in the first subsection is similar to the one used in the proof of Theorem 2.1, with one additional simple trick that simplifies the computation. The idea is to insist on getting each labelled  $k$  vertex graph  $F \in \mathcal{F}(k)$  as an induced subgraph of the labelled random graph  $G = G(n, 0.5)$  on the set of vertices  $[n]$  by an order preserving mapping, namely, the embedding of  $H$  in  $G$  respects the order of the labels. Note that for such embeddings, for every fixed labelled  $F$  and every set  $K$  of  $k$  vertices of  $G$ , the probability that the labelled induced subgraph of  $G$  on  $K$  is (monotonely) isomorphic to  $F$  is exactly  $2^{-\binom{k}{2}}$ , independently of the number of automorphisms of  $F$ .

#### 3.1 A tight estimate

Let  $n$  be the smallest integer that satisfies

$$\binom{n}{k} 2^{-\binom{k}{2}} \geq 4k^4. \quad (3)$$

As in subsection 2.1, since the ratio between  $\binom{n+1}{k}$  and  $\binom{n}{k}$  is  $\frac{n+1}{n-k+1}$  which is very close to 1 for the relevant parameters, the left-hand-side of (3) for this smallest  $n$  is  $1 + o(1)$  and thus, by Stirling's Formula

$$n = \frac{k}{e} 2^{(k-1)/2} (1 + O(\frac{\log k}{k})),$$

and in particular,  $n = (1 + o(1)) \frac{k}{e} 2^{(k-1)/2}$ .

**Theorem 3.1** *Let  $n$  be as above and let  $G = (V, E) = G(n, 0.5)$  be the random graph on a set  $V = \{1, 2, \dots, n\}$  of  $n$  labelled vertices. Then, with high probability every labelled  $F \in \mathcal{F}(k)$  is an induced subgraph of  $G$  with an order preserving embedding. In particular,  $G$  is an induced universal graph for  $\mathcal{F}(k)$  with high probability.*

**Proof:** Let  $F$  be a fixed member of  $\mathcal{F}(k)$  considered as a labelled graph on  $[k] = \{1, 2, \dots, k\}$ , and let  $G$  be the random graph  $G(n, 0.5)$  on  $[n] = \{1, 2, \dots, n\}$ , where  $n$  is as chosen in (3). For every subset  $K \subset V$  of size  $|K| = k$  let  $X_K$  denote the indicator random variable whose value is 1 if the induced subgraph of  $G$  on  $K$  is isomorphic to  $F$

with a monotone embedding and let  $X = \sum_K X_K$ , where the summation is over all subsets  $K \subset V$  of cardinality  $k$ . Thus  $X$  is the number of copies of  $F$  in  $G$ . The expectation of each  $X_K$  is clearly

$$E(X_K) = 2^{-\binom{k}{2}},$$

and thus, by linearity of expectation,

$$E(X) = \binom{n}{k} 2^{-\binom{k}{2}} \geq 4k^4$$

where the last inequality follows from (3).

As before, say that two copies of  $F$  in  $G$  have a nontrivial intersection if they share at least two vertices. Put  $\mu = E(X)$ , then  $\mu = (1 + o(1))4k^4$ . Let  $Z$  denote the random variable  $Z = \sum_{K, K'} X_K X_{K'}$  where the summation is over all (ordered) pairs of  $k$ -subsets  $K, K'$  of  $V$  that satisfy  $2 \leq |K \cap K'| \leq k - 1$ . Thus,  $Z$  is the number of pairs of copies of  $F$  in  $G$  that have a nontrivial intersection. We next compute the expectation of  $Z$  and show that it is much smaller than  $\mu$  (and hence also much smaller than  $\mu^2$ ). Put  $\Delta = E(Z)$  and note that  $\Delta = \sum_{j=2}^{k-1} \Delta_j$  where  $\Delta_j$  is the expected number of pairs  $K, K'$  with  $X_K = X_{K'} = 1$  and  $|K \cap K'| = j$ .

**Claim:** For each  $2 \leq j \leq k - 1$

$$\Delta_j \leq \mu \frac{1}{n^{0.48}} \tag{4}$$

**Proof of Claim:** Consider two possible cases, as follows.

**Case 1:**  $2 \leq j \leq 3k/4$ .

In this case

$$\Delta_j \leq \binom{n}{k} 2^{-\binom{k}{2}} \binom{k}{j} \binom{n-k}{k-j} 2^{-\binom{k}{2} + \binom{j}{2}}.$$

Indeed, there are  $\binom{n}{k}$  ways to choose the set  $K$ , and  $\binom{k}{j} \binom{n-k}{k-j}$  ways to choose  $K'$  with  $|K \cap K'| = j$ . There is a unique way to place a copy of  $F$  monotonely in  $K$  and a unique way to place a copy of  $F$  in  $K'$ . This is an overcount, as these two copies have to agree on the edges in their common part. This determines all the edges in the induced graph on  $K$  and on  $K'$ , and the probability that  $G$  indeed has exactly these edges and nonedges is

$$2^{-\binom{k}{2}} \cdot 2^{-\binom{k}{2} + \binom{j}{2}}$$

Therefore,

$$\frac{\Delta_j}{\mu^2} \leq \frac{\binom{k}{j} \binom{n-k}{k-j} 2^{\binom{j}{2}}}{\binom{n}{k}} \leq \left( \frac{k^2 2^{(j-1)/2}}{n} \right)^j \leq \left( \frac{1}{n^{1/4-0.005}} \right)^j \leq \frac{1}{n^{0.49}}.$$

Here we used the fact that  $k = (2 + o(1)) \log_2 n$  and that since  $j \leq 3k/4$  it follows that  $2^{(j-1)/2} \leq n^{3/4+o(1)}$ . Recall that  $\mu = (1 + o(1))4k^4 \leq n^{0.01}$  and thus

$$\frac{\Delta_j}{\mu} \leq \mu \frac{\Delta_j}{\mu^2} \leq \frac{1}{n^{0.48}}$$

as claimed.

**Case 2:**  $j = k - i$ ,  $i \leq k/4$ .

In this case we also have

$$\Delta_j \leq \binom{n}{k} \binom{k}{j} \binom{n-k}{k-j} 2^{-2\binom{k}{2} + \binom{j}{2}}.$$

Indeed, there are

$$\binom{n}{k} \binom{k}{j} \binom{n-k}{k-j}$$

ways to choose the sets  $K, K'$  having intersection  $j = k - i$ , and for each such choice there is at most one way to place copies of  $F$  in each of them monotonely. The probability that this coincides with all edges and nonedges of the induced subgraph of  $G$  on  $K$  and on  $K'$  is  $2^{-2\binom{k}{2} + \binom{j}{2}}$ . Since  $\mu \geq 4k^4 > 1$  this implies that

$$\begin{aligned} \frac{\Delta_j}{\mu^2} &< \frac{\Delta_j}{\mu} \leq \binom{k}{j} \binom{n-k}{k-j} 2^{-\binom{k}{2} + \binom{j}{2}} \\ &\leq \binom{k}{i} \binom{n-k}{i} 2^{-i(k-i)} \leq (kn 2^{-(k-i)})^i \leq \frac{1}{n^{0.5-o(1)}} \leq \frac{1}{n^{0.48}}. \end{aligned}$$

This completes the proof of the claim.

Returning to the proof of the theorem, note that the variance of the random variable  $X$  satisfies

$$\text{Var}(X) \leq E(X) + \sum_{K, K'} \text{Cov}(X_K, X_{K'})$$

where the summation is over all ordered pairs  $K, K'$  where  $2 \leq |K \cap K'| \leq k - 1$  (since for all other pairs the covariance is zero). Since

$$\text{Cov}(X_K, X_{K'}) = E(X_K X_{K'}) - E(X_K)E(X_{K'}) \leq E(X_K X_{K'})$$

it follows from the claim above that

$$\text{Var}(X) \leq \mu + \Delta \leq (1 + o(1))\mu.$$

Therefore, by Chebyshev's Inequality, the probability that  $X \geq 3\mu/4$  is (much) bigger than  $3/4$ . By Markov's inequality, the probability that  $\Delta \leq \mu/4$  is also (much) bigger than  $3/4$  and hence with probability larger than  $1/2$ , both events happen simultaneously, that is, the number of copies of  $F$  in  $G$  is at least  $3\mu/4$  and the number of pairs of copies of  $F$  with nontrivial intersection is smaller than  $\mu/4$ . By removing one copy of  $F$  from each pair with a nontrivial intersection we conclude that if this is the case, then  $G$  contains a family of at least  $\mu/2$  copies of  $F$  with no two having a nontrivial intersection.

Let  $h(G)$  be the maximum cardinality of a family of copies of  $F$  in  $G$  in which no two members have a nontrivial intersection, and let  $Y$  be the random variable  $Y = h(G)$ . We next apply Talagrand's Inequality (Theorem 2.2) to deduce that the probability that  $Y$  is zero is tiny.

By the above discussion, the probability that  $Y$  is at least  $\mu/2$  exceeds  $1/2$ . It is also clear that the value of  $Y = h(G)$  can change by at most 1 if we add or delete one edge to  $G$ , and that  $h$  is  $f$ -certifiable where  $f(s) = s \binom{k}{2}$ . Therefore, by Theorem 2.2 with  $b = \mu/2$  and  $t = \sqrt{\mu}/k$  we conclude that the probability that  $Y = 0$  is smaller than

$$e^{-\mu/4k^2}$$

which is much smaller than  $2^{-k^2}$ . As  $Y = 0$  if and only if there is no (monotone) copy of  $F$  in  $G$ , and as the total number of labelled graphs  $F$  in  $\mathcal{F}(k)$  is  $2^{\binom{k}{2}}$  we conclude that  $G$  is induced universal for  $\mathcal{F}(k)$  with high probability. This completes the proof of Theorem 3.1.  $\square$

**Remark:** The quantity  $\binom{n}{k} 2^{-\binom{k}{2}}$  is exactly the expected number of cliques of size  $k$  in  $G = G(n, 0.5)$ . Therefore, if this number is  $o(1)$  then the probability that  $G$  contains such a clique is  $o(1)$ , by Markov's Inequality. It follows that the smallest  $n$  such that  $G(n, 0.5)$  is induced universal for  $\mathcal{F}(k)$  with high probability is at least  $(1+o(1)) \frac{k}{e} 2^{(k-1)/2}$ . By Theorem 3.1 this is tight up to the  $o(1)$ -term and the smallest  $n$  is in fact  $(1+o(1)) \frac{k}{e} 2^{(k-1)/2}$ . Note that the above estimate suffices to show that given  $n$ , the largest  $k$  so that the random graph  $G = G(n, 0.5)$  is induced universal for  $\mathcal{F}(k)$  with high probability is always one of two consecutive numbers and for "most" values of  $n$  it is a single number. It is well known that this is the case for the clique number of  $G$ , as proved in [24], [10]. The result here shows that for most values of  $n$ , with high probability, the value  $k$  which is the size of the largest clique in  $G$  is equal to the largest  $k$  so that  $G$  is induced universal for  $\mathcal{F}(k)$  (and for

every  $n$  these two numbers differ by at most 1 with high probability). The reason is that since the clique of size  $k$  and its complement have the largest number of automorphisms among all graphs of size  $k$ , once the random graph contains them, it typically contains also all the graphs on  $k$  vertices with a smaller number of automorphisms. A more precise version of this statement appears in the next subsection. Here “most” means that for any large  $N$ , if  $n$  is chosen randomly and uniformly among the numbers between  $N$  and  $2N$ , then with probability that tends to 1 as  $N$  tends to infinity there is a unique  $k$  so that the largest clique of  $G = G(n, 0.5)$  is  $k$  with high probability and  $G$  is induced universal for  $\mathcal{F}(k)$  (with the same  $k$ ) with high probability. Indeed, this is the case whenever  $n$  satisfies the following: if  $k_0$  is the largest  $k$  for which  $\binom{n}{k}2^{-\binom{k}{2}} \geq 1$  then in fact  $\binom{n}{k_0}2^{-\binom{k_0}{2}} \geq 4k_0^4$ .

It is worth noting that there are values of  $n$  for which the probability that the size of the largest clique of  $G = G(n, 0.5)$  is one more than the largest  $k$  for which  $G$  is induced universal for  $\mathcal{F}(k)$ , is bounded away from zero and one. Indeed, if  $\lambda > 0$  is bounded away from 0 and 1 and  $\binom{n}{k}2^{-\binom{k}{2}} = \lambda$ , then one can use the Stein-Chen method to conclude that the probability that the size of the maximum clique in  $G$  is  $k$ , is  $(1 + o(1))(1 - e^{-\lambda})$ . This is also the probability that the size of the maximum independent set in  $G$  is  $k$ , and these two events are nearly independent. In this case with probability  $(1 + o(1))e^{-\lambda}(1 - e^{-\lambda})$  the size of the maximum independent set is smaller by 1 than that of the maximum clique, and the largest  $k$  for which  $G$  is induced universal for  $\mathcal{F}(k)$  is, with high probability, the smaller of these two numbers. We omit the detailed argument, but mention that it is very similar to the one appearing in [2], Section 2.

### 3.2 A hitting time result

Erdős and Rényi [18] introduced the study of the evolution of the random graph, in which one considers a graph process starting with the empty graph on  $n$  labelled vertices and continuing by adding all  $\binom{n}{2}$  edges of the complete graph on these vertices one by one, according to a uniformly chosen random permutation. For any monotone graph property one can then study the hitting time of the property, that is, the smallest  $i$  so that after adding  $i$  edges the resulting graph satisfies the property. There are several known results showing that the hitting times of two monotone properties are, with high probability, identical. The best known result of this type, proved independently by Bollobás [9] and by Ajtai, Komlós and Szemerédi [1] (see also [22] for a statement of the result), is that the hitting times for having minimum degree 2 and being Hamiltonian are identical, with high probability. Another early result appears in [12], where it is shown that the hitting times for having minimum degree  $k$  and being  $k$ -connected are identical, with high probability.



Here we suggest a variant of the random graph process initiated by Erdős and Rényi, which we call the *vertex random graph process* (with parameter  $p$ ). For a probability  $p \in [0, 1]$  let  $G_1, G_2, G_3 \dots$  be the infinite sequence of random graphs defined as follows.  $G_1$  is the graph with 1 vertex  $v_1$ , and for each  $i \geq 1$ ,  $G_{i+1}$  is obtained from  $G_i$  by adding to  $G_i$  a new vertex  $v_{i+1}$ , where for each  $j \leq i$  randomly and independently,  $v_j v_{i+1}$  forms an edge with probability  $p$ . Therefore,  $G_n$  is distributed like the usual binomial random graph  $G(n, p)$ . When  $p = 0.5$  we do not mention the parameter  $p$  and refer to this process as the vertex random graph process. For a hereditary property  $P$ , the hitting time for not having  $P$  is the smallest  $i$  so that  $G_i$  does not satisfy  $P$ . In this terminology, we prove the following strengthening of Theorem 3.1.

**Theorem 3.2** *Let  $G_1, G_2, \dots$  be the vertex random graph process defined above and let  $k > 2$  be an integer. Then, with high probability (that is, with probability that tends to 1 as  $k$  tends to infinity) the hitting time for containing  $K_k$  and its complement  $\overline{K_k}$  as induced subgraphs is the same as that of the property of being induced universal for  $\mathcal{F}(k)$ . That is, with high probability as soon as  $G_i$  contains a clique and an independent set of size  $k$ , it already contains all graphs on  $k$  vertices as induced subgraphs.*

The proof combines a modification of the argument in the proof of Theorem 3.1 with some group theoretic tools that supply an upper bound for the number of  $k$  vertex graphs with very large automorphism groups, as stated in the following lemma. As before we assume here, whenever this is needed, that  $k$  is sufficiently large.

**Lemma 3.3** (i) *A  $k$ -vertex graph  $H$  has  $k!$  automorphisms if and only if  $H = K_k$  or  $H = \overline{K_k}$ . In any other case the number of automorphisms is at most  $(k-1)!$*   
(ii) *For any constant  $b \geq 1$  there is a constant  $B = B(b)$  so that the number of  $k$ -vertex graphs with at least  $k!/k^b$  automorphisms is at most  $B$ .*

**Proof:** Let  $H$  be a  $k$ -vertex graph, let  $A$  denote its group of automorphisms and put  $s = |A|$ . Thus  $A$  is a group of permutations acting on the set of  $k$  vertices of  $H$ , which we denote by  $[k] = \{1, 2, \dots, k\}$ . Clearly if  $A$  is doubly transitive then  $H$  is either the complete graph  $K_k$  or the empty graph  $\overline{K_k}$ . If  $A$  is primitive but not doubly transitive then, as proved by Babai in [6],  $|A| \leq e^{4\sqrt{k} \ln^2 k}$  which is much smaller than  $k!/k^b$  for any fixed  $b$  provided  $k$  is sufficiently large. (Note that we do not need the nearly tight result of Babai here, the estimates in [8] (see also [30], [16]) or in [27] suffice).

If  $A$  is not primitive, but is transitive, then there is partition of the set  $[k]$  into blocks of equal size exceeding 1, call it  $t$ , so that each permutation in  $A$  preserves the block

structure. Therefore, in this case,  $|A| = s \leq (k/t)!(t!)^{k/t}$ . Without trying to optimize the estimate we show that in this case

$$s \leq \frac{k!}{2^{k/2-1}}. \quad (5)$$

Indeed, since both  $t$  and  $k/t$  are between 2 and  $k/2$  we have

$$\begin{aligned} \frac{s}{k!} &\leq \frac{(k/t)!(t!)^{k/t}}{k!} = \frac{[t(t-1)\cdots 2]^{k/t}}{k(k-1)\cdots(k/t+1)} \\ &\leq \frac{[t(t-1)\cdots 2]^{k/t-1}}{k(k-1)\cdots(k/t+t)} \leq \frac{[(t+2)/2]^{(t-1)(k/t-1)}}{(t+2)^{k-k/t-t+1}} \leq (1/2)^{k-k/t-t+1} \leq (1/2)^{k/2-1}, \end{aligned}$$

as claimed.

Finally, if  $A$  is not transitive, and its largest orbit is of size  $m < k$  then  $s \leq (k-m)!m!$

Part (i) follows, since by the above arguments if  $A$  is not doubly transitive then if it is primitive then  $s$  is far smaller than  $(k-1)!$ , this is the case also if it is transitive but not primitive, and finally if it is not transitive then  $s$  is at most  $(k-1)!$

To prove part (ii) note that if  $s$  is at least  $k!/k^b$  and  $A$  is not doubly transitive then it cannot be primitive and cannot be transitive but not primitive, as in both cases  $s$  is much smaller. Thus  $A$  is not transitive. If its largest orbit is of size smaller than  $k-b$ , then  $s \leq (b+1)!(k-b-1)! < k!/k^b$  hence the largest orbit must be of size  $m \geq k-b$ . The group  $A$  must act primitively and hence doubly transitively on this orbit, since otherwise, again,  $s$  is far too small. Thus  $H$  contains a clique or an independent set of size  $m \geq k-b$  on the set  $W$  of the vertices in this orbit. Let the other vertices of  $H$  be  $u_1, u_2, \dots, u_g$ , where  $g \leq b$ . Split the vertices of  $W$  into at most  $2^g$  disjoint subsets according to their neighbors in  $U = \{u_1, u_2, \dots, u_g\}$ . If the largest subset is of size smaller than  $k-b-1$  then, again,  $s$  can be at most some  $C(b)(k-b-1)! < k!/k^b$ , contradiction. Thus the largest subset is of size at least  $k-b$ . There are at most

$$2^{\binom{g}{2}} < 2^{b^2/2}$$

possibilities for the induced subgraph of  $H$  on  $U = \{u_1, u_2, \dots, u_g\}$ , and less than

$$\binom{2^g + b}{b} < 2^{b^2}$$

possibilities to choose the numbers of vertices of  $W$  in each of the above subsets. The product of these two bounds is an upper bound for the number  $B = B(b)$  of  $k$ -vertex graphs with at least  $k!/k^b$  automorphisms, completing the proof of the lemma.  $\square$

**Lemma 3.4** *Let  $H$  be a  $k$ -vertex graph, suppose it is neither complete nor empty, and let  $s$  denote the number of its automorphisms. Assume  $s \geq \frac{k!}{k^5}$  and let  $n$  be an integer so that*

$$\binom{n}{k} 2^{-\binom{k}{2}} = \frac{1 + o(1)}{\sqrt{k}}.$$

*Then the probability that the random graph  $G = G(n, 0.5)$  contains no induced copy of  $H$  is at most  $\frac{1+o(1)}{\sqrt{k}}$ .*

**Proof:** By Lemma 3.3, part (i),  $s \leq (k-1)!$  Thus, the expected number of induced copies of  $H$  in  $G$ , which we denote by  $\mu$ , satisfies

$$\mu = \binom{n}{k} \frac{k!}{s} 2^{-\binom{k}{2}}, \quad (1 + o(1))\sqrt{k} \leq \mu \leq (1 + o(1))k^{4.5}.$$

As in the proof of Theorem 3.1, the expected number of pairs of induced copies of  $H$  with a nontrivial intersection is at most  $\Delta = \sum_{j=2}^{k-1} \Delta_j$ , where

$$\Delta_j \leq \binom{n}{k} \frac{k!}{s} 2^{-\binom{k}{2}} \binom{k}{j} \binom{n-k}{k-j} \frac{k!}{s} 2^{-\binom{k}{2} + \binom{j}{2}}.$$

Considering separately the cases  $j \leq 3k/4$  and  $j = k - i, i \leq k/4$  it is easy to check, following the computation in the proof of Theorem 3.1 with the obvious modifications, that  $\Delta < \frac{\mu}{n^{0.4}}$  (with room to spare). Therefore, if  $X$  is the random variable counting the number of induced copies of  $H$  in  $G$  then the expectation of  $X$  is  $\mu \geq (1 + o(1))\sqrt{k}$  and its variance is at most  $\mu + \Delta = (1 + o(1))\mu$ . The desired result follows from Chebyshev's Inequality.  $\square$

**Lemma 3.5** *Let  $H$  be a  $k$ -vertex graph with  $s \leq \frac{k!}{k^5}$  automorphisms. Let  $n$  be, as before, an integer so that*

$$\binom{n}{k} 2^{-\binom{k}{2}} = \frac{1 + o(1)}{\sqrt{k}}.$$

*Then the probability that the random graph  $G = G(n, 0.5)$  contains no induced copy of  $H$  is at most  $e^{-k^2}$*

**Proof:** Here too the proof is similar to that of Theorem 3.1, with one extra twist, as follows. Define a real  $p \in (0, 1)$  so that

$$\binom{n}{k} \frac{k!}{s} 2^{-\binom{k}{2}} p = 4k^4.$$

Note that by the assumptions on  $n$  and  $s$ ,  $p$  is at most  $\frac{4+o(1)}{\sqrt{k}}$  and  $\frac{k!}{s}p = (4 + o(1))k^{4.5}$ . Let  $\mathcal{H}$  be a random collection of induced copies of  $H$  in the random graph  $G = G(n, 0.5)$  obtained by picking, randomly and independently, each induced copy of  $H$  to be a member of  $\mathcal{H}$  with probability  $p$ . The expected number of copies of  $H$  in  $\mathcal{H}$ , denoted by  $\mu$ , is exactly

$$\mu = \binom{n}{k} \frac{k!}{s} 2^{-\binom{k}{2}} p = 4k^4.$$

Let  $Z$  be the random variable counting the number of ordered pairs of members of  $\mathcal{H}$  with a nontrivial intersection. Then the expectation of  $Z$  is  $\Delta = \sum_{j=2}^{k-1} \Delta_j$  where

$$\Delta_j \leq \binom{n}{k} \frac{k!}{s} 2^{-\binom{k}{2}} p \binom{k}{j} \binom{n-k}{k-j} \frac{k!}{s} 2^{-\binom{k}{2} + \binom{j}{2}} p$$

As before, it is not difficult to check that each  $\Delta_j$  is smaller than  $\mu \frac{1}{n^{0.41}}$  and thus  $\Delta \leq \frac{\mu}{n^{0.4}} = o(\mu) = o(\mu^2)$ .

Therefore, by Chebyshev's Inequality, the probability that the size of  $\mathcal{H}$  is at least  $3\mu/4 = 3k^4$  is (much) bigger than  $3/4$ . By Markov's inequality, the probability that  $\Delta \leq \mu/4$  is also (much) bigger than  $3/4$  and hence with probability larger than  $1/2$ , both events happen simultaneously, that is, the size of  $\mathcal{H}$  is at least  $3\mu/4$  and the number of pairs of members of  $\mathcal{H}$  with nontrivial intersection is smaller than  $\mu/4$ . By removing one copy of  $H$  from each pair with a nontrivial intersection we conclude that if this is the case, then  $G$  contains a family of at least  $\mu/2 = 2k^4$  copies of  $H$  with no two having a nontrivial intersection.

Let  $h(G)$  be the maximum cardinality of a family of copies of  $H$  in  $G$  in which no two members have a nontrivial intersection, and let  $Y$  be the random variable  $Y = h(G)$ .

By the above discussion, the probability that  $Y$  is at least  $\mu/2 = 2k^4$  exceeds  $1/2$ . It is also clear that the value of  $Y = h(G)$  can change by at most 1 if we add or delete one edge to  $G$ , and that  $h$  is  $f$ -certifiable where  $f(s) = s \binom{k}{2}$ . Therefore, by Talagrand's Inequality (Theorem 2.2) with  $b = \mu/2$  and  $t = \sqrt{\mu}/k$  we conclude that the probability that  $Y = 0$  is smaller than

$$e^{-\mu/4k^2} = e^{-k^2}.$$

Since if  $Y > 0$  then  $G$  contains an induced copy of  $H$  this completes the proof of the lemma.  $\square$

**Proof of Theorem 3.2:** For a given (large)  $k$  let  $n$  be the smallest integer so that

$$\binom{n}{k} 2^{-\binom{k}{2}} \geq \frac{1}{\sqrt{k}}.$$

It is easy to see that in fact the left hand side is  $\frac{1+o(1)}{\sqrt{k}}$ . By Markov's Inequality, the probability that in the vertex random graph process the graph number  $n$ ,  $G_n$ , contains a clique of size  $k$  is at most  $\frac{1+o(1)}{\sqrt{k}} = o(1)$ . On the other hand we claim that with high probability  $G_n$  contains an induced copy of every  $k$  vertex graph  $H$  besides  $K_k, \overline{K}_k$ . For the graphs  $H$  with at least  $\frac{k!}{k^5}$  automorphisms this follows from Lemma 3.4 and Lemma 3.3, part (ii), as there are only  $O(1)$  such graphs and the probability of not containing each given one is at most  $\frac{1+o(1)}{\sqrt{k}}$ . For the graphs  $H$  with a smaller number of automorphisms this follows from Lemma 3.5, as there are less than  $2^{k^2/2}$  such graphs and the probability of not containing a given one is at most  $e^{-k^2}$ . This completes the proof.  $\square$

Theorem 3.2 and its proof together with the Stein-Chen method provide a sharp estimate for the probability that the random graph  $G = G(n, 0.5)$  is induced universal for  $\mathcal{F}(k)$ . This is stated in Theorem 1.2 in Section 1. The proof, which is a short consequence of the proof of Theorem 3.2 follows.

**Proof of Theorem 1.2:** Fix a small  $\epsilon > 0$ . By Markov's Inequality the probability that  $G$  contains a copy of  $K_k$  is at most  $\lambda$ , hence if  $\lambda < \epsilon$  then  $(1 - e^{-\lambda})^2$  (which is close to 0) is indeed a good approximation of the desired probability. Similarly, if  $\lambda > 1/\epsilon$  is large, then by the known results of [10], [24] (proved by the second moment method) the probability that  $G$  contains a copy of  $K_k$  and of  $\overline{K}_k$  is very close to 1 and the quantity  $(1 - e^{-\lambda})^2$  is a good approximation of that. By Theorem 3.2 and its proof the probability that  $G$  is induced universal for  $\mathcal{F}(k)$  is very close to the probability it contains the two graphs above. If  $\lambda$  is neither tiny nor huge the desired result follows from this fact and the Stein-Chen method which gives that the probability that  $G$  contains a copy of  $K_k$  is  $(1 + o(1))(1 - e^{-\lambda})$ , that this is also the probability that it contains an induced copy of  $\overline{K}_k$ , and that these two events are nearly independent. The detailed computation required for applying the Stein-Chen method is omitted. As mentioned in the remark following the proof of Theorem 3.1 this computation is very similar to the one in [2], Section 2.  $\square$

## 4 Extensions

The proof of Theorem 1.1 can be extended to supply similar tight estimates for several related questions. In this section we describe some examples. In most of them the adaptation of the proof is simple, the only case that requires several additional ideas is that of bipartite graphs, described in subsection 4.5.

## 4.1 Tournaments

A *tournament* on  $k$  vertices is an orientation of a complete graph on  $k$  vertices, that is, a directed graph on  $k$  vertices so that for every pair  $u, v$  of distinct vertices there is either a directed edge from  $u$  to  $v$  or a directed edge from  $v$  to  $u$  (but not both). It is clear that the number of tournaments on  $k$  labelled vertices is  $2^{\binom{k}{2}}$ , and therefore the number of pairwise non-isomorphic tournaments on  $k$  vertices is at least

$$\frac{2^{\binom{k}{2}}}{k!}.$$

Let  $\mathbf{T}_k$  denote the set of all tournaments on  $k$  vertices. Call a tournament  $G$  *induced universal* for  $\mathbf{T}_k$  if it contains every member of  $\mathbf{T}_k$  as an induced sub-tournament, and let  $t(k)$  denote the minimum possible number of vertices of such a universal tournament. The obvious counting argument shows that  $t(k) \geq 2^{(k-1)/2}$  and Moon [26] showed that  $t(k) \leq O(k2^{k/2})$ . This has been improved in [4], where it is proved that  $t(k) \leq 16 \cdot 2^{\lceil k/2 \rceil}$ . Our method here suffices to determine  $t(k)$  up to a lower order additive term.

**Theorem 4.1** *The minimum possible number of vertices  $t(k)$  of a tournament that contains a copy of every  $k$ -vertex tournament satisfies*

$$2^{(k-1)/2} \leq t(k) \leq 2^{(k-1)/2} \left(1 + O\left(\frac{\log^{3/2} k}{\sqrt{k}}\right)\right).$$

Therefore  $t(k) = (1 + o(1))2^{(k-1)/2}$ .

The proof is similar to that of Theorem 1.1. We first show, using Talagrand's Inequality, that a random tournament on  $n$  vertices, where  $n$  is as in (1), contains, with high probability, a copy of every tournament on  $k$  vertices in which any induced subgraph has less than  $k^8 \sqrt{k \log k}$  automorphisms. Next we prove that there is a much smaller tournament that contains a copy of all the  $k$ -vertex tournaments containing a subgraph with that many automorphisms. This is done by proceeding as in Section 2, using the large automorphism group to describe an appropriate labeling scheme for these tournaments. The desired universal tournament is the vertex disjoint union of the random tournament with the smaller one above. We omit the details.

## 4.2 Directed graphs

Let  $\mathcal{D}(k)$  denote the set of all directed graphs on  $k$  vertices. Any two vertices  $u, v$  in a member  $D \in \mathcal{D}(k)$  can be either nonadjacent, or connected by a directed edge from  $u$  to

$v$ , or by one from  $v$  to  $u$ , or by both. Therefore the cardinality of  $\mathcal{D}(k)$  is at least

$$\frac{4^{\binom{k}{2}}}{k!}.$$

A directed graph  $G$  is induced universal for  $\mathcal{D}(k)$  if every member of  $\mathcal{D}(k)$  is an induced subgraph of  $G$ . Let  $d(k)$  denote the minimum possible number of vertices of such a graph. Simple counting shows that  $d(k) \geq 2^{k-1}$ . In [4] the authors prove that this is tight up to a constant factor, proving that  $d(k) \leq 16 \cdot 2^{k-1}$  and mention the open problem of closing the gap between the upper and lower bound. Our method here gives the following tight estimate.

**Theorem 4.2** *The minimum possible number of vertices  $d(k)$  of a directed graph that contains a copy of every  $k$ -vertex directed graph satisfies*

$$2^{k-1} \leq d(k) \leq 2^{k-1} \left(1 + O\left(\frac{\log^{3/2} k}{\sqrt{k}}\right)\right).$$

Therefore  $d(k) = (1 + o(1))2^{k-1}$ .

The proof is a straightforward modification of the ones of Theorem 1.1 and Theorem 4.1.

### 4.3 Oriented graphs

Let  $\mathcal{O}(k)$  denote the set of all oriented graphs on  $k$  vertices. Any two vertices  $u, v$  in a member  $O \in \mathcal{O}(k)$  can be either nonadjacent, or connected by a directed edge from  $u$  to  $v$ , or by one from  $v$  to  $u$ , but not by both. Therefore the cardinality of  $\mathcal{O}(k)$  is at least

$$\frac{3^{\binom{k}{2}}}{k!}.$$

An oriented graph  $G$  is induced universal for  $\mathcal{O}(k)$  if every member of  $\mathcal{O}(k)$  is an induced subgraph of  $G$ . Let  $r(k)$  denote the minimum possible number of vertices of such a graph. Simple counting shows that  $r(k) \geq 3^{\binom{k-1}{2}}$ . Our technique shows that this is tight up to a small additive error. The proof, which is a simple variant of the previous ones, is omitted.

**Theorem 4.3** *The minimum possible number of vertices  $r(k)$  of an oriented graph that contains an induced copy of every  $k$ -vertex oriented graph satisfies*

$$3^{\binom{k-1}{2}} \leq r(k) \leq 3^{\binom{k-1}{2}} \left(1 + O\left(\frac{\log^{3/2} k}{\sqrt{k}}\right)\right).$$

Therefore  $r(k) = (1 + o(1))3^{\binom{k-1}{2}}$ .

#### 4.4 Edge colored complete graphs

For a fixed positive integer  $r$ , let  $\mathcal{K}(k, r)$  denote the set of all complete graphs on  $k$  vertices whose edges are colored by the  $r$  colors  $\{1, 2, \dots, r\}$ . The number of members of  $\mathcal{K}(r)$  is thus at least

$$\frac{r^{\binom{k}{2}}}{k!}.$$

Note that the case  $r = 2$  is equivalent to that of all undirected graphs on  $k$  vertices. Let  $kr(k)$  denote the minimum possible number of vertices of an induced universal graph  $G$  for  $\mathcal{K}(r, k)$ , that is, the minimum number of vertices of a complete  $r$ -edge colored graph that contains every member of  $\mathcal{K}(r, k)$  as an induced subgraph. Counting shows that  $kr(k) \geq r^{\binom{k-1}{2}}$ . Our method here gives

**Theorem 4.4** *For every fixed  $r$  and large  $k$ , the function  $kr(k)$  satisfies*

$$r^{\binom{k-1}{2}} \leq kr(k) \leq r^{\binom{k-1}{2}} \left(1 + O\left(\frac{\log^{3/2} k}{\sqrt{k}}\right)\right).$$

Therefore  $kr(k) = (1 + o(1))r^{\binom{k-1}{2}}$ .

#### 4.5 Bipartite graphs

Let  $\mathcal{B}(k)$  denote the set of all bipartite graphs  $B = (U_1, U_2, E)$  on  $k$  vertices. Here the vertex classes of  $B$  are  $U_1, U_2$ . Call  $B$  a  $(k_1, k_2)$ -bipartite graph if  $|B_1| = k_1$  and  $|B_2| = k_2$  (where  $k_1 + k_2 = k$ ). A bipartite graph  $G$  with vertex classes  $V_1, V_2$ , each of size  $n$ , is an induced universal graph for  $\mathcal{B}(k)$  if for any member  $B = (U_1, U_2, E)$  of  $\mathcal{B}(k)$  there are sets  $U'_1 \subset V_1$  and  $U'_2 \subset V_2$  and bijections from  $U_1$  to  $U'_1$  and from  $U_2$  to  $U'_2$  which map  $B$  to an isomorphic induced subgraph of  $G$ . Note that we insist here that the vertices of  $U_1$  are mapped to vertices in  $V_1$ , and the same holds for  $U_2, V_2$ . Allowing arbitrary mappings from the vertices of  $B$  to those of  $G$  is also possible, and does not cause any significant changes, hence we prefer to consider the definition given above. Let  $b(k)$  denote the smallest possible  $n$  so that an induced universal bipartite graph as above for  $\mathcal{B}(k)$  exists. Since this graph has to contain all  $(\lfloor k/2 \rfloor, \lceil k/2 \rceil)$ -bipartite graphs as induced subgraphs, a simple counting argument shows that for even  $k$ ,  $b(k) \geq 2^{k/4}$  whereas for odd  $k$ ,

$$b(k) \geq 2^{(k^2-1)/4k} = 2^{k/4} (1 - O(1/k)).$$

In [23] it is shown that  $b(k) \leq O(k^2 2^{k/4})$  and in [4] it is proved that  $b(k) \leq c 2^{k/4}$  for some absolute constant  $c$  (which is not specified in the paper, but is certainly less than 100.) Here we show that the lower bound is tight, up to a lower order additive term.



**Theorem 4.5** *The function  $b(k)$  satisfies*

$$2^{k/4}(1 - O(1/k)) \leq b(k) \leq 2^{k/4}(1 + O(\frac{\log^{3/2} k}{\sqrt{k}}))$$

*Therefore  $b(k) = (1 + o(1))2^{k/4}$ .*

The proof is similar to the previous ones, but requires a few additional arguments. We first note that as shown in [4], it is easy to deal with all unbalanced bipartite graphs. This is stated in the following lemma.

**Lemma 4.6** ([4], **Theorem 8.1**) *For every  $r$  there is a bipartite graph  $G_r$  with classes of vertices  $V_1, V_2$ , each of size at most  $n = 2k \cdot 2^{k/4 - r^2/k}$ , that contains every  $(k/2 - r, k/2 + r)$ -bipartite graph  $(U_1, U_2, E)$  as an induced subgraph (with  $U_1$  embedded in  $V_1$  and  $U_2$  in  $V_2$ .)*

The proof is by orienting the edges of the complete bipartite graph with classes of vertices of sizes  $k/2 - r$  and  $k/2 + r$  so that every outdegree is at most  $k/4 - r^2/k + 1$ . Such an orientation exists by Fact 2 in subsection 2.2. We can then label each vertex by its number (which is a number not exceeding  $k$ ) and by the bits describing its adjacency relations to its outneighbors in the above orientation, in order. This assigns to each vertex a label with at most  $\log_2 k + k/4 - r^2/k + 1$  bits, and the labels of any two vertices suffice to determine whether or not they are adjacent. The existence of the desired graph  $G_r$  now follows from the standard simple connection between adjacency labeling schemes and induced universal graphs.

Note that by taking the vertex disjoint union of the graphs  $G_r$  for all  $r$  (positive or negative) satisfying, say,  $2\sqrt{k \log k} \leq |r| (< k/2)$  the total number of vertices in each side is less than  $2^{k/4}/k^2$  which is smaller than  $(1 + o(1))b(k)/k^2$ . It thus remains to deal with the  $(k_1, k_2)$ -bipartite graphs in which  $k_1$  and  $k_2$  are close to each other, hence we assume from now on that each of them is at least, say,  $0.45k$ .

As in the proof of Theorem 1.1, we deal separately with bipartite graphs in which every large induced subgraph has a relatively small number of automorphisms, which we call asymmetric, and with the others, called symmetric. The asymmetric graphs will be contained in a random bipartite graph, and the symmetric ones in a smaller graph, constructed using an appropriate adjacency labelling scheme. We proceed with the details. In the rest of this section an automorphism of a bipartite graph  $B = (U_1, U_2, E)$  with vertex classes  $U_1, U_2$  means an automorphism that maps each  $U_i$  to itself. Recall that in all  $(k_1, k_2)$  bipartite graphs considered from now on  $k_1 + k_2 = k$  and each  $k_i$  is between  $0.45k$  and  $0.55k$ .

Call a  $(k_1, k_2)$ -bipartite graph on  $k_1 + k_2 = k$  vertices *asymmetric* if any induced subgraph of it on at least  $0.9k$  vertices has at most  $k^{8m}$  automorphisms, where  $m = 2\sqrt{k \log k}$ . Note that, in particular, the number of automorphisms of any such graph is at most  $k^{8m}$ . Let  $\mathcal{B}'(k)$  denote the family of all asymmetric bipartite graphs on  $k$  vertices.

Let  $n$  be the smallest integer that satisfies the following inequality

$$\binom{n}{\lfloor k/2 \rfloor} \binom{n}{\lceil k/2 \rceil} \frac{\lfloor k/2 \rfloor! \lceil k/2 \rceil!}{k^{16m}} 2^{-\lfloor k/2 \rfloor \cdot \lceil k/2 \rceil} \geq 1. \quad (6)$$

When increasing  $n$  to  $n+1$ , the left-hand-side of the last inequality increases by a factor of  $1+o(1)$  for the relevant parameters, and thus it follows that the above smallest  $n$  satisfies

$$n = 2^{k/4}(1 + O(1/k)).$$

In particular,  $n = (1 + o(1))2^{k/4}$ .

**Theorem 4.7** *Let  $n$  be as above and let  $G = (V_1, V_2, E) = G(n, n, 0.5)$  be the random bipartite graph with vertex classes  $V_1, V_2$ , each of size  $n$ , where each pair of vertices  $v_1 \in V_1$  and  $v_2 \in V_2$  forms an edge randomly and independently with probability  $1/2$ . Then, with high probability  $G$  is an induced universal bipartite graph for  $\mathcal{B}'(k)$ .*

As in the proof of Theorem 2.1, the proof here is based on Talagrand's Inequality (Theorem 2.2).

We need the following lemma.

**Lemma 4.8** *Let  $B \in \mathcal{B}'(k)$  have  $s$  automorphisms. Let the sizes of the vertex classes of  $B$  be  $k_1, k_2$  (and recall that each  $k_i$  is at least  $0.45k$ .) Suppose  $j_1, j_2$  are integers satisfying  $1 \leq j_1 = k_1 - i_1 \leq k_1$ ,  $1 \leq j_2 = k_2 - i_2 \leq k_2$ ,  $j_1 + j_2 = j = k - i$  and assume  $1 \leq i \leq 0.1k$ . Let  $K_1, K'_1$  be two sets of vertices, each of size  $k_1$ , and let  $K_2, K'_2$  be two sets of vertices, each of size  $k_2$ , where  $|K_1 \cap K_2| = j_1$ ,  $|K'_1 \cap K'_2| = j_2$  and  $K_1, K'_1$  do not intersect  $K_2, K'_2$ . Then the number of ways to choose the edges and nonedges among all  $2k_1k_2 - j_1j_2$  pairs consisting of a vertex of  $K_1$  and a vertex of  $K_2$  or a vertex of  $K'_1$  and a vertex of  $K'_2$  so that the induced subgraph on  $K_1 \cup K_2$  is isomorphic to  $B$  and the induced subgraph on  $K'_1 \cup K'_2$  is also isomorphic to  $B$  (and both isomorphisms map the first vertex class of  $B$  to  $K_1$  and to  $K'_1$ ) is at most*

$$\frac{k_1!k_2!}{s} k^{i_1+i_2} k^{8m}.$$

**Proof:** There are exactly  $\frac{k_1!k_2!}{s}$  ways to place a copy of  $B$  on  $K_1 \cup K_2$ . For each such fixed copy, we bound the number of embeddings of  $B$  in  $K'_1 \cup K'_2$ . There are at most

$$k_1(k_1 - 1) \cdots (k_1 - i_1 + 1) k_2(k_2 - 1) \cdots (k_2 - i_2 + 1) < k^{i_1+i_2}$$

ways to choose the vertices of  $B$  mapped to the vertices of  $K'_1 - K_1$  and to the vertices of  $K_2 - K'_2$ . Fix a set  $T$  of these  $i$  vertices and their embedding. In order to complete the embedding, the induced subgraph of the copy of  $B$  placed in  $K_1 \cup K_2$  on the set of vertices  $(K_1 \cap K'_1) \cup (K_2 \cap K'_2)$  has to be isomorphic to the induced subgraph of  $B$  on  $V(B) - T$ . If so, then the number of ways to embed these  $k - i$  vertices is the number of automorphisms of this induced subgraph of  $B$ , which is, by the definition of  $\mathcal{B}'(k)$ , at most  $k^{8m}$ .  $\square$

**Proof of Theorem 4.7:** Let  $B$  be a fixed member of  $\mathcal{B}'(k)$ , let  $k_1, k_2$  denote the number of vertices in its vertex classes, and let  $s$  be the size of its automorphism group. Then

$$s \leq k^{8m} = k^{16\sqrt{k \log k}}.$$

Let  $G = (V_1, V_2, E)$  be the random bipartite graph  $G(n, n, 0.5)$ , where  $n$  is as chosen in (6). For every pair of subsets  $K_1 \subset V_1$ ,  $K_2 \subset V_2$  of sizes  $|K_1| = k_1$ ,  $|K_2| = k_2$ , let  $X_{K_1, K_2}$  denote the indicator random variable whose value is 1 if the induced subgraph of  $G$  on  $K_1 \cup K_2$  is isomorphic to  $B$  and let  $X = \sum_{K_1, K_2} X_{K_1, K_2}$ , where the summation is over all pairs of subsets  $K_1 \subset V_1$  of cardinality  $k_1$  and  $K_2 \subset V_2$  of cardinality  $k_2$ . Thus  $X$  is the number of copies of  $B$  in  $G$ . The expectation of each  $X_{K_1, K_2}$  is clearly

$$E(X_{K_1, K_2}) = \frac{k_1!k_2!}{s} 2^{-k_1 k_2}.$$

Thus, by linearity of expectation,

$$E(X) = \binom{n}{k_1} \binom{n}{k_2} \frac{k_1!k_2!}{s} 2^{-k_1 k_2} \geq k^{8m}$$

where in the last inequality we used (6) and the fact that  $s \leq k^{8m}$ . Note also that since

$$k_1 k_2 \geq \frac{k^2}{4} - \frac{k^2}{400}$$

it follows that the expectation of  $X$  is at most  $(1 + o(1))k^{16m} 2^{k^2/400} < n^{0.02}$  (even if  $s = 1$ ).

We say that two copies of  $B$  in  $G$  have a *nontrivial intersection* if they share at least one vertex in each vertex class. Put  $\mu = E(X)$ . Let  $Z$  denote the random variable  $Z = \sum_{K, K'} X_{K_1, K_2} X_{K'_1, K'_2}$  where the summation is over all (ordered) pairs  $K = (K_1, K_2)$  and  $K' = (K'_1, K'_2)$  that satisfy  $K_1 \cap K'_1 \neq \emptyset$  and  $K_2 \cap K'_2 \neq \emptyset$ . Thus,  $Z$  is the number of pairs of copies of  $B$  in  $G$  that have a nontrivial intersection. We next compute the expectation of  $Z$  and show that it is much smaller than  $\mu$  (and hence also much smaller than  $\mu^2$ ). Put  $\Delta = E(Z)$ . Then  $\Delta = \sum_{j_1, j_2} \Delta_{j_1, j_2}$ , where the summation is over all  $(j_1, j_2)$  satisfying  $1 \leq j_1 \leq k_1$ ,  $1 \leq j_2 \leq k_2$  and  $j_1 + j_2 < k_1 + k_2 = k$ . Here  $\Delta_{j_1, j_2}$  is the

expected number of pairs  $K = (K_1, K_2), K' = (K'_1, K'_2)$  with  $X_{K_1, K_2} = X_{K'_1, K'_2} = 1$  and  $|K_1 \cap K'_1| = j_1, |K_2 \cap K'_2| = j_2$ .

**Claim:** For all admissible  $j_1, j_2$

$$\Delta_{j_1, j_2} \leq \mu \frac{1}{n^{0.1}} \quad (7)$$

**Proof of Claim:** Consider two possible cases, as follows.

**Case 1:**  $j_1 + j_2 \leq 0.9k$ .

In this case

$$\Delta_{j_1, j_2} \leq \binom{n}{k_1} \binom{n}{k_2} \binom{k_1}{j_1} \binom{k_2}{j_2} \binom{n-k_1}{k_1-j_1} \binom{n-k_2}{j_2-k_2} \frac{k_1!k_2!}{s} \frac{k_1!k_2!}{s} 2^{-2k_1k_2+j_1j_2}.$$

Indeed, there are  $\binom{n}{k_1} \binom{n}{k_2}$  ways to choose the sets  $K_1, K_2$ . For each such choice there are

$$\binom{k_1}{j_1} \binom{k_2}{j_2} \binom{n-k_1}{k_1-j_1} \binom{n-k_2}{j_2-k_2}$$

ways to choose  $K'_1, K'_2$  with  $|K_1 \cap K'_1| = j_1, |K_2 \cap K'_2| = j_2$ . There are  $\frac{k_1!k_2!}{s}$  ways to place a copy of  $B$  in  $K_1 \cup K_2$  and  $\frac{k_1!k_2!}{s}$  ways to place a copy of  $B$  in  $K'_1 \cup K'_2$  (this is an overcount, as these two copies have to agree on the edges in their common part). This determines all the edges and nonedges in the induced subgraph on  $K_1 \cup K_2$  and on  $K'_1 \cup K'_2$ . The probability that the bipartite graph  $G$  indeed has exactly these edges is

$$2^{-2k_1k_2+j_1j_2}.$$

Therefore,

$$\begin{aligned} \frac{\Delta_{j_1, j_2}}{\mu^2} &\leq k_1^{j_1} k_2^{j_2} \left(\frac{k_1}{n}\right)^{j_1} \left(\frac{k_2}{n}\right)^{j_2} 2^{j_1j_2} < \left(\frac{k^2}{n}\right)^{j_1+j_2} 2^{(j_1+j_2)/2} = \left[\frac{k^2 2^{(j_1+j_2)/4}}{n}\right]^{j_1+j_2} \\ &< \left(\frac{1}{n^{0.1-o(1)}}\right)^2 < \frac{1}{n^{0.15}}. \end{aligned}$$

Here we used the fact that  $k = (4 + o(1)) \log_2 n$  and that  $2^{(j_1+j_2)/4} \leq 2^{0.9k/4}$ . Recall that  $\mu \leq n^{0.02}$  and thus

$$\frac{\Delta_{j_1, j_2}}{\mu} \leq \mu \frac{\Delta_{j_1, j_2}}{\mu^2} < \frac{1}{n^{0.1}}$$

as claimed.

**Case 2:**  $j_1 = k_1 - i_1, j_2 = k_2 - i_2$   $1 \leq i = i_1 + i_2 \leq 0.1k$ .

In this case we have, by Lemma 4.8

$$\Delta_{j_1, j_2} \leq \binom{n}{k_1} \binom{n}{k_2} \binom{k_1}{j_1} \binom{n-k_1}{k_1-j_1} \binom{k_2}{j_2} \binom{n-k_2}{k_2-j_2} \frac{k_1!k_2!}{s} k^{i_1+i_2} k^{8m} 2^{-2k_1k_2+j_1j_2}.$$

Indeed, there are

$$\binom{n}{k_1} \binom{n}{k_2} \binom{k_1}{j_1} \binom{n-k_1}{k_1-j_1} \binom{k_2}{j_2} \binom{n-k_2}{k_2-j_2}$$

ways to choose the sets  $K_1, K_2, K'_1, K'_2$  with the required intersections, and by Lemma 4.8 for each such choice there are at most  $\frac{k_1!k_2!}{s} k^{i_1+i_2} k^{8m}$  ways to place copies of  $B$  in each of them. The probability that this coincides with all edges and nonedges of the induced subgraph of  $G$  on  $K_1 \cup K_2$  and on  $K'_1 \cup K'_2$  is  $2^{-2k_1k_2+j_1j_2}$ .

Since  $\mu \geq k^{8m} > 1$  this implies that

$$\begin{aligned} \frac{\Delta_{j_1, j_2}}{\mu^2} &< \frac{\Delta_{j_1, j_2}}{\mu} \leq \binom{k_1}{i_1} \binom{n-k_1}{i_1} \binom{k_2}{i_2} \binom{n-k_2}{i_2} k^{i_1+i_2} k^{8m} 2^{-k_1k_2+j_1j_2} \\ &< (k^2n)^{i_1+i_2} n^{o(1)} 2^{-i_2(k_1-0.5i_1)-i_1(k_2-0.5i_2)} \\ &< (k^2n)^{i_1+i_2} 2^{-0.4k(i_1+i_2)} n^{o(1)} < (n^{-0.6+o(1)})^i < n^{-0.1}. \end{aligned}$$

This completes the proof of the claim.

Returning to the proof of the theorem, note that the variance of the random variable  $X$  satisfies

$$\text{Var}(X) \leq E(X) + \sum_{K, K'} \text{Cov}(X_{K_1, K_2}, X_{K'_1, K'_2})$$

where the summation is over all ordered pairs  $K = (K_1, K_2), K' = (K'_1, K'_2)$  with nontrivial intersections (since for all other pairs the covariance is zero). Since

$$\text{Cov}(X_{K_1, K_2}, X_{K'_1, K'_2}) \leq E(X_{K_1, K_2} X_{K'_1, K'_2})$$

it follows from the claim that

$$\text{Var}(X) \leq \mu + \Delta \leq (1 + o(1))\mu.$$

Therefore, by Chebyshev's Inequality, the probability that  $X \geq 3\mu/4$  is (much) bigger than  $3/4$ . By Markov's inequality, the probability that  $\Delta \leq \mu/4$  is also (much) bigger than  $3/4$  and hence with probability larger than  $1/2$ , both events happen simultaneously, that is, the number of copies of  $B$  in  $G$  is at least  $3\mu/4$  and the number of pairs of copies of  $B$  with nontrivial intersection is smaller than  $\mu/4$ . By removing one copy of  $B$  from each

pair with a nontrivial intersection we conclude that if this is the case, then  $G$  contains a family of at least  $\mu/2$  copies of  $B$  with no two having a nontrivial intersection.

Let  $h(G)$  be the maximum cardinality of a family of copies of  $B$  in  $G$  in which no two members have a nontrivial intersection, and let  $Y$  be the random variable  $Y = h(G)$ . We next apply Talagrand's Inequality (Theorem 2.2) to deduce that the probability that  $Y$  is zero is tiny.

By the above discussion, the probability that  $Y$  is at least  $\mu/2$  exceeds  $1/2$ . It is also clear that the value of  $Y = h(G)$  can change by at most 1 if we add or delete one edge to  $G$ , and that  $h$  is  $f$ -certifiable where  $f(s) = s \cdot k_1 k_2 \leq s k^2/4$ . Therefore, by Theorem 2.2 with  $b = \mu/2$  and  $t = \sqrt{2\mu}/k$  we conclude that the probability that  $Y = 0$  is smaller than

$$e^{-\mu/2k^2}.$$

This is much much smaller than one over the cardinality of  $\mathcal{B}'(k)$ , which is less than  $k2^{k^2/4}$ . As  $Y = 0$  if and only if there is no copy of  $B$  in  $G$ , we conclude that  $G$  is induced universal for  $\mathcal{B}'(k)$  with high probability. This completes the proof of Theorem 4.7.  $\square$

In order to complete the proof of Theorem 4.5 it suffices to show that there is bipartite graph with at most, say,  $2^{k/4}/k$  vertices which is induced universal for the symmetric bipartite graphs, that is, for all  $(k_1, k_2)$ -bipartite graphs with each  $k_i$  being at least  $0.45k$  that have induced subgraphs on at least  $0.9k$  vertices with at least  $k^{8m}$  automorphisms, where  $m = 2\sqrt{k \log k}$ . This is similar to the proof of Theorem 2.7. Here is a sketch of the argument. By Lemma 2.4 it follows that any such symmetric  $(k_1, k_2)$ -bipartite graph  $F = (U_1, U_2, E)$  has an automorphism with at least  $2m$  fixed points and at least  $6m$  non-fixed points. Such an automorphism must contain at least  $m$  fixed points in one vertex class and at least  $3m$  non-fixed points in the other. (Indeed, it has at least  $m$  fixed points in some vertex class and at least  $3m$  non-fixed points in some class. If these happen to be the same class, then the other class contains either many fixed points or many non-fixed points, implying what's needed.) Without loss of generality assume there are  $m$  fixed points in  $U_1$  and  $3m$  non-fixed points in  $U_2$ . As in Corollary 2.5 this implies that there is a set  $A = \{a_1, a_2, \dots, a_m\}$  of  $m$  vertices in  $U_1$  and two disjoint sets  $B = \{b_1, b_2, \dots, b_m\}$  and  $C = \{c_1, c_2, \dots, c_m\}$  in  $U_2$  such that for any  $1 \leq i, j \leq m$ ,  $a_i b_j$  is an edge of  $F$  if and only if  $a_i c_j$  is an edge of  $F$ .

The next ingredient needed is an appropriate bipartite analogue of Lemma 2.6, which follows.

**Lemma 4.9** *Let  $H$  be the bipartite  $(k_1, k_2)$ -graph obtained from the complete bipartite graph with vertex classes  $U_1, U_2$ , ( $|U_i| = k_i$ ), by omitting all edges connecting  $A$  and  $C$ ,*

where  $A$  is a subset of cardinality  $m$  of  $U_1$ , and  $B, C$  are disjoint subsets of cardinality  $m$  of  $U_2$ , and  $m = 2\sqrt{k \log k}$ . Then there is an orientation  $H'$  of  $H$  in which all edges between  $A$  and  $B$  are oriented from  $A$  to  $B$  and the maximum outdegree of a vertex in  $H'$  is at most

$$\frac{k_1 k_2}{k} - \frac{m^2}{k} + O(1) \left( < \frac{k}{4} - 3 \log k \right).$$

The proof is by applying Fact 2 used in the proof of Lemma 2.6. Let  $D_1 = U_1 - A$  and  $D_2 = U_2 - (B \cup C)$ . By Fact 2 for any real  $x \in (0, 1)$  there is an orientation of the edges of the complete bipartite graph with vertex classes  $A$  and  $D_2$  so that the outdegree of each vertex of  $A$  is at most  $x|D_2| + 1 = x(k_2 - 2m) + 1$  and the outdegree of each vertex of  $D_2$  is at most  $(1 - x)m + O(1)$ . Similarly, for any real  $y \in (0, 1)$  there is an orientation of the edges of the complete bipartite graph between  $B \cup C$  and  $D_1$  so that the outdegree of every vertex of  $B \cup C$  is at most  $y|D_1| + 1 = y(k_1 - m) + 1$  and the outdegree of each vertex of  $D_1$  is at most  $(1 - y)2m + O(1)$ . Finally, for any  $z \in (0, 1)$  there is an orientation of all edges connecting  $D_1$  and  $D_2$  so that each vertex of  $D_1$  has outdegree at most  $z(k_2 - 2m) + 1$  and each vertex of  $D_2$  has outdegree at most  $(1 - z)(k_1 - m) + O(1)$ .

Recalling that each  $k_i$  is between  $0.45k$  and  $0.55k$ , and that  $m = o(k)$ , it follows that there are  $x, y, z \in (0, 1)$ , all rather close to  $1/2$ , so that all 4 quantities

$$m + x(k_2 - 2m), \quad y(k_1 - m), \quad (1 - y)2m + z(k_2 - 2m), \quad \text{and} \quad (1 - x)m + (1 - z)(k_1 - m)$$

which represent (up to an  $O(1)$  additive error) the outdegrees of the vertices in  $A, B \cup C, D_1$  and  $D_2$  respectively, are equal. This means that in our orientation, in which all edges between  $A$  and  $B$  are oriented from  $A$  to  $B$ , the outdegrees of all vertices are essentially the same, and hence each of them is the average outdegree, which is

$$\frac{k_1 k_2}{k} - \frac{m^2}{k},$$

up to an additive constant error. This establishes the lemma.  $\square$

The last lemma can be used, as in the proof of theorem 2.7, to show that for any fixed  $k_1, k_2$  there is an adjacency labelling scheme for the above symmetric bipartite graphs in which each label has at most  $\frac{k}{4} - 2 \log k$  bits. This supplies an induced universal bipartite graph for the required symmetric bipartite graphs, whose total number of vertices is at most  $2^{k/4}/k$ . Together with Theorem 4.7, this completes the proof of Theorem 4.5.

## 5 Concluding remarks

- Our upper and lower bounds for the various induced-universal graphs considered here differ by lower order additive terms. It may be interesting to try and get even tighter estimates. How close are the counting lower bounds to the correct values ?

To be specific, consider the function  $f(k)$  discussed in Theorem 1.1, namely, the minimum possible number of vertices in an induced-universal graph for the set  $\mathcal{F}(k)$  of all  $k$ -vertex undirected graphs. The trivial counting lower bound shows that  $n = f(k)$  must satisfy

$$\binom{n}{k} \geq |\mathcal{F}(k)|.$$

Equality would hold here if there would have been a graph  $G$  on  $n = (1+o(1))2^{(k-1)/2}$  vertices containing every graph  $F \in \mathcal{F}(k)$  as an induced subgraph exactly once. It is not difficult to show that this is impossible, and in fact any graph  $G$  of this size must contain at least  $2^{\Omega(k)}$  induced copies of some  $k$ -vertex graphs. This is proved in the following proposition.

**Proposition 5.1** *For any positive constant  $c$  and all sufficiently large  $k$  the following holds. For any graph  $G = (V, E)$  on at least  $2^{ck}$  vertices there is a graph  $H$  on  $k$  vertices so that  $G$  contains at least  $2^{(c/2+o(1))k}$  induced copies of  $H$ .*

**Proof:** By the standard proof of Ramsey's Theorem there are vertices  $v_1, v_2, \dots, v_m$ , where  $m = (ck - \log k)/2$  and a set  $U$  of at least  $k$  vertices containing none of the vertices  $v_i$  so that one of the following holds. Either

- (i) There are no edges among the vertices  $v_i$  and no edges connecting any  $v_i$  to any vertex of  $U$ , or
- (ii) The vertices  $v_i$  form a clique and each  $v_i$  is adjacent to all vertices of  $U$ .

Indeed, starting with  $U_0 = V$  choose arbitrarily a vertex  $x_1 \in U_0$  and let  $U_1$  be either the set of all its neighbors in  $U_0$  or the set of all its non-neighbors, whichever is bigger. Next choose  $x_2 \in U_1$  and let  $U_2$  be the set of all its neighbors in  $U_1$  or all its non-neighbors, whichever is bigger. Proceeding in the same way, after  $2m - 1$  steps we get a set  $\{x_1, x_2, \dots, x_{2m-1}\}$  of vertices and a set  $U = U_{2m-1}$  of more than  $k$  vertices of  $G$ . If in at least  $m$  of the steps we have chosen non-neighbors, we get (i), else we get (ii).



To complete the proof fix a set  $U'$  of  $k - m/2$  vertices of  $U$  and observe that the induced graphs on the union of  $U'$  with any set of  $m/2$  of the vertices  $v_i$  are all isomorphic.  $\square$

In particular, if  $G$  is an induced universal graph for  $\mathcal{F}(k)$  with  $(1 + o(1))2^{(k-1)/2}$  vertices, whose existence is proved in Theorem 1.1, then by the above proposition  $G$  contains at least  $2^{(1+o(1))k/4}$  induced copies of some  $k$ -vertex graphs. (The proof in fact shows that this is the case for any subgraph consisting of  $m/2$  vertices  $v_i$  and any  $k - m/2$  vertices of  $U$ .) Note that counting shows that most  $k$ -vertex graphs appear much less, that is, only  $2^{o(k)}$  times, as induced subgraphs of  $G$ .

- It is not difficult to show that the maximum number of automorphisms of a tournament on  $k$  vertices is only exponential in  $k$  - much smaller than the maximum number of automorphisms of a  $k$ -vertex undirected graph (which is  $k!$ ) Indeed, let  $g(k)$  denote the maximum number of automorphisms of a tournament on  $k$  vertices, and let  $T$  be a tournament attaining this maximum. Like any tournament,  $T$  contains a vertex  $v$  whose outdegree  $d$  satisfies  $k/4 + O(1) \leq d \leq 3k/4 + O(1)$ . If an automorphism of  $T$  maps  $v$  to  $u$ , then its outneighbors are mapped to the outneighbors of  $u$  and its inneighbors are mapped to the inneighbors of  $u$ . This shows that  $g(k) \leq kg(d)g(k - 1 - d)$  where  $d$  is as above, and it is easy to verify that this gives an exponential upper bound for  $g(k)$ . This, together with the arguments in the proof of Theorem 2.1, imply that there is an absolute constant  $c > 1$  so that a random tournament with  $c2^{(k-1)/2}$  vertices is induced universal for the set  $\mathbf{T}_k$  of all tournaments on  $k$  vertices with high probability. Although this does not imply the sharp statement of Theorem 4.1, it shows that for tournaments, unlike for undirected graphs, the random construction is larger than the best one only by a (small) constant factor.
- The discussion in subsection 3.2 suggests that it may be interesting to study the vertex random graph process introduced there and in particular to investigate hitting time results for this process. Similar processes exist for random directed graphs, oriented graphs, tournaments, and even permutations, and appear to be interesting as well.
- All the induced universal graphs appearing in the theorems in the paper contain a large random part, namely, the constructions here are not explicit, unlike the ones in [25], [26], [4]. It would be interesting to find explicit constructions of the same sizes.

**Acknowledgment:** I would like to thank Michael Krivelevich, Alexander Magazinov, Oleg Pikhurko, Aner Shalev, Or Zamir and Uri Zwick for helpful discussions and comments.

## References

- [1] M. Ajtai, J. Komlós and E. Szemerédi, First occurrence of Hamilton cycles in random graphs, in: *Cycles in graphs* (Burnaby, B.C., 1982), 173–178, North-Holland Math. Stud., 115, North-Holland, Amsterdam, 1985.
- [2] N. Alon, Bipartite decomposition of random graphs, *J. Combinatorial Theory, Ser. B* 113 (2015), 220-235.
- [3] N. Alon and J. H. Spencer, *The Probabilistic Method, Fourth Edition*, Wiley, 2016, xv+375 pp.
- [4] S. Alstrup, H. Kaplan, M. Thorup and U. Zwick, Adjacency Labeling Schemes and Induced-Universal Graphs, *Proc. STOC 2015*, 625-634.
- [5] S. Alstrup, S. Dahlgaard and M. Knudsen, Optimal Induced Universal Graphs and Adjacency Labeling for Trees, *Proc. FOCS 2015*, 1311-1326.
- [6] L. Babai, On the orders of uniprimitive permutation groups, *Ann. of Math.* 113 (1981), 553-568.
- [7] L. Babai, Automorphism groups, isomorphism, reconstruction, in: *Handbook of Combinatorics*, Vol. 1, 2, Elsevier, Amsterdam, 1995, 1447-1540.
- [8] A. Bochert, Ueber die Transitivitätsgrenze der Substitutionengruppen, welche die alternirende ihres Grades nicht enthalten. (German) *Math. Ann.* 33 (1889), no. 4, 572–583.
- [9] B. Bollobás, The evolution of sparse graphs, *Graph theory and combinatorics* (Cambridge, 1983), 35–57, Academic Press, London, 1984.
- [10] B. Bollobás and P. Erdős, Cliques in random graphs, *Math. Proc. Cambridge Philos. Soc.* 80 (1976), no. 3, 419–427.
- [11] B. Bollobás and A. Thomason, Graphs which contain all small graphs, *European Journal of Combinatorics*, 2(1) (1981), 13–15.

- [12] B. Bollobás and A. Thomason, Random graphs of small order. *Random graphs '83* (Poznan 1983), 47–97, North-Holland Math. Stud., 118, North-Holland, Amsterdam, 1985.
- [13] G. R. Brightwell and Y. Kohayakawa, Ramsey properties of orientations of graphs, *Random Structures and Algorithms* 4 (1993), 413–428.
- [14] S. Butler, Induced-universal graphs for graphs with bounded maximum degree, *Graphs and Combinatorics* 25 (2009), 461–468.
- [15] F. R. K. Chung, Universal graphs and induced-universal graphs, *Journal of Graph Theory* 14 (1990), 443–454.
- [16] J. D. Dixon and B. Mortimer, *Permutation Groups*, Springer-Verlag, New York, 1996.
- [17] L. Esperet, A. Labourel and P. Ochem, On induced-universal graphs for the class of bounded-degree graphs, *Inf. Process. Lett.* 108 (2008), 255–260.
- [18] P. Erdős and A. Rényi, On the evolution of random graphs, *Publ. Math. Inst. Hungar. Acad. Sci.* 5 (1960), 17–61.
- [19] C. Gavaille and A. Labourel, Shorter implicit representation for planar graphs and bounded treewidth graphs, *Proc. ESA 2007*, pages 582–593, Springer, 2007.
- [20] S. Kannan, M. Naor and S. Rudich, Implicit representation of graphs, *SIAM J. Discrete Math.* 5 (1992), 596–603.
- [21] J. Kempe, L. Pyber and A. Shalev, Permutation groups, minimal degrees and quantum computing, *Groups Geom. Dyn.* 1 (2007), no. 4, 553–584.
- [22] J. Komlós and E. Szemerédi, Limit distributions for the existence of Hamilton circuits in a random graph, *Discrete Mathematics* 43 (1983), 55–63.
- [23] V. V. Lozin and G. Rudolf, Minimal universal bipartite graphs, *Ars Comb.* 84 (2007), 345–356.
- [24] D. W. Matula, *The largest clique size in a random graph*, Southern Methodist University, Dallas, Texas, 1976.
- [25] J. W. Moon, On minimal  $n$ -universal graphs, *Proceedings of the Glasgow Mathematical Association*, 7(1) (1965), 32–33.

- [26] J. W. Moon, *Topics on Tournaments*, New York, 1968.
- [27] C. Praeger and J. Saxl, On the order of primitive permutation groups, *Bull. London Math. Soc.* 12, (1980), 303-308.
- [28] R. Rado, Universal graphs and universal functions, *Acta. Arith.* (1964), 331–340.
- [29] V.G. Vizing, Some unsolved problems in graph theory, *Russian Mathematical Surveys*, 23(6) (1968), 125–141.
- [30] H. Wielandt, *Finite Permutation Groups*, Acad. Press, New York, 1964.