

Out-colourings of Digraphs

N. Alon* J. Bang-Jensen† S. Bessy‡

July 13, 2017

Abstract

We study vertex colourings of digraphs so that no out-neighbourhood is monochromatic and call such a colouring an **out-colouring**. The problem of deciding whether a given digraph has an out-colouring with only two colours (called a 2-out-colouring) is \mathcal{NP} -complete. We show that for every choice of positive integers r, k there exists a k -strong bipartite tournament which needs at least r colours in every out-colouring. Our main results are on tournaments and semicomplete digraphs. We prove that, except for the Paley tournament P_7 , every strong semicomplete digraph of minimum out-degree at least 3 has a 2-out-colouring. Furthermore, we show that every semicomplete digraph on at least 7 vertices has a 2-out-colouring if and only if it has a **balanced** such colouring, that is, the difference between the number of vertices that receive colour 1 and colour 2 is at most one. In the second half of the paper we consider the generalization of 2-out-colourings to vertex partitions (V_1, V_2) of a digraph D so that each of the three digraphs induced by respectively, the vertices of V_1 , the vertices of V_2 and all arcs between V_1 and V_2 have minimum out-degree k for a prescribed integer $k \geq 1$. Using probabilistic arguments we prove that there exists an absolute positive constant c so that every semicomplete digraph of minimum out-degree at least $2k + c\sqrt{k}$ has such a partition. This is tight up to the value of c .

1 Introduction

A **k -partition** of a (di)graph $G = (V, A)$ is a partition of V into k disjoint non-empty subsets V_1, \dots, V_k . The complexity of deciding whether one can partition the vertex set of a (di)graph into two or more non-empty subsets of vertices, such that the sub(di)graphs induced by these sets satisfies prescribed properties is a difficult problem. In [7, 9] the complexity of 120 such problems concerning 2-partitions of digraphs was settled.

Thomassen [18] proved that every digraph with minimum out-degree at least 3 has a 2-partition (V_1, V_2) such that each of the subdigraph induced by these two sets has minimum out-degree at least 1, see also [1] for an extension for partitioning into more parts. One can decide in polynomial time whether a given digraph has a 2-partition such that each set has minimum out-degree at least 1 (see e.g. [7]). Another result by Thomassen on digraphs with no even directed cycle [19] implies that there is no lower bound on the minimum out-degree of a digraph that implies that a digraph D has a 2-partition (V_1, V_2) such that the bipartite digraph $D(V_1, V_2)$ induced by the arcs between V_1, V_2 has minimum out-degree at least one. Still, as was shown in [6], it can be decided in polynomial time whether a given digraph has a 2-partition V_1, V_2 such that $D(V_1, V_2)$ has minimum out-degree at least one (if we require higher out-degrees, the problem becomes \mathcal{NP} -complete [5]).

In the language of 2-partitions of digraphs a 2-out-colouring is the same thing as a 2-partition (V_1, V_2) such that each vertex has an out-neighbour in both sets. That is, we want both of the properties above. This is the same thing as searching for a 2-colouring with no monochromatic edges of the hypergraph $\mathcal{H}_D = (V, \mathcal{E})$ on $|V|$ edges that we obtain from a given digraph $D = (V, A)$ by letting the edges in \mathcal{E} correspond to each of the $|V|$ out-neighbourhoods in D . It is well-known [14] that it is

*Sackler School of Mathematics and Blavatnik School of Computer Science, Tel Aviv University, Tel Aviv, Israel, Email: nogaa@tau.ac.il. Research supported in part by a BSF grant, an ISF grant and a GIF grant.

†Department of Mathematics and Computer Science, University of Southern Denmark, Odense DK-5230, Denmark, Email: jbj@imada.sdu.dk. Financial support: Danish research council, grant number 1323-00178B.

‡LIRMM, Université de Montpellier, Montpellier, France. Email: stephane.bessy@lirmm.fr. Financial support: OSMO project, Occitanie regional council.

\mathcal{NP} -complete to decide whether one can 2-colour the vertices of a hypergraph \mathcal{H} in such a way that no edge is monochromatic. As we show below this problem remains \mathcal{NP} -complete for hypergraphs like \mathcal{H}_D that are the **out-neighbourhood hypergraph** of some digraph D .

Our main focus is on tournaments and semicomplete digraphs, that is, respectively, orientations of complete graphs and digraphs with no pair of non-adjacent vertices. Tournaments form the most well-studied class of digraphs and despite their restricted structure a lot of deep results exist and many challenging problems remain for this class (see e.g. [10] for a comprehensive list of results on tournaments and semicomplete digraphs). In this paper we show, among other results, that with only one exception on 7 vertices, every semicomplete digraph with minimum out-degree at least 3 has a 2-out-colouring and give a polynomial algorithm for finding such a partition or certifying its nonexistence in an arbitrary semicomplete digraph. In the second part of the paper we consider a generalization of 2-out-colourings where we do not only want that each vertex has at least one out-neighbour in both sets but now we want at least k out-neighbours in each set for a prescribed number k . Using probabilistic arguments we prove that every tournament whose minimum out-degree is sufficiently large as a function of k has such a partition. The bound we give on this function is asymptotically best possible.

The paper is organized as follows. We first provide some notation and show that even for highly structured digraphs, such as symmetric digraphs and bipartite tournaments, high out-degree is not sufficient to guarantee the existence of an out-colouring with r colours for any fixed r . In Section 3 we show that deciding whether a digraph has a 2-out-colouring is \mathcal{NP} -complete, even when the input is restricted to be a symmetric digraph with high out-degree or to be semicomplete bipartite. In Section 4 we characterize all tournaments which have a 2-out-colouring. Using these results we characterize also in Section 5 all semicomplete digraphs admitting a 2-out-colouring. In Section 6 we prove that with very few exceptions, if a semicomplete digraph has a 2-out-colouring, then it also has a balanced one, that is, the sizes of the two sides of the 2-partition differ by at most one. In Section 7 we study the generalization of 2-out-colourings where we want a 2-partition such that each vertex has at least k out-neighbours in both sets. Using probabilistic methods we give asymptotically best possible bounds for the minimum out-degree which will guarantee that a tournament with this minimum out-degree has a 2-partition as above. Finally, in Section 8 we mention some further consequences of our results.

2 Terminology and preliminaries

Notation follows [8]. For any digraph $D = (V, E)$ we use the notation $u \rightarrow v$ if the arc uv belongs to E and we say that u **dominates** v and say that v is an **out-neighbour** of u and u is an **in-neighbour** of v . The set of out-neighbours (in-neighbours) of a vertex v is denoted by $N^+(v)$ ($N^-(v)$) and we denote by $\delta^+(D)$ the minimum out-degree of the digraph D , that is the minimum size of an out-neighbourhood. For two disjoint sets of vertices A and B of D we write $A \Rightarrow B$ if $a \rightarrow b$ for every $a \in A$ and $b \in B$. A set $X \subset V$ is an **in-dominating set** in $D = (V, A)$ if every vertex in $V - X$ dominates at least one vertex in X .

A (u, v) -path is a directed path from u to v . A digraph is **strongly connected** (or **strong**) if it contains a (u, v) -path for every ordered pair of distinct vertices u, v . A digraph D is **k -strong** if for every set S of less than k vertices the digraph $D - S$ is strong. A **strong component** of a digraph D is a maximal subdigraph of D which is strong. A strong component is **trivial**, if it has order 1. An **initial** (resp. **terminal**) strong component of D is a strong component X with no arcs entering (resp. leaving) X in D . It is easy to see that for a tournament T which is not strong, we can order the strong components uniquely as D_1, \dots, D_k , $k \geq 2$ so that all arcs are directed from D_i to D_j for $1 \leq i < j \leq k$. In particular T has exactly one initial component D_1 and exactly one terminal component D_k .

A **k -colouring** γ of a digraph $D = (V, E)$ is a mapping $\gamma : V \rightarrow \{1, \dots, k\}$ of its vertex set. A **k -colouring** γ of D is a **k -out-colouring** if no out-neighbourhood is monochromatic, that is for every vertex u of D there exist two out-neighbours v and w of u with $\gamma(u) \neq \gamma(v)$. For any k -colouring of D a vertex x will be **good** if its out-neighbourhood is not monochromatic. So a k -colouring is a k -out-colouring if all vertices of D are good.

For a 2-colouring $\gamma : V \rightarrow \{1, 2\}$ of a digraph D we will denote by V_i the set $\{x \in V : \gamma(x) = i\}$, for $i = 1, 2$. As the set V_1 totally defines γ we will often just specify it to describe the 2-colouring γ

and we will simply say that it **defines** γ .

The following observation shows that even for some classes of highly structured digraphs, for every integer r there does not exist a number K so that all digraphs from this class and with minimum out-degree at least K have an r -out-colouring. A **bipartite tournament** is a digraph that can be obtained from a complete bipartite graph B by assigning an orientation to each edge of B .

Proposition 2.1 *For all positive integers k, r there exists a k -strong bipartite tournament $B_{k,r}$ with $\delta^+(B_{k,r}) = k$ which has no r -out-colouring.*

Proof: Let U be an independent set on kr vertices, let $X_1, X_2, \dots, X_{\binom{kr}{k}}$ be an ordering of the distinct k -subsets of U and let V be an independent set of $\binom{kr}{k}$ vertices $v_1, v_2, \dots, v_{\binom{kr}{k}}$ where v_i corresponds to X_i for each $i \in [\binom{kr}{k}]$. The vertex set of $B_{k,r}$ is $U \cup V$ and the arc set consist of the arcs from v_i to X_i , $i \in [\binom{kr}{k}]$ and all other arcs between vertices in U and V go from U to V . It is easy to check that $B_{k,r}$ is k -strong. No matter how we r -colour U there will be a monochromatic k -set X_i , showing that $B_{k,r}$ has no r -out-colouring. \diamond

By adding a new set W inducing a digraph with chromatic number at least $p - 2$ and all possible arcs from U to W and from W to V , we see that there is no bound on the out-degree that guarantees a r -out-colouring, even when the digraph is p -chromatic.

We shall use the following classical result due to Moon.

Theorem 2.2 (Moon) [15] *Every strong tournament is vertex-pancyclic.*

3 Complexity of 2-out-colouring

Every undirected graph $G = (V, E)$ corresponds to the symmetric digraph $D = \overset{\leftrightarrow}{G}$ that we obtain from G by replacing every edge of G by a directed 2-cycle.

A **total dominating** set of a graph $G = (V, E)$ is a set of vertices S such that every vertex of G has a neighbour in S . Note that a graph G has a partition into two total dominating sets $(S, V - S)$ if and only if $\overset{\leftrightarrow}{G}$ has a 2-out-colouring. Thus the result of [13] that deciding whether a graph has a 2-partition into total dominating sets is \mathcal{NP} -complete, implies the following.

Theorem 3.1 *Deciding whether a given digraph has a 2-out-colouring is \mathcal{NP} -complete.*

For completeness we give a short proof of the following strengthening of Theorem 3.1

Theorem 3.2 *For every integer K , deciding whether a given symmetric digraph of minimum out-degree at least K , has a 2-out-colouring is \mathcal{NP} -complete.*

Proof: Let $\mathcal{H} = (X, E)$ be a hypergraph. We consider its bipartite representation G . That is, the vertices of G are $X \cup E$ and there is an edge from $x \in X$ to $e \in E$ iff $x \in e$. We add to G a vertex z linked to all vertices of X and then replace every edge by a 2-cycle to obtain the digraph $D_{\mathcal{H}}$. It is clear that $D_{\mathcal{H}}$ is a symmetric digraph, and we claim that \mathcal{H} is 2-colourable if and only if $D_{\mathcal{H}}$ admits a 2-out-colouring. Indeed if \mathcal{H} is 2-colourable then we keep this colouring on the set of vertices of $D_{\mathcal{H}}$ corresponding to X , colour the vertices of $D_{\mathcal{H}}$ corresponding to E by 1 and z by 2. Then we obtain a 2-out-colouring of $D_{\mathcal{H}}$. Conversely, if $D_{\mathcal{H}}$ has a 2-out-colouring, as the out-neighbours of every vertex belonging to E contains different colours, the colouring of X corresponds to a 2-colouring of the hypergraph \mathcal{H} .

The claim now follows from the \mathcal{NP} -completeness of the 2-coloring problem for hypergraphs [14], mentioned in the introduction, and the easy fact that the hypergraph 2-colouring problem remains \mathcal{NP} -complete when every vertex is in at least K hyperedges and all hyperedges have size at least K . \diamond

We show below that even for highly structured digraphs with many arcs, the problem is still hard.

Theorem 3.3 *It is NP-complete to decide whether a bipartite tournament with minimum out-degree 3 admits a 2-out-colouring.*

Proof: Let \mathcal{F} be an instance of MONOTONE NAE-3-SAT¹ with variables x_1, x_2, \dots, x_n and clauses C_1, C_2, \dots, C_m . By reordering if necessary, we may assume that C_1 and C_2 contain no common variable. Let $U = \{u_1, u_2, \dots, u_n\}$ and $V = \{c_1, c_1^*, c_1^{**}, c_2, c_2^*, c_2^{**}, c_3, \dots, c_{m-1}, c_m, \}$ be two disjoint vertex sets. For each clause $C_i, i \in [m]$ we add 3 arcs from c_i to the vertices $u_{i_1}, u_{i_2}, u_{i_3}$ corresponding to the three literals $x_{i_1}, x_{i_2}, x_{i_3}$ of C_i and let all remaining vertices in U have an arc to c_i . Finally we let the arcs incident to c_i^*, c_i^{**} be copies of the arcs incident to c_i for $i = 1, 2$ (so we have 3 vertices for each of C_1, C_2 and one for each other clause).

We claim that the resulting bipartite tournament $B = B(\mathcal{F})$ has a 2-out-colouring if and only there is a truth assignment ϕ to the variables such that every clause has either one or two true literals. Suppose first that B has a 2-out-colouring. Then we let variable x_i be true precisely if u_i receives colour 1 under this colouring. Since c_j has precisely 3 out-neighbours in B it follows that this truth assignment will satisfy one or two variables of each clause. Conversely, suppose ϕ is a truth assignment to the variables so that C_j has either one or two true literals for $j \in [m]$. For each $i \in [n]$ such that $\phi(x_i) = \text{true}$ we colour u_i by 1 and for all i such that $\phi(x_i) = \text{false}$ we colour u_i by 2. Now all clause vertices have both colours in their out-neighbourhoods, so we just have to make sure the same holds for the vertices $u_i, i \in [n]$. We obtain this by colouring all $c_j, j \in [m]$ as well as c_1^*, c_2^* by colour 1 and c_1^{**}, c_2^{**} by colour 2. Since every u_i either dominates both of c_1^*, c_1^{**} or both of c_2^*, c_2^{**} , every vertex of U is good also and this colouring is a 2-out-colouring of $B(\mathcal{F})$. \diamond

4 Out-colourings of tournaments

In this section we focus on 2-out-colourings of tournaments.

Let T be a tournament which is not strongly connected and denote by C the terminal component of T . As $T \setminus C \Rightarrow C$ the following holds.

Observation 1 *If γ is a k -out-colouring of the terminal component C of a tournament T , then every extension of γ to the vertices of $T \setminus C$ leads to a k -out-colouring of T .*

Lemma 4.1 *Every tournament T on $n \geq 5$ vertices contains either an in-dominating vertex or an in-dominating cycle of size at most $n - 2$.*

Proof: First assume that T is not strongly connected and denote by X the terminal component of T . If X only contains one vertex, then this vertex is an in-dominating vertex of T . Otherwise if $|X| \geq 4$ then by Theorem 2.2, T contains a cycle on $|X| - 1$ vertices which is an in-dominating cycle of T of size at most $n - 2$. And if $|X| = 3$ then a Hamiltonian 3-cycle of X forms also an in-dominating cycle of T of size at most $n - 2$.

Now if T is strongly connected, by Theorem 2.2 it contains a cycle C on $n - 2$ vertices. Let us denote by x and y the vertices of $T \setminus C$. If x and y have both an out-neighbour on C , then C is an in-dominating cycle of T . Otherwise, as T is strongly connected, it means for instance that y has an out-neighbour on C and that we have $x \rightarrow y$ and $C \Rightarrow x$. In this case if we let z be an out-neighbour of y on C , xyz is an in-dominating cycle of T of size $3 \leq n - 2$. \diamond

4.1 2-out-colouring of tournaments with minimum out-degree 2

The **rotational tournament on 5 vertices** denoted by RT_5 has vertex set $\{1, 2, 3, 4, 5\}$ and is the union of the two directed cycles 12345 and 13524. It is easy to check that RT_5 has no 2-out-colouring. Moreover we define the tournament T_7 on 7 vertices. It contains two 3-cycles C and C' and a vertex

¹Recall that NAE-3-SAT is the variant of 3-SAT where we seek a truth assignment t such that each clause has both a false and a true literal under t . The further restriction monotone NAE-3-SAT means that we restrict to instances with no negated variables. This problem is still \mathcal{NP} -complete [16].

z such that $C' \Rightarrow z$, $z \Rightarrow C$ and $C \Rightarrow C'$. It is also easy to check that T_7 has no 2-out-colouring. A tournament T on $n \geq 6$ vertices with $\delta^+(T) = 2$ belongs to **the family** \mathcal{G}_1 if there exists a subtournament T' of T on $n - 3$ vertices such that: T' has an in-dominating vertex w , $T \setminus T'$ is a 3-cycle C , $C \Rightarrow z$ where z is a vertex of T' different from w and with out-degree at least 2, and $T' \setminus z \Rightarrow C$. Notice that if T belongs to \mathcal{G}_1 then T has no 2-out-colouring. Indeed in such a colouring all the vertices of C must receive the same colour (different from the colour of z) and then the out-neighbourhood of w would be monochromatic. The tournaments RT_5 , T_7 and the family \mathcal{G}_1 are depicted in Figure 1 (for the more general case of semicomplete digraphs).

Theorem 4.2 *A tournament T with $\delta^+(T) = 2$ admits a 2-out-colouring except if it belongs to the family \mathcal{G}_1 or if its terminal strong component is RT_5 or T_7 .*

Proof: By Observation 1 we can assume that T is strongly connected. We deal with two different cases.

Case 1: T has exactly one vertex with out-degree 2. In this case T admits a 2-out-colouring. Indeed denote by a the vertex of T with out-degree 2 and by b and c its out-neighbours with $b \rightarrow c$. Let also c_1 and c_2 be two out-neighbours of c . Consider the 2-colouring defined by $\{a, c, c_1\}$. As every vertex different from a has degree at least 3, it is a 2-out-colouring of T except if there exists a vertex d_1 whose out-neighbourhood is exactly $\{a, c, c_1\}$. Now the 2-colouring defined by $\{a, c, c_2\}$ is a 2-out-colouring of T : If this was not the case, then there would be a new vertex d_2 with $N^+(d_2) = \{a, c, c_2\}$, but then d_1 dominates d_2 , contradicting our conclusion above.

Case 2: T contains at least two vertices of degree 2. We observe two different sub-cases.

Case 2.1: No two vertices of out-degree 2 have a common out-neighbour. In this case, denote by a and b two vertices of out-degree 2 with $a \rightarrow b$. Denote by c the second out-neighbour of a and by d and e the two out-neighbours of b with $d \rightarrow e$. Notice that we have $c \rightarrow b$ and $\{d, e\} \Rightarrow a$. First consider the 2-colouring of T defined by $\{a, b, e\}$. It is a 2-out-colouring of T except if there exists a vertex x with its out-neighbours included in $\{a, b, e\}$. In this case the out-neighbourhood of x is exactly $\{a, b, e\}$ otherwise x has out-degree 2 and has a common out-neighbour with a or b which is excluded in this sub-case. In particular we have $x \in N^-(a) \cup N^-(b)$ and every vertex different from a , b and e dominates x . So now the 2-colouring with first set $\{a, b, d\}$ is a 2-out-colouring of T .

Case 2.2: There exist two vertices of out-degree 2 with a common out-neighbour. Call by a and b two such vertices with $a \rightarrow b$ and denote by d their common out-neighbour and by c the second out-neighbour of b . Also denote by X the set $N^-(a) \cap N^-(b) = V(T) \setminus \{a, b, c, d\}$. First assume that $c \rightarrow d$. If c has an out-neighbour e in X and $d \rightarrow e$ then $\{b, c, e\}$ defines a 2-out-colouring of T . If d does not dominate e then $\{b, c, e, f\}$ defines a 2-out-colouring of T where f is an out-neighbour of d . So we can assume that $X \Rightarrow \{a, b, c\}$ and that c has out-degree 2 also. If there is $w \in X$ whose out-neighbourhood is exactly $\{a, b, c\}$ then T belongs to the family \mathcal{G}_1 , with $T' = T[X \cup \{d\}]$, $C = abc$ and d playing the role of z . Otherwise let e_3 be an out-neighbourhood of d and consider the 2-colouring defined by $\{a, b, c, e_3\}$. If it is not a 2-out-colouring of T , it means that there exist e_2 whose out-neighbourhood is exactly $\{a, b, c, e_3\}$ (it cannot be just $\{a, b, c\}$ otherwise T belongs to \mathcal{G}_1). Similarly if $\{a, b, c, e_2\}$ does not define a 2-out-colouring of T there exists e_1 whose out-neighbourhood is $\{a, b, c, e_2\}$. Now either $\{a, b, c, e_1\}$ defines a 2-out-colouring of T or $T[a, b, c, d, e_1, e_2, e_3]$ is the terminal component of T and isomorphic to T_7 . In this later case T does not admit a 2-out-colouring. Finally, we treat the case where $d \rightarrow c$. Assume that there exist e and f two different vertices in X such that $c \rightarrow e$ and $d \rightarrow f$. Then $\{b, c, e\}$ defines a 2-out-colouring of T . Thus we may assume that c and d are vertices of out-degree 2 and they have a common out-neighbour e in X . If $X \setminus \{e\} = \emptyset$ then $T = RT_5$, otherwise $X \setminus \{e\} \Rightarrow \{a, b, c, d\}$ and as T is strongly connected e must have an out-neighbour in X . So we check that $\{a, b, c\}$ defines a 2-out-colouring of T . \diamond

4.2 2-out-colouring of tournaments with minimum out-degree at least 3

Lemma 4.3 *Every tournament T with $\delta^+(T) \geq 3$ and which has an in-dominating set of size 2 admits a 2-out-colouring*

Proof: Assume that $\{a, b\}$ is an in-dominating set of T with a dominating b . Let c be an out-neighbour of a . We consider the 2-colouring of T defined by $\{a, b, c\}$. If it is not a 2-out-colouring of

T it means that there exists a vertex d whose out-neighbourhood is exactly $\{a, b, c\}$. So we choose another out-neighbour c' of b and check that $\{a, b, c'\}$ defines a 2-out-colouring of T (as every vertex of $T \setminus \{a, b, c, d\}$ dominates either a or b and dominates d). \diamond

The **Paley tournament on 7 vertices** denoted by P_7 has vertex set $\{1, 2, 3, 4, 5, 6, 7\}$ and is the union of the three directed cycles 1234567, 1357246 and 1526374. It is easy to check that P_7 has no 2-out-colouring.

Theorem 4.4 *Every tournament T with $\delta^+(T) \geq 3$ and whose terminal strong component is different from P_7 admits a 2-out-colouring.*

Proof: By Observation 1 we can assume that T is strongly connected. Consider a vertex x of T with minimum out-degree. If $T[N^+(x)]$ has an in-dominating vertex y , then $\{x, y\}$ is an in-dominating set of T of size 2 and we conclude with Lemma 4.3. We study now different cases:

First assume that $\delta^+(T) \geq 5$. In particular $d^+(x) \geq 5$ and by Lemma 4.1 $T[N^+(x)]$ contains an in-dominating cycle C of size at most $d^+(x) - 2$. So $C \cup x$ defines a 2-out-colouring of T . Indeed it is clear that every vertex has an out-neighbour coloured by 1. And if a vertex y has only out-neighbours coloured by 1, it means that $d^+(y) \leq |C| + 1 \leq d^+(x) - 2 + 1 < d^+(x)$, a contradiction.

Now assume that $\delta^+(T) = 4$. We have $d^+(x) = 4$ and as $T[N^+(x)]$ does not contain any in-dominating vertex, $T[N^+(x)]$ contains an in-dominating 3-cycle. So we denote $N^+(x) = \{a, b, c, d\}$ with abc being a 3-cycle and $d \rightarrow a$. We consider the 2-colouring of T defined by $\{x, a, b, c\}$. If it is not a 2-out-colouring it means that there exists $y \in N^-(x)$ such that $N^+(y) = \{x, a, b, c\}$. In this case let e be an out-neighbour of b lying in $N^-(x)$ and consider the 2-colouring of T defined by $\{x, a, b, e\}$. As every vertex of $N^-(x) \setminus y$ dominates y , this colouring is a 2-out-colouring of T .

Finally we assume that $\delta^+(T) = 3$. We have $d^+(x) = 3$ and as $T[N^+(x)]$ does not contain any in-dominating vertex, $T[N^+(x)]$ is 3-cycle, denoted by abc . By Lemma 4.1, we may assume that $N^+(a) \cap N^+(b) \neq \emptyset$, $N^+(b) \cap N^+(c) \neq \emptyset$, $N^+(c) \cap N^+(a) \neq \emptyset$. If a, b and c have a common out-neighbour y in $N^-(x)$ then consider the 2-out-colouring of T defined by $\{x, a, y\}$. If it is not a 2-out-colouring of T , then $\{x, b, y\}$ defined a 2-out-colouring of T . Similarly if a and b have two common out-neighbours y and y' , then $\{x, a, y\}$ or $\{x, a, y'\}$ defines a 2-out-colouring of T . Thus by symmetry, it means that a and b have exactly one common out-neighbour, called d , that b and c have exactly one common out-neighbour, called f and that c and a have exactly one common out-neighbour, called e . Using that every vertex of $N^-(x) \setminus d$ dominates a or b , we can deduce the following. If $d \rightarrow f$ then $\{x, c, d, e\}$ defines a 2-out-colouring of T . So by symmetry we can assume that def is a 3-cycle of T . If $\{a, b, c\}$ has an out-neighbour in $N^-(x) \setminus \{d, e, f\}$, say that $c \rightarrow y$ with $y \in N^-(x) \setminus \{d, e, f\}$, then $\{x, c, d, y\}$ defines a 2-out-colouring of T . Finally if $N^-(x) \setminus \{d, e, f\} \neq \emptyset$ then as T is strongly connected it means that for instance $d \rightarrow y$ for some $y \in N^-(x) \setminus \{d, e, f\}$. But then $\{x, c, d, e\}$ defines a 2-out-colouring of T . Otherwise $N^-(x) \setminus \{d, e, f\} = \emptyset$ and T is exactly the Paley tournament. \diamond

Together Theorem 4.2, Theorem 4.4 and the fact that our proofs are constructive provide the following corollary.

Corollary 4.5 *A tournament T with $\delta^+(T) \geq 2$ admits a 2-out-colouring if and only if its terminal strong component is different from RT_5 , T_7 and P_7 and T does not belong to the family \mathcal{G}_1 . Consequently, in polynomial time, one can decide whether or not a tournament admits a 2-out-colouring and find a 2-out-colouring when it exists.*

5 Out-colourings of semicomplete digraphs

We extend the results of the previous section to semicomplete digraphs.

5.1 2-out-colourings of semicomplete digraphs with minimum out-degree 2

Figure 1 defines a family \mathcal{G} of semicomplete digraphs. A digraph belongs to \mathcal{G} if its terminal strong component is CD_3 , RT_5 , RT_5^a , RT_5^b , RT_5^{bb} or RT_5^c or if it belongs to \mathcal{G}_1 or \mathcal{G}_2 . A digraph belongs to \mathcal{G}_1 if it contains a vertex w whose out-neighbourhood is a 2- or 3-cycle C such that the out-neighbourhood of $V(C)$ is exactly one vertex z different from w . Finally a semicomplete digraph T belongs to the family \mathcal{G}_2 if its terminal strong component contains a 2- or 3-cycle C which dominates a 2- or 3-cycle C' which dominates a vertex z dominating C .




Figure 1: The semicomplete digraphs CD_3 , RT_5 , RT_5^a , RT_5^b , RT_5^{bb} and RT_5^c and the families \mathcal{G}_1 and \mathcal{G}_2 (red arcs could be replaced by 2-cycles and non oriented red edges could be oriented in any direction).

Lemma 5.1 *None of the semicomplete digraphs in \mathcal{G} admit a 2-out-colouring. Moreover for each semicomplete digraph $T \in \mathcal{G}$ if we add an arc to T , then either the resulting digraph admits a 2-out-colouring or it belongs to \mathcal{G} .*

Proof: We will not provide the complete proof of the statement but just give some important remarks on how to obtain the result. We leave it to the reader to check that if we add an extra arc to one of the six digraphs in $\mathcal{G} - \mathcal{G}_1 - \mathcal{G}_2$, then we either get a new digraph in \mathcal{G} or the resulting digraph has a 2-out-colouring. However we pay attention to the family \mathcal{G}_1 where the most technical case occurs. Let T be a digraph of \mathcal{G}_1 and xy be an arc to add to T . If xy is incident neither to C nor to w then $T + xy$ is clearly a member of \mathcal{G}_1 . The same holds if $x = z$ and $y \in C$. If $x \in C$ or $xy = wz$ then we check that $T + xy$ admits a 2-out-colouring. Finally let us assume that $x = w$ and that y belongs to $X = V(T) \setminus (V(C) \cup \{w, z\})$. If z has a second out-neighbour in X different from y then $\{w, y, z\}$ defines a 2-out-colouring of $T + xy$. So assume now that y is the only out-neighbour of z in X . If y has an out-neighbour t in X then $\{z, y, t\}$ defines a 2-out-colouring of $T + xy$. In the remaining case $V(C) \cup \{w, y, z\}$ forms the terminal strong component of $T + xy$ and contains the 2-cycle wy which dominates C which dominates z which in turn dominates the 2-cycle wy . Then $T + xy$ belongs to \mathcal{G}_2 . Finally let us just mention that adding an arc to any digraph of \mathcal{G}_2 leads to a digraph admitting a 2-out-colouring. \diamond

Theorem 5.2 *A semicomplete digraph T with $\delta^+(T) = 2$ admits a 2-out-colouring unless its terminal strong component is CD_3 , RT_5 , RT_5^a , RT_5^b , RT_5^{bb} or RT_5^c or if it belongs to \mathcal{G}_1 or \mathcal{G}_2 .*

Proof: We prove the result by induction on the number of 2-cycles in T . If T does not contain any 2-cycle then the result holds by Theorem 4.2. Assume now that T has a 2-cycle xy and suppose first that the digraph $T - xy$ still satisfies $\delta^+(T - xy) = 2$. If $T - xy$ admits a 2-out-colouring then

so does T . Otherwise by induction it means that $T - xy$ belongs to the family \mathcal{G} . By Lemma 5.1 T also belongs to \mathcal{G} or admits a 2-out-colouring.

Now by symmetry we can assume that both x and y have out-degree exactly 2 in T . We look at two sub-cases: either x and y have common out-neighbour or not. Assume first that x and y both dominate a vertex z . If there exists $w \in V(T)$ such that $N_T^+(w) = \{x, y\}$, then either $w = z$ and the terminal component of T is CD_3 or $w \neq z$ and T belongs to the family \mathcal{G}_1 . In both cases T does not admit a 2-out-colouring. Otherwise it means that $\delta^+(T \setminus \{x, y\}) \geq 1$. If z dominates x or y then $\{x, y\}$ defines a 2-out-colouring of T . If z has an out-neighbour u such that $\delta^+(T \setminus \{x, y, u\}) \geq 1$ then $\{x, y, u\}$ defines a 2-out-colouring of T . Otherwise it is easy to check that the out-neighbourhood of z is exactly a 2- or a 3-cycle C dominated by $V(T) \setminus (V(C) \cup \{x, y\})$. But in this case T belongs to \mathcal{G}_2 . For the other sub-case, assume that x and y do not have a common out-neighbour. So we respectively denote by z and t the second out-neighbour of x and y . In particular, z dominates y and t dominates x . Without loss of generality we can assume that z dominates t . If $\delta^+(T \setminus \{x, y\}) \geq 1$, then $\{x, y\}$ defines a 2-out-colouring of T . Otherwise it means that there exists $v \in T$ with $N^+(v) = \{x, y\}$. Notice that $v \neq z$ as $z \rightarrow t$. We treat together the cases $v = t$ and $v \neq t$. If z has an out-neighbour u in $V(T) \setminus \{x, y, t, v\}$ then $\{x, z, u\}$ defines a 2-out-colouring of T . Otherwise the out-neighbourhood of z is exactly $\{v, t, y\}$ and T belongs to the family \mathcal{G}_1 with tv being the 2- or 3-cycle C and x playing the role of w . \diamond

5.2 2-out-colourings of semicomplete digraphs with minimum out-degree at least 3

Lemma 5.3 *Every semicomplete digraph T with $\delta^+(T) \geq 3$ which admits an in-dominating set of size 2 has a 2-out-colouring.*

Proof: Let $\{a, b\}$ be an in-dominating set of T . If $T[a, b]$ is a 2-cycle, then $\{a, b\}$ defines a 2-out-colouring of T , as $\delta^+(T) \geq 3$. Otherwise assume that a dominates b and that b does not dominate a . Let c be an out-neighbour of b . We know that c dominates a or b and so if $\{a, b, c\}$ does not define a 2-out-colouring of T , it means that there exists $d \in V(T)$ such that $N^+(d) = \{a, b, c\}$. As a is not an out-neighbour of b , there exists a vertex $c' \notin \{a, b, c, d\}$ such that $b \rightarrow c'$. As previously if $\{a, b, c'\}$ does not define a 2-out-colouring of T there exists $d' \in V(T)$ with $N^+(d') = \{a, b, c'\}$. As d' does not dominate d we must have $d' = c$. Now if $b \rightarrow d$ then bd is a in-dominating 2-cycle of T , and we conclude as in the first part of the proof. Otherwise there exists $c'' \notin \{a, b, c, d, c'\}$ such that $b \rightarrow c''$. To conclude we check that $\{a, b, c''\}$ defines a 2-out-colouring of T . \diamond

Theorem 5.4 *Every semicomplete digraph T with $\delta^+(T) \geq 3$ admits a 2-out-colouring unless its terminal strong component is P_7 .*

Proof: We prove the result by induction on the number of 2-cycles in T . If T does not contain any 2-cycle then the result holds by Theorem 4.4. Besides it is easy to show the Paley tournament on 7 vertices plus one arc admits a 2-out-colouring. Indeed without loss of generality, this semicomplete digraph has vertex set $\{1, \dots, 7\}$ and arc set $\{ij : j - i = 1, 2 \text{ or } 4 \pmod{7}\} \cup \{21\}$ and admits the 2-out-colouring defined by $\{3, 4, 6\}$. So assume that T contains a 2-cycle ab . When we remove from T the arc ab if we obtain a semicomplete digraph with minimum out-degree at least 3 (including P_7) then T admits a 2-out-colouring by induction or by the previous remark. Otherwise it means that a has out-degree exactly 3. The same holds for b and we denote by S the set $(N^+(a) \cup N^+(b)) \setminus \{a, b\}$. Thus we have $2 \leq |S| \leq 4$. If we have $|S| = 4$ then S is an in-dominating set of T of size 2 and we conclude with Lemma 5.3. If $|S| = 2$ then a and b have two common out-neighbours c and d . Assume that c dominates d , then $\{a, d\}$ is an in-dominating set of T of size 2 and we conclude again with Lemma 5.3. Finally if $|S| = 3$ then a and b has a common out-neighbour c and we denote by d the third out-neighbour of a and by e the one of b . As $|S| = 3$ we have $d \neq e$. If d and e both dominate c , then $\{a, c\}$ is an in-dominating set of T and we use Lemma 5.3. Otherwise we can assume without loss of generality that c dominates d . As a does not dominate e , e has to dominate a . Once again $\{a, d\}$

is an in-dominating set of T and we use Lemma 5.3 to conclude. \diamond

As the proofs of Theorem 5.2 and Theorem 5.4 are constructive provide we obtain the following corollary.

Corollary 5.5 *A semicomplete digraph T with $\delta^+(T) \geq 2$ admits a 2-out-colouring unless its terminal strong component is CD_3 , RT_5 , RT_5^a , RT_5^b , RT_5^{bb} , RT_5^c or P_7 or if it belongs to \mathcal{G}_1 or \mathcal{G}_2 . Consequently, in polynomial time, one can decide whether or not a semicomplete digraph admits a 2-out-colouring and find a 2-out-colouring when it exists.*

To conclude this section, notice that every exception listed in the previous statement can be 3-out-coloured. So we obtain the following.

Corollary 5.6 *Every semicomplete digraph T with $\delta^+(T) \geq 2$ admits a 3-out-colouring.*

6 Balanced 2-out-colourings of semicomplete digraphs

A 2-out-colouring with colour classes V_1 and V_2 is **balanced** if we have $||V_1| - |V_2|| \leq 1$.

The next figure defines three families of semicomplete digraphs \mathcal{T}_1 , \mathcal{T}_2 and \mathcal{T}_3 containing respectively two, three and five digraphs. It is not difficult to check that all of the digraphs in \mathcal{T}_1 , \mathcal{T}_2 and \mathcal{T}_3 have a 2-out-colouring (the one indicated by the two columns of vertices) and that none of them have a balanced 2-out-colouring.




Figure 2: The families \mathcal{T}_1 , \mathcal{T}_2 and \mathcal{T}_3 of semicomplete digraphs which admit a 2-out-colouring but no balanced 2-out-colouring (red arcs could be replaced by 2-cycles and non oriented red edge could be oriented in any direction).

Theorem 6.1 *Every semicomplete digraph T which admits a 2-out-colouring also admits a balanced 2-out-colouring except if T belongs to \mathcal{T}_1 , \mathcal{T}_2 or \mathcal{T}_3 .*

Proof: Let T be a semicomplete digraph which admits a 2-out-colouring. If $|T| \leq 5$ then any 2-out-colouring of T is balanced. So we assume that $|T| \geq 6$ and among all the 2-out-colourings of T we consider one, denoted by γ , inducing a partition (V_1, V_2) with $||V_1| - |V_2||$ as small as possible. Assume that $||V_1| - |V_2|| > 1$ and that we have $|V_2| > |V_1| + 1$. So, let Y_2 be the set $\{v_2 \in V_2 : d_{V_2}^+(v_2) = 1\}$, let X_2 be $N_{V_2}^+(Y_2)$ and for every vertex $v_2 \in V_2$ let Z_{v_2} be the set $\{v_1 \in V_1 : N_{V_2}^+(v_1) = \{v_2\}\}$. Notice that as T is semicomplete, we have $|X_2| = |Y_2| \leq 3$ and that if a vertex $v_2 \in V_2 \setminus X_2$ satisfies $Z_{v_2} = \emptyset$ then $V_1 \cup \{v_2\}$ also defines a 2-out-colouring of T , contradicting the choice of γ . So we have $Z_{v_2} \neq \emptyset$ for every $v_2 \in V_2 \setminus X_2$. Moreover, by definition, all the sets Z_{v_2} are disjoint and we have $|V_1| \geq \sum_{v_2 \in V_2 \setminus X_2} |Z_{v_2}| \geq |V_2 \setminus X_2| = |V_2| - |X_2|$. So we have $|X_2| = 2$ or $|X_2| = 3$.

Below we will show that we can either easily reach a contradiction to the optimality of γ or we may choose one vertex x in V_1 and two vertices y and z in V_2 such that the 2-colouring γ' defined by $(V_1 \setminus \{x\}) \cup \{y, z\}$ is a 2-out-colouring of T , contradicting also the choice of γ . In order to define γ' we

will also select another vertex x' in V_1 . If V_1 contains a vertex with out-degree exactly 1 in V_1 then we choose x' to be such a vertex and x the only out-neighbour of x' in V_1 . Otherwise we choose x' to be any vertex of V_1 and x one out-neighbour of x' in V_1 . In any case, every vertex of $V_1 \setminus \{x'\}$ has at least one out-neighbour in $V_1 \setminus \{x\}$. Then we consider several cases.

First assume that $|Y_2| = 2$ and that $T[Y_2]$ is 2-cycle ab . In this case we have $X_2 = Y_2$ and $V_2 \setminus X_2$ dominates X_2 . Thus we also have $|V_2| = |V_1| + 2$ and $|Z_{v_2}| = 1$ for every vertex $v_2 \in V_2 \setminus X_2$ and then $V_1 = \cup_{v_2 \in V_2 \setminus X_2} Z_{v_2}$. Notice that the arcs from V_1 to V_2 form a matching. So we choose $y \in V_2$ such that $Z_y = \{x'\}$ and $z = a$. Let us check that in this case γ' is 2-out-colouring of T . Every vertex of $V_2 \setminus X_2$ dominates X_2 and hence dominates a vertex from $V_1(\gamma')$ and one from $V_2(\gamma')$. The same holds for the vertices of X_2 because they dominate x' and x . Every vertex of $V_1 \setminus x'$ has still one out-neighbour in $V_1 \setminus \{x\} \subset V_1(\gamma')$ and one in $V_2 - \{y\} \subset V_2(\gamma')$. And finally x' dominates x which lies in $V_2(\gamma')$ and y which lies in $V_1(\gamma')$. So γ' is a 2-out-colouring of T , a contradiction.

Next assume that $|Y_2| = 2$ also but that $T[Y_2]$ is not a 2-cycle. In this case we must also have $|V_2| = |V_1| + 2$ and $|Z_{v_2}| = 1$ for every vertex $v_2 \in V_2 \setminus X_2$ and then $V_1 = \cup_{v_2 \in V_2 \setminus X_2} Z_{v_2}$. We denote by a and b the vertices of Y_2 with $a \rightarrow b$ and we denote by c the only out-neighbour of b , that is, $X_2 = \{b, c\}$. Notice that every vertex of $V_2 \setminus \{a, b, c\}$ dominates $\{a, b\}$. If $|V_1| \geq 3$ then every vertex of V_2 has at least two out-neighbours in V_1 and we can choose $y \in V_2$ such that $Z_y = \{x'\}$ and $z = a$. As previously, we check that γ' is 2 out-colouring of T , using especially that the only arcs from V_1 to V_2 is the matching $\{uv : Z_v = \{u\}, v \in V_2 \setminus X_2\}$ and so that every vertex of V_2 has at least one out-neighbour in $V_1(c')$. So we must have $|V_1| = 2$ and then V_1 is a 2-cycle de with $d \rightarrow a$ for instance. Denote by f the vertex of V_2 with $Z_f = \{e\}$. So T must contain the following arcs: fa , fb , ca and all the arcs from V_2 to V_1 except possibly ad and fe . If T has a 2-out-colouring such that d and e receive the same colour, then a , b , c and f are forced to receive the other colour and the 2-out-colouring is not balanced. If d and e have different colours in a balanced 2-out-colouring, then this 2-out-colouring must have bipartition $(\{a, b, d\}, \{c, e, f\})$. Every vertex dominates a vertex in each part except possibly f . But if f does not dominate c or e then T belongs to the family \mathcal{T}_1 .

Finally, assume that $|Y_2| = 3$. In this case Y_2 is a 3-cycle, denoted by abc , and $Y_2 = X_2$ is dominated by $V_2 \setminus X_2$. If $|V_2 \setminus X_2| = |V_1|$, then as previously, we have $V_1 = \cup_{v_2 \in V_2 \setminus X_2} Z_{v_2}$. We choose x , x' and y as before and let $z = a$, then we check that the resulting colouring γ' is a balanced 2-out-colouring of T . Thus we must have $|V_1| = |V_2 \setminus X_2| + 1$. In this case we have $|Z_{v_2}| = 1$ for every vertex $v_2 \in V_2$ except possibly for one vertex v_2 where we can have $|Z_{v_2}| = 2$. So we pick one vertex in each Z_{v_2} and denote by w the only vertex of V_1 not chosen. Now we denote by Y_1 the set $\{v_1 \in V_1 : d_{V_1}^+(v_1) = 1\}$, and by X_1 the set $N_{V_1}^+(Y_1)$. If there exists $x_1 \neq w$ in $V_1 \setminus X_1$, it means that every vertex in V_1 has an out-neighbour in $V_1 \setminus x_1$. Then we choose y and z in $X_2 = \{a, b, c\}$ such that $N_{V_2}^+(w) \not\subseteq \{y, z\}$. Now it is easy to check that the 2-colouring defined by $(V_1 \setminus \{x_1\}) \cup \{y, z\}$ is a 2-out-colouring of T . So we must have that $V_1 \subseteq X_1 \cup \{w\}$. In particular, as $|X_1| = |Y_1| \leq 3$, we have $|V_1| \leq 4$. We will first look at two particular cases. In **case A** we assume that $T[X_1]$ is a 2-cycle or a 3-cycle and that w dominates all vertices of X_1 . Then, we choose $x' \in X_1$, x the out-neighbour of x' in $T[X_1]$, y the vertex of V_2 such that $x' \in Z_y$ and $z \in X_2$. It is easy to check that $(V_1 \setminus \{x\}) \cup \{y, z\}$ defines a 2-out-colouring of T . In **case B** we assume that $T[V_1]$ has size 3 and contains a 3-cycle wx_1y_1 . We denote by x_2 (resp. y_2) the only out-neighbour of x_1 (resp. y_1) in V_2 . If $N_{V_2}^+(w) = \{x_2\}$ then we check that $(V_1 \setminus \{w\}) \cup \{y_2, a\}$ defines a 2-out-colouring of T . If $N_{V_2}^+(w) \neq \{x_2\}$ then we can choose $z \in \{a, b, c\}$ such that $N_{V_2}^+(w) \cap (V_2 \setminus \{x_2, z\}) \neq \emptyset$. So we check that $(V_1 \setminus \{y_1\}) \cup \{x_2, z\}$ defines a 2-out-colouring of T .

Now, if $|V_1| = 4$ then $X_1 = Y_1 = V_1 \setminus w$ and $T[V_1]$ is a 3-cycle dominated by w , situation we treated in *case A*. If $|V_1| = 3$ then either $|X_1| = 2$ or $|X_1| = 3$. In the former case, if $Y_1 = X_1$ then $T[V_1 \setminus w]$ is a 2-cycle dominated by w , corresponding to *case A*. If $|X_1| = 2$ and $Y_1 \neq X_1$ then $T[V_1]$ contains a 3-cycle, case we already settled in *case B*. And if $|V_1| = |X_1| = 3$ then $T[V_1]$ is exactly a 3-cycle, situation corresponding to *case B*. Finally if $|V_1| = 2$ then $T[V_1]$ is a 2-cycle. We denote by wd this 2-cycle and by e the only out-neighbour of d in V_2 . If w has at least one out-neighbour and at least one in-neighbour in X_2 , then we can assume that w dominates a and is dominated by b . So T has a balanced 2-out-colouring defined by $\{a, b, w\}$, contradicting the choice of γ . Otherwise X_2 dominates w or is dominated by w and there are no digons between w and X_2 . In these case we can see that T belongs to the family \mathcal{T}_2 or \mathcal{T}_3 . \diamond

From the proof of the above theorem we can derive the following corollary.

Corollary 6.2 *There exists a polynomial algorithm which given a semicomplete digraph $T \notin \mathcal{T}_i$, $i \in [3]$ and a 2-out-colouring of T , returns a balanced 2-out-colouring of T*

7 2-partitions of tournaments with out-degree at least k to both sets

In this section, for a given 2-partition (V_1, V_2) we let $D\langle V_i \rangle$ denote the induced subgraph of D on V_i and let $D\langle V_1, V_2 \rangle$ denote the spanning bipartite subgraph whose edges are all edges of D with an end vertex in V_1 and an end vertex in V_2 . Recall that we call a 2-partition (V_1, V_2) balanced if $||V_1| - |V_2|| \leq 1$.

Our aim is to use probabilistic methods to obtain results about sufficient conditions, in terms of minimum out-degrees, for the existence of 2-partitions where all vertices have at least k out-neighbours in both sets. Our main result is the following

Theorem 7.1 *There exist two absolute positive constants c_1, c_2 so that the following holds.*

1. *Let $T = (V, E)$ be a tournament with minimum out-degree at least $2k + c_1\sqrt{k}$. Then there is a balanced partition $V = V_1 \cup V_2$ of V so that $\delta^+(T\langle V_1 \rangle), \delta^+(T\langle V_2 \rangle)$ and $\delta^+(T\langle V_1, V_2 \rangle)$ are all at least k .*
2. *For infinitely many values of k there is a tournament with minimum out-degree at least $2k + c_2\sqrt{k}$ so that for any partition $V = V_1 \cup V_2$ of V into two disjoint sets, at least one of the quantities $\delta^+(T\langle V_1 \rangle), \delta^+(T\langle V_2 \rangle), \delta^+(T\langle V_1, V_2 \rangle)$ is smaller than k .*

In order to illustrate the usefulness of the probabilistic method, we first give short probabilistic proof of the following corollary of Theorems 4.4 and 6.1. Recall that the constant 4 is best possible by the Paley tournament P_7 .

Corollary 7.2 *Let $T = (V, E)$ be a tournament with minimum out-degree at least 4. Then there is a balanced partition $V = V_1 \cup V_2$ of V so that $\delta^+(T\langle V_1 \rangle), \delta^+(T\langle V_2 \rangle)$ and $\delta^+(T\langle V_1, V_2 \rangle)$ are all at least 1.*

7.1 Proofs

We need the following simple statement.

Lemma 7.3 *Let $(n_i)_{i \geq i_0}$ be a sequence of integers, put $N_j = \sum_{i=i_0}^j n_i$ and let $(p_i)_{i \geq i_0}$ be a non-increasing sequence of non-negative reals. Suppose $N_j \leq B_j$ for all $j \geq i_0$. Then the sum*

$$S = \sum_{i \geq i_0} n_i p_i$$

satisfies

$$S \leq B_{i_0} p_{i_0} + \sum_{i \geq i_0} (B_{i+1} - B_i) p_{i+1}.$$

Proof: Let S be as above, then

$$\begin{aligned} S &= N_{i_0} p_{i_0} + (N_{i_0+1} - N_{i_0}) p_{i_0+1} + (N_{i_0+2} - N_{i_0+1}) p_{i_0+2} + \dots \\ &= N_{i_0} (p_{i_0} - p_{i_0+1}) + N_{i_0+1} (p_{i_0+1} - p_{i_0+2}) + \dots \\ &\leq B_{i_0} (p_{i_0} - p_{i_0+1}) + B_{i_0+1} (p_{i_0+1} - p_{i_0+2}) + \dots = B_{i_0} p_{i_0} + \sum_{i \geq i_0} (B_{i+1} - B_i) p_{i+1}, \end{aligned}$$

as needed. ◇

Proof of Corollary 7.2: Let n_i denote the number of vertices of T with out-degree i , and put $N_j = \sum_{i \leq j} n_i$. Thus $N_3 = 0$. Note that $N_i \leq 2i + 1$ for all i , as the average out-degree in any induced subgraph of T on more than $2i + 1$ vertices exceeds i . We consider two possible cases.

Case 1: $n_4 \geq 2$. Let x, y be two vertices of out-degree 4 in T and let M be an arbitrary near perfect matching in T containing the edge connecting x and y and two additional edges x_1, x_2 and y_1, y_2 where x_1, x_2 are out-neighbors of x and y_1, y_2 are out-neighbors of y (it is easy to check that there are always three disjoint edges as above). For each edge cd of M , randomly and independently, place either c in V_1 and d in V_2 or c in V_2 and d in V_1 , where each of these choices are equally likely. If $|V|$ is odd place the remaining vertex, uncovered by M , randomly and uniformly either in V_1 or in V_2 . Note that by construction V_1 and V_2 are of nearly equal sizes. In addition both x and y have an out-neighbor in V_1 and an out-neighbor in V_2 . Moreover, each of the vertices that is neither an out-neighbor of x nor of y has both x and y as out-neighbors, and hence has at least one out-neighbor in V_1 and at least one in V_2 . The only remaining vertices are the out-neighbors of x and y (besides x and y). There are at most 7 such vertices. For each such vertex v let A_v denote the event that v fails to have an out-neighbor in V_1 or in V_2 . If the out-neighbors of v contain an edge of M , the probability of this event is 0. Else, its probability is exactly $2^{1-d^+(v)} \leq 1/8$, where $d^+(v)$ is the out-degree of v . Thus, by the union bound, the probability that some event A_v holds is at most $7/8 < 1$, showing that with positive probability no event A_v holds, that is, there is a partition with the desired properties in this case.

Case 2: $n_4 \leq 1$. Put $p_i = 2^{1-i}$ and let M be a near perfect matching as before, including an edge connecting two out-neighbors of the unique vertex of out-degree 4, if there is such a vertex. Let V_1 and V_2 be chosen randomly, as before, by the random process splitting the endpoints of each edge of M randomly and independently between V_1 and V_2 . Thus V_1 and V_2 are of nearly equal sizes and if there is a vertex of degree 4 then it has out-neighbors in V_1 and in V_2 . It remains to deal with the other vertices. For each vertex v let A_v be the event defined in Case 1. We have to show that with positive probability no event A_v holds. Using the fact that the sequence p_i is decreasing it follows, by Lemma 7.3, that the sum of probabilities of all these events is at most

$$11p_5 + \sum_{i \geq 5} ((2i + 3) - (2i + 1))p_{i+1} = 11 \cdot 2^{-4} + \sum_{i \geq 5} 2^{1-i} = 13/16 < 1.$$

This proves the existence of the desired partition.

We proceed with the proof of the second part of Theorem 7.1. For a prime q which is 3 modulo 4, the quadratic residue tournament P_q is the tournament whose vertices are the integers modulo q where (i, j) is a directed edge iff $i - j$ is a quadratic residue modulo q .

Lemma 7.4 *Let $P_q = (V, E)$ be as above. Then for any function $f : V \mapsto \{-1, 1\}$ there is a vertex $v \in V$ so that $|\sum_{u \in N^+(v)} f(u)| > \frac{1}{2}\sqrt{q}$.*

Proof: It is easy and well known (c.f., e.g., [3], Chapter 9) that every vertex of P_q has out-degree and in-degree $(q - 1)/2$ and any two vertices of it have exactly $(q - 3)/4$ common in-neighbors (and out-neighbors). Let $A = A_q$ be the adjacency matrix of P_q , that is, the 0/1 matrix whose rows and columns are indexed by the vertices of P_q , where $A_{ij} = 1$ iff (i, j) is a directed edge. By the above comment each diagonal entry of $A^t A$ is $(q - 1)/2$ and each other entry is $(q - 3)/4$. Thus the eigenvalues of $A^t A$ are $(q - 1)/2 + (q - 1)(q - 3)/4 = (q - 1)^2/4$ (with multiplicity 1) and $(q - 1)/2 - (q - 3)/4 = (q + 1)/4$ (with multiplicity $(q - 1)$). This implies that

$$\|Af\|_2^2 = f^t A^t A f \geq (q + 1)/4 \|f\|_2^2 = q(q + 1)/4.$$

It follows that there is an entry of Af whose square is at least $(q + 1)/4$, completing the proof. \diamond

Note that by the above Lemma, for any partition of the vertices of P_q into two disjoint (not necessarily nearly equal) sets V_1 and V_2 there is a vertex v of P_q so that the number of its out-neighbors in V_1 differs from that in V_2 by more than $\sqrt{q}/4$. This implies the assertion of part (ii) of Theorem 7.1 for infinitely many values of k .

Before proving the assertion of Theorem 7.1 we describe a short proof of the following weaker result.

Proposition 7.5 *Let $T = (V, E)$ be a tournament with minimum out-degree at least*

$$2k + (1 + o(1))\sqrt{2k \ln k}.$$

Then there is a balanced partition $V = V_1 \cup V_2$ of V so that $\delta^+(T\langle V_1 \rangle)$, $\delta^+(T\langle V_2 \rangle)$ and $\delta^+(T\langle V_1, V_2 \rangle)$ are all at least k . (The $o(1)$ -term above tends to zero as k tends to infinity.)

Proof: The proof is similar to that of Theorem 7.2. We assume, whenever this is needed, that k is sufficiently large. Put $m = 2k + (1 + \epsilon)\sqrt{2k \ln k}$. Let n_i denote the number of vertices of T with out-degree i , and put $N_j = \sum_{i \leq j} n_i$. Thus $N_s = 0$ for all $s < m$ and $N_s \leq 2s + 1$ for all s . Let M be an arbitrary near perfect matching in T . For each edge cd of M , randomly and independently, place either c in V_1 and d in V_2 or c in V_2 and d in V_1 , where each of these choices are equally likely. If $|V|$ is odd place the remaining vertex, uncovered by M , randomly and uniformly either in V_1 or in V_2 . Note that by construction V_1 and V_2 are of nearly equal sizes. For each vertex v of T let A_v be the event that v has less than k out-neighbors in V_1 or less than k out-neighbors in V_2 . Let $d(\geq m)$ be the out-degree of v . Note that A_v is exactly the event that the number of out-neighbors of v in V_1 differs from that in V_2 by more than $d - 2k$. Let X_v be the random variable whose value is this difference. If the set of out-neighbors of v contains no edge of the matching M then X_v is the sum of d independent uniform $-1, 1$ variables. By the Chernoff Inequality (c.f., e.g., [3], Theorem A.1.2) the probability of the event A_v is at most $2e^{-(d-2k)^2/2d}$. If the set of out-neighbors contains t edges of the matching then X_v is the sum of only $d - 2t$ independent uniform $-1, 1$ variables and the probability is even smaller. Put $p_d = 2e^{-(d-2k)^2/2d}$. It is easy to check that the sequence p_d is decreasing for all $d \geq 2k$ (indeed the function $g(x) = -(x - 2k)^2/2x$ is decreasing on the interval $[2k, +\infty[$). Therefore, by Lemma 7.3 and the union bound, the probability that at least one of the events A_v holds is at most

$$P \leq 2 \cdot (2m + 1)e^{-(m-2k)^2/2m} + 2 \sum_{d > m} p_d.$$

In the sum above each of the terms p_d for $d \leq 4k$ is at most p_m , and each of the terms p_d for bigger d is at most $2e^{-d/8}$. Therefore

$$P \leq (4m + 2 + 2 \cdot 4k)e^{-(m-2k)^2/2m} + 2 \sum_{d > 4k} 2e^{-d/8} < (1 + o(1))16ke^{-(m-2k)^2/2m} + 40e^{-4k/8} < 1,$$

where in the last inequality we have used the assumption that k is large. It follows that with positive probability none of the events A_v holds, completing the proof. \diamond

We proceed with the proof of part (i) of Theorem 7.1. An equivalent formulation of this part is that for any tournament $T = (V, A)$ with minimum out-degree at least $2k + c_1\sqrt{k}$ there is a function $f : V \mapsto \{-1, 1\}$ so that $|\sum_{v \in V} f(v)| \leq 1$ and for every vertex v of T with out-degree $d = |N^+(v)|$

$$\left| \sum_{u \in N^+(v)} f(u) \right| \leq d - 2k.$$

Indeed, given such an f we can simply define $V_1 = f^{-1}(1)$ and $V_2 = f^{-1}(-1)$. This resembles results about discrepancy of set systems (see, e.g., [3], Chapter 13). In particular, for the special case in which the number of vertices of the tournament is at most, say, $10k$, the above follows from the six standard deviations result of Spencer [17], which asserts that the discrepancy of any hypergraph with m vertices and m edges is at most $6\sqrt{m}$. The general case requires some work, we prove it by combining a variant of the partial coloring idea of [17] (see also [11]) with the main result of [17]. In what follows we make no attempt to optimize the absolute constants.

Lemma 7.6 *Let \mathcal{F} be a family of subsets of $[n] = \{1, 2, \dots, n\}$, and suppose that each set $F \in \mathcal{F}$ is of size at least $2k + 1000\sqrt{k}$. Suppose, further, that for every s there are less than $3s$ members of \mathcal{F} of size at most s . Then there is a function $f : [n] \mapsto \{-1, 0, 1\}$ such that*

1. $|\sum_{i=1}^n f(i)| \leq 1$
2. For every $F \in \mathcal{F}$ satisfying $|F| \leq 100k$, $|\sum_{i \in F} f(i)| \leq 200\sqrt{k}$
3. For every $F \in \mathcal{F}$ of size $|F| > 100k$, $|f^{-1}(1) \cap F| \geq k$ and $|f^{-1}(-1) \cap F| \geq k$.

Proof : Without loss of generality assume that n is even (otherwise add a point). The proof can be described using the pigeonhole principle, but it is cleaner to present a version applying some simple properties of the entropy function. Recall that the (binary) entropy of a random variable X getting values x_i with probabilities p_i for $i \in I$ is

$$H_2(X) = \sum_{i \in I} p_i \log_2(1/p_i).$$

It is well known (see, e.g., [3], Chapter 15) that if $X = (X_1, X_2, \dots, X_q)$ is a vector then

$$H_2(X) \leq \sum_{i=1}^q H_2(X_i) \tag{1}$$

Another fact we need is that

$$\text{If } p_i \leq 2^{-r} \text{ for all } i \in I \text{ then } H_2(X) \geq \sum_{i \in I} p_i \cdot r = r \tag{2}$$

Let $g : [n] \mapsto \{-1, 1\}$ be a random function defined as follows. For each $i \leq n/2$ randomly, uniformly and independently put $g(2i-1) = 1, g(2i) = -1$ or $g(2i-1) = -1, g(2i) = 1$. Put $\mathcal{F}_1 = \{F \in \mathcal{F}, |F| \leq 100k\}$. For each $F \in \mathcal{F}_1$ define

$$t(F) = \lfloor 0.5 + \frac{\sum_{i \in F} g(i)}{20\sqrt{|F|}} \rfloor$$

Note that for each such F , $t(F)$ is very likely to be 0. It is 1 or -1 only with probability smaller than e^{-50} , and more generally it is i or $-i$ with probability smaller than e^{-50i^2} for all $i \geq 1$. Therefore, the entropy of the random variable $t(F)$ is smaller than

$$\log_2 \frac{1}{1 - 2e^{-50}} + \sum_{i \geq 1} 2 \cdot \log_2 e \cdot (50i^2)e^{-50i^2}$$

which is much smaller than $1/30000$. Let X be the (vector valued) random variable defined by $X = (t(F) : F \in \mathcal{F}_1)$. By (1), since the total number of members of \mathcal{F}_1 is smaller than $300k$, the entropy of the vector X is smaller than $0.01k$. It follows by (2) that there is a specific value of the vector X obtained with probability at least $2^{-0.01k}$. Fix such a vector X . In what follows we show that we can choose a pair g_1, g_2 of functions g as above that give this vector so that $f = (g_1 - g_2)/2$ satisfies the assertion of the lemma.

We need the following claim.

Claim 1 *Let F be a fixed member of $\mathcal{F} - \mathcal{F}_1$. Then the number of pairs of functions $g_1, g_2 : [n] \mapsto \{-1, 1\}$ defined in the previous paragraph so that there are less than k elements j of F with $g_1(j) = 1 = -g_2(j)$ or that there are less than k elements j of F with $g_1(j) = -1 = -g_2(j)$ is smaller than $2^{n-0.1|F|}$.*

Proof of Claim 1: Let s be the number of indices i so that $\{2i-1, 2i\} \subset F$. Then there are $|F| - 2s$ elements of F whose mate in the matching $\{2i-1, 2i\}$, ($i \leq n/2$) is not in F . Thus there are exactly $2^{s+|F|-2s} = 2^{|F|-s}$ ways to choose the values of $g_1(j)$ for all $j \in F$ and the same number of ways to choose the values of $g_2(j)$ for $j \in F$. We next bound the number of choices in which there are less than k elements j of F satisfying $g_1(j) = 1 = -g_2(j)$. This number is at most

$$M = \sum_{j+\ell < k} \binom{s}{j} 2^s \binom{|F|-2s}{\ell} 3^{|F|-2s-\ell} < \binom{|F|}{k} 2^s 3^{|F|-2s}. \tag{3}$$

Indeed, for each j, ℓ with $j + \ell < k$ there are $\binom{s}{j}$ ways to choose j pairs $\{2i - 1, 2i\}$ contained in F in which g_1, g_2 do not agree. Once these are chosen there are still 2^s ways to choose the actual values of g_1, g_2 on these s pairs (agreeing on j pairs and disagreeing on $s - j$). There are then $\binom{|F| - 2s}{\ell}$ ways to select the elements $j \in F$ that do not belong to these pairs for which $g_1(j) = 1 = -g_2(j)$. Finally there are 3 possibilities for the values of $g_1(j)$ and $g_2(j)$ for each other element j .

Since $|F| \geq 100k$ it follows that $\binom{|F|}{k} \leq 2^{H_2(0.01)|F|} < 2^{0.1|F|}$. Here we used the known fact that for every pair of integers $a > b > 0$, $\binom{a}{b} \leq 2^{H_2(b/a)a}$ where

$$H_2(x) = x \log_2 \frac{1}{x} + (1 - x) \log_2 \frac{1}{1 - x}$$

is the binary entropy of the number x , $0 < x < 1$, which is the binary entropy of the indicator random variable attaining the value 1 with probability x and the value 0 with probability $1 - x$. The above inequality follows, for example, from the assertion of Corollary 15.7.3 in [3] by taking $n = a$ with \mathcal{F} being the family of all subsets of cardinality b of $\{1, 2, \dots, a\}$.

Plugging in (3) we conclude that

$$M \leq 2^{0.1|F|} 2^s 3^{|F| - 2s} < 2^{0.1|F|} 4^{|F| - s} (3/4)^{|F|} < 2^{-0.3|F|} 4^{|F| - s} < 2^{-0.11|F|} 4^{|F| - s}$$

Each choice of the values of g_1, g_2 on the $|F| - s$ pairs of elements $\{2i - 1, 2i\}$ intersecting F can be completed to the full values of g_1, g_2 in $4^{n/2 - |F| + s}$ ways. Thus the number of pairs g_1, g_2 for which there are less than k elements j with $g_1(j) = 1 = -g_2(j)$ is smaller than

$$2^{-0.11|F|} 4^{|F| - s} \cdot 4^{n/2 - |F| + s} = 2^{n - 0.11|F|}$$

By symmetry, the number of pairs g_1, g_2 so that there are less than k elements j with $g_1(j) = -1 = -g_2(j)$ satisfies the same inequality. This provides the assertion of Claim 1. \diamond

Returning to the proof of Lemma 7.6 recall that we have fixed a value of X such that there are at least $2^{n/2 - 0.01k}$ choices for the function g giving the value of X . Hence there are at least $2^{n - 0.02k}$ choices for an ordered pair of functions g_1, g_2 giving this value. By Claim 1, among these choices, the number of pairs that have, in some set $F \in \mathcal{F} - \mathcal{F}_1$, less than k elements j with $g_1(j) = 1 = -g_2(j)$ or less than k elements j with $g_2(j) = 1 = -g_1(j)$ is at most

$$\sum_{F \in \mathcal{F} - \mathcal{F}_1} 2^{n - 0.1|F|} < \sum_{s \geq 100k} (3s) 2^{n - 0.1s} < 2^{n - 0.02k}$$

(with a lot of room to spare).

In the above inequality we used the fact that for $A = \sum_{s=r}^{\infty} s q^s$, $qA = \sum_{s=r}^{\infty} s q^{s+1}$, and thus

$$A - qA = r q^r + \sum_{s=r+1}^{\infty} q^s = r q^r + \frac{q^{r+1}}{1 - q}$$

So we obtain

$$A = \frac{r q^r}{1 - q} + \frac{q^{r+1}}{(1 - q)^2}$$

Taking $q = 2^{-0.1}$ and $r = 100k$ gives the required bound.

Since there are $2^{n - 0.02k}$ choices for the ordered pair of functions g_1, g_2 , each giving the value of X , it follows that there is a pair g_1, g_2 for which the event described in Claim 1 does not happen for any $F \in \mathcal{F} - \mathcal{F}_1$. Fix such g_1, g_2 and define $f = (g_1 - g_2)/2$. We claim that f satisfies the assertion of Lemma 7.6. Indeed, by construction it satisfies property 1. Property 2 follows from the fact that both g_1 and g_2 give the same vector X implying that for every $F \in \mathcal{F}_1$

$$\lfloor 0.5 + \frac{\sum_{i \in F} g_1(i)}{10\sqrt{|F|}} \rfloor = \lfloor 0.5 + \frac{\sum_{i \in F} g_2(i)}{10\sqrt{|F|}} \rfloor$$

Property 3 follows from the fixed choice of the functions g_1, g_2 . This completes the proof of the lemma. \diamond

Proof of Theorem 7.1, part (i): Let $[n] = \{1, 2, \dots, n\}$ denote the set of vertices of T and let \mathcal{F} be the family of out-neighborhoods of its vertices. Note that it satisfies the assumptions of Lemma 7.6. Let f be as in the conclusion of the lemma. Put all elements in $f^{-1}(1)$ in V_1 and all elements in $f^{-1}(-1)$ in V_2 (the partition of the elements in $f^{-1}(0)$ will be determined in what follows.) Note that the partial assignment above already ensures, by property 3, that all vertices with out-degree exceeding $100k$ have at least k out-neighbors in V_1 as well as in V_2 .

It remains to assign the vertices i for which $f(i) = 0$ values in $\{-1, 1\}$ ensuring that we do not increase the discrepancy of the members of \mathcal{F}_1 by too much and maintaining the property $|\sum f(i)| \leq 1$. We then define $V_1 = f^{-1}(1), V_2 = f^{-1}(-1)$. This is done by applying the result of [17] that implies that the discrepancy of any set system of m sets (of any sizes) is at most $12\sqrt{m}$, (see [3], Corollary 13.3.4). Apply that to the family consisting of the intersections of the sets in \mathcal{F}_1 with $f^{-1}(0)$ together with one additional set: the set $f^{-1}(0)$. Adding this set ensures that our resulting partition is nearly balanced, and we can now change the values of some $O(\sqrt{k})$ elements (among those in $f^{-1}(0)$) arbitrarily, to make the split precisely balanced without changing the discrepancy of any set F in \mathcal{F}_1 by more than $2 \cdot 12\sqrt{300k + 1}$ (an addition of at most $12\sqrt{300k + 1}$ corresponding the discrepancy of $F \cap f^{-1}(0)$ and another addition of at most this quantity corresponding to the final arbitrary modification of values that makes the split balanced). This completes the proof. \diamond

7.2 Consequences of our methods

We conclude this section with some observations about further consequences of the methods and proofs we used.

- The proofs carry over with no change to semicomplete digraphs.
- The same proof works to split a tournament or a semicomplete digraph to r parts with each vertex having at least k out-neighbors in each part. Minimum out-degree $(1 + o(1))rk$ suffices. Getting best possible bounds for small r and k , e.g., $r \leq 10$ and $k = 1$, may be difficult.
- The same proof works to split an oriented graph with no independent set of size s . Here any set of size exceeding $(s - 1)(2r + 1)$ contains a vertex of out-degree exceeding r , hence we have an upper bound on the number of vertices of out-degree at most r and can repeat the probabilistic argument to get a similar result.
- The probabilistic proof, with no real essential change, can handle simultaneously in-degrees and out-degrees, establishing the following.

Theorem 7.7 *There exists an absolute constant c so that the following holds for every positive integer k . Let $T = (V, A)$ be a tournament with minimum out-degree at least $2k + c\sqrt{k}$ and minimum in-degree at least $2k + c\sqrt{k}$. Then there is a balanced partition $V = V_1 \cup V_2$ of V so that every vertex has at least k out-neighbors in V_1 and in V_2 , and every vertex has at least k in-neighbors in V_1 and in V_2 .*

8 Further remarks

We have shown in Section 7, using probabilistic arguments, that every tournament whose minimum out-degree is sufficiently large as a function of k has a partition of the vertices into two sets of nearly equal sizes so that each vertex has at least k out-neighbours in each set. The bound we give on this function is asymptotically best possible. After the completion of this paper we learned from the authors of [20] that they have independently proved the existence of such a function. Their proof is also

probabilistic, but the bound they get for the function is weaker, roughly twice our bound.

A digraph is k -out-regular if all out-degrees are k . As mentioned in the introduction Thomassen [19] showed that for every integer $k \geq 1$ there exist k -out-regular digraphs with no even cycle. As mentioned in [2] this implies that for every k there is an oriented graph with minimum out-degree at least k which admits no splitting into two parts with every vertex having an out-neighbor in the other part. On the other hand, as shown here and as mentioned above, such splittings always exist for tournaments and for graphs with bounded independence numbers, provided the minimum out-degree is large enough. It may be interesting to give additional natural classes of oriented graphs for which such a splitting is possible.

Theorem 8.1 *Every planar digraph D with $\delta^+(D) \geq 2$ admits a 4-out-colouring.*

Proof: By removing arcs we can assume that D is 2-out-regular. Now we define the non-oriented graph G_D on $V(D)$ with edge set $\{N^+(x) : x \in V(D)\}$. The graph G_D is planar otherwise it would contain a K_5 or $K_{3,3}$ minor by Kuratowski's Theorem (see e.g. [12]). But the underlying graph of D would also contain such a minor contradicting the fact that D is planar. So, using the Four Colour Theorem [4], we obtain a proper 4-colouring of G_D which correspond to a 4-out-colouring of D . \diamond

Notice that a planar digraph D such that G_D contains K_4 for instance achieves the bound given by Theorem 8.1.

Proposition 8.2 *There exists a polynomial algorithm for deciding whether a 2-out-regular digraph has a 2-out-colouring.*

Proof: This follows from the fact that a 2-out-regular digraph D has a 2-out-colouring if and only if the graph G_D defined above is bipartite. This can be checked in linear time. \diamond

As mentioned in the introduction, there is a polynomial algorithm for deciding whether a digraph D has a 2-partition (V_1, V_2) such that each of $D\langle V_i \rangle$, $i = 1, 2$ have out-degree at least one. It was shown in [6] that, despite Thomassen's examples mentioned above, it is also polynomial to decide whether D has a 2-partition so that $D\langle V_1, V_2 \rangle$ has minimum out-degree at least 1. So, given that 2-out-colouring is \mathcal{NP} -complete, it is natural to ask about the complexity of deciding whether D has a 2-partition such that $D\langle V_1, V_2 \rangle$ and $D\langle V_1 \rangle$ both have out-degree at least 1, but we do not require this for $D\langle V_2 \rangle$. The following result shows that this is also \mathcal{NP} -complete.

Theorem 8.3 *It is \mathcal{NP} -complete to decide whether a given digraph D has a 2-partition (V_1, V_2) so that $D\langle V_1, V_2 \rangle$ and $D\langle V_1 \rangle$ both have minimum out-degree at least one.*

Proof: Call a 2-partition (V_1, V_2) **nice** if $D\langle V_1, V_2 \rangle$ and $D\langle V_1 \rangle$ both have minimum out-degree at least one. Let X denote the digraph on 6 vertices $\{a, b, v, v', \bar{v}, \bar{v}'\}$ and the following 12 arcs $\{ab, ba, av', b\bar{v}', vv', v'v, \bar{v}\bar{v}', \bar{v}'\bar{v}, v\bar{v}, \bar{v}v', v'\bar{v}', \bar{v}'v\}$. It is not difficult to check that X has a nice 2-partition and that for every such partition (V_1, V_2) we have $v, v' \in V_1$ and $\bar{v}, \bar{v}' \in V_{3-i}$ for $i = 1$ or $i = 2$ and both are possible. Now let \mathcal{F} be an instance of NAE-3-SAT with variables x_1, x_2, \dots, x_n and clauses C_1, C_2, \dots, C_m . Form a digraph $R = R(\mathcal{F})$ as follows: take n copies X_1, X_2, \dots, X_n of X and give the vertices in the i th copy subscript i , i.e. in X_i the vertex v_i corresponds to the vertex v . Associate the vertices v_i, \bar{v}_i of X_i with the variable x_i and its negation \bar{x}_i , respectively. Now for each clause C_j we add two new vertices d_j, c_j , the arc $d_j c_j$ and three arcs from c_j to those vertices in the X copies that correspond to the literals of C_j . Hence if $C_j = (x_4 \vee \bar{x}_6 \vee x_8)$, then we add the three arcs $c_j v_4, c_j \bar{v}_6$ and $c_j v_8$. This completes the description of R . As the vertices d_j , $j \in [m]$ have out-degree one, they must all belong to V_2 in any nice 2-partition and this forces all the vertices c_1, c_2, \dots, c_m to belong to V_1 for all nice 2-partitions. Assume now that R has a nice 2-partition (V_1, V_2) . By the remark above, each vertex c_j belongs to V_1 and hence must have an out-neighbour in both sets and it is easy to check that if we set x_i to be true whenever $v_i \in V_1$ and false otherwise, we obtain a truth assignment satisfying one or two literals of every clause. Conversely, given a truth assignment

ϕ satisfying at least one but never three literals of any clause, we obtain a nice 2-partition by putting $\{d_1, \dots, d_m\}$ in V_2 , $\{c_1, c_2, \dots, c_m\}$ in V_1 , for each $i \in [n]$ putting $v_i \in V_1$ and $\bar{v}_i \in V_2$ if $\phi(x_i) = 'True'$ and $v_i \in V_2$ and $\bar{v}_i \in V_1$ if $\phi(x_i) = 'False'$ and finally distributing the rest of the vertices of the X copies in V_1, V_2 (as we know we can). \diamond

References

- [1] N. Alon. Disjoint directed cycles. *J. Combin. Theory Ser. B*, 68(2):167–178, 1996.
- [2] N. Alon. Splitting digraphs. *Combin. Probab. Comput.*, 15:933–937, 2006.
- [3] N. Alon and J.H. Spencer. *The Probabilistic Method, Fourth Edition*. Wiley, 2016.
- [4] K. Appel and W. Haken. Every planar map is four colorable. *Bull. Amer. Math. Soc.*, 82(5):711–712, 1976.
- [5] J. Bang-Jensen, S. Bessy, F. Havet, and A. Yeo. Bipartite spanning subdigraphs induced by 2-partitions, 2017, submitted.
- [6] J. Bang-Jensen, S. Bessy, F. Havet, and A. Yeo. Out-degree reducing 2-partitions of digraphs, 2017, submitted.
- [7] J. Bang-Jensen, N. Cohen, and F. Havet. Finding good 2-partitions of digraphs II. Enumerable properties. *Theoretical Computer Science*, 640:1–19, 2016.
- [8] J. Bang-Jensen and G. Gutin. *Digraphs: Theory, Algorithms and Applications, 2nd Edition*. Springer-Verlag, London, 2009.
- [9] J. Bang-Jensen and F. Havet. Finding good 2-partitions of digraphs I. Hereditary properties. *Theoretical Computer Science*, 636:85–94, 2016.
- [10] J. Bang-Jensen and F. Havet. Tournaments and semicomplete digraphs. In J. Bang-Jensen and G. Gutin, editors, *Classes of directed graphs*, chapter 2. Springer Verlag, London, to appear.
- [11] J. Beck. Roth’s estimate of the discrepancy of integer sequences is nearly optimal. *Combinatorica*, pages 319–325, 1981.
- [12] J.A. Bondy and U.S.R. Murty. *Graph Theory*, volume 244 of *Graduate Texts in Mathematics*. Springer-Verlag, Berlin, 2008.
- [13] P. Heggernes and Telle J.A. Partitioning graphs into generalized dominating sets. *Nordic J. Comput.*, 5:128–142, 1998.
- [14] L. Lovász. Coverings and coloring of hypergraphs. *Congr. Numer.*, 8:3–12, 1973.
- [15] J.W. Moon. On subtournaments of a tournament. *Can. Math. Bull.*, 9:297–301, 1966.
- [16] T.J. Schaefer. The complexity of satisfiability problems. In *Proceedings of the 10th Annual ACM Symposium on Theory of Computing (STOC 10)*, pages 216–226, New York, 1978. ACM.
- [17] J. H. Spencer. Six standard deviations suffice. *Transactions of the Amer. Math. Soc.*, 289:679–706, 1985.
- [18] C. Thomassen. Disjoint cycles in digraphs. *Combinatorica*, 3(3-4):393–396, 1983.
- [19] C. Thomassen. Even cycles in directed graphs. *Eur. J. Combin.*, 6(1):85–89, 1985.
- [20] D. Yang, Y. Bai, G. Wang, and J. Wu. On splitting digraphs. ArXiv:1707.03600, 2017.