

# Inverting Primes in Weinstein geometry

Joint w Sylwan, Tanaka

## I Localization in topology

Sullivan, Quillen... for any set of primes  $P$ ,  
there is a functor


$$(\ )_P : \text{Spaces} \rightarrow \text{Spaces}$$

" (w/htpy,  $\pi_i = 0$ )

1) invariants localize  $H_i(X_P; \mathbb{Z}) \cong H_i(X; \mathbb{Z}) \otimes \mathbb{Z}[\frac{1}{P}]$


2) construction of  $X_P$  depends on CW presentation  
of  $X$  - but homotopy type independent

3) idempotent  $(X_p)_p \cong X_p$



## II Symplectic localization

• symplectic analog of CW complex is  
Weinstein domain  $X^{2n}$



•  $P$  set of primes

Thm (Abouzaid-Segal) given  $X^{2n}$ ,  $n \geq 6$ ,  
exists  $X_p^{2n}$  diffeo to  $X$  s.t.

symplectic cohomology  $\leftarrow SH(X) \leftarrow$  not iso  $\leftarrow H^*(X)$

$SH(X_p) \cong SH(X) \otimes \mathbb{Z}[\frac{1}{p}]$   $\leftarrow H^*(X_p)$

Idea: modify vanishing cycles in  
Lefschetz fibration for  $X$

- not clear if  $X_p$  is independent of  
loc fib / Weinstein presentation of  $X$
- not clear if  $(X_p)_p \cong X_p$

Thm (Cieliebak-Eliashberg, Murphy)

given  $X^{2n}$ ,  $n \geq 3$ , exists  $X_{flex}^{2n}$  **flexible**

- $W(X_{flex}) \cong 0$ , **wrapped Fukaya**
- $X_{flex}$  is d. hco to  $X$
- h-principle: if  $X, Y$  d. hco,  $X_{flex}, Y_{flex}$  are symplectomorphic

$\Rightarrow X_{flex}$  is independent of pres of  $X$   
and  $(X_{flex})_{flex} \cong X_{flex}$  symplectomorphism  
or Weinstein homotopy

- $( )_{flex}$  is local modification  
of attaching spheres

Ex,  $TS^n \rightarrow TS^n_{flex}$



$\int$  Weinstein homotopy

flex  $\longrightarrow$



loose chat

Weinstein hty  
by Murphy's



think singular Lagrangian

Thm (L. with Sylvan) given  $X^{2n}$ ,  $n \geq 5$ , exists


- a Weinstein subdomain  $X_p \subset X^{2n}$  d.f.f. to  $X$
- $X_p \subset X_q$  if  $p \supset q$  or  $0 \in p$

- wrapped Fukaya  $TWW(X_p) \cong TWW(X) \otimes \mathbb{Z}[\frac{1}{p}]$

- local modification of attaching spheres

-  $X_0 = X_{\text{lex}}$ ,  $W(X|\mathbb{Z}[\frac{1}{0}]) \cong 0 \cong W(X_{\text{lex}})$

Ex.



order cannot be reversed

if  $T^2 \subset T^3$ , then Uiterbo ring map

$$SH(T^3; \mathbb{Z}/3) \not\rightarrow SH(T^2; \mathbb{Z}/3)$$

$$\cong SH(T^2; \mathbb{Z}/3)$$

Thm (L. w. Sullivan) if  $M^n$  is simply-connected and spin, then any Weinstein subdomain  $X \subset TM$  has  $TwW(X) \cong TwW(TM_p)$  for some set of primes  $p$

- not true in general that any  $X_0 \subset X$  has  $W(X_0) \cong W(X)[\frac{1}{p}]$

Ex.  $TN \subset TN \hookrightarrow TM$

Q. how unique is  $X_p$ ?

- don't affect Fukaya cat
- Two "symplectically trivial" operations
- subcritical handle attachment,  $n$ -principle  
 $X^n \rightsquigarrow X^n \cup H^k, \quad k < n$
  - stabilization ← sector  
 $X^n \rightsquigarrow X^n \times T^*D^1$

Thm (L. w. Sylvan, Tanaka)

- if  $X, X'$  are symplectomorphic  
 then  $X_p, X'_p$  are symplectomorphic  
 up to stabilization and subcritical handles
- $(X_p)_p \cong X_p$  up to stabilization  
 and subcritical handles
- functoriality  $\rightarrow$  construct  $W_{\text{ein}}^{\text{stab}}[\text{sub}^*]$

# III Construction

· given a Lagrangian disk  $D^n \subset X^n$   
can form subdomain  $X \setminus D \subset X$



cutting out

twisted complexes

Theorem (Ganatra-Pardon-Sherlock)  
 $\text{Tw}W(X \setminus D) \cong \text{Tw}W(X) / D$

→  
categorical localization  
by object  $D$

and Lagrangian cocores  $C_x \subset X^n$   
generate, i.e.  $\text{Tw}W(X) \cong \text{Tw}C_x$


## Abouzaid-Seidel disks

· given smooth subdomain  $U \subset S^{n-1}$ ,  
 $D_U^n \subset T^*D^n$  such that  
reduced, symplectic twisted complex

$$D_U^n \cong \tilde{C}^\infty(U) \otimes T^*D^n \text{ in } \text{Tw}W(T^*D^n)$$



•  $D_n = \Gamma df_u$ ,  $f_u: D^n \rightarrow \mathbb{R}$ ,  $\lim_{x \rightarrow \partial D^n} f(x) = -\infty$




• take  $U \subset S^{n-1}$  to be  $p$ -Moore space so  $\tilde{C}^n(U) \cong \mathbb{Z}[1] \xrightarrow{p} \mathbb{Z}$ , call  $D_p$

$U: \text{[scribble]} \xrightarrow{S^1 \cup D^2} \mathbb{R}^5 \hookrightarrow S^5$   
 $\downarrow S^4$

Carving out

• remove  $D_p$  near cocores  $C_x$  of  $X$



so that  $X_p, X$  are d.f.feo

(and reattach flexible handles)

• then  $W(X_p) = W(X \setminus D_p)$   
 $\stackrel{\text{GPS}}{\cong} W(X) / D_p$   
 $\cong W(X) / \{\mathbb{Z}[1] \xrightarrow{p} \mathbb{Z}\} \otimes C_x$



$$\begin{aligned} &\cong W(X) / \{\mathbb{Z}[1] \xrightarrow{p} \mathbb{Z}\} \otimes W(X) \\ &\cong W(X) \left[ \frac{1}{p} \right] \end{aligned}$$

forcing  $\{\mathbb{Z}[1] \xrightarrow{p} \mathbb{Z}\}$  to be acyclic  
 $\Leftrightarrow$  making  $p$  be invertible

### IV Classifying subdomains

$X \subset T^*M$  Weinstein subdomain

$\Leftrightarrow$  exists disks  $D_i$  s.t.

$$T^*M \setminus D_i = X \cup \text{subcritical handles}$$

$\rightarrow Tw W(X) \cong W(T^*M) / D_i$  for  
 some Lagrangian disks  $D_i$

$$W(T^*M) \xrightarrow{\text{disks}} W(T^*M) \xrightarrow{\text{embedded}} Tw W(T^*M) \cong Tw T_2^*M \cong Tw C_{-1}(M)$$

Q. which twisted complexes are iso to actual Lagrangians in  $X$ ?

Ex.  $S^n \cong T_2^*S^n[n] \xrightarrow{\gamma} T_2^*S^n, \gamma \in HW^{i-n}(T_2^*S^n, T_2^*S^n)$   
 higher degree

$$D_2 \cong T_2^* S^n [1] \xrightarrow{\cong} T_2^* S^n \quad \mathbb{Z} \in \text{HW}^0(T_2^* S^n, T_2^* S^n)$$

↑ " 2-Id


2-chiral complex  $\otimes T_2^* S^n$

Thm (L. w Sylvan) if  $M$  is spin, simply-connected,  
 any Lagrangian LCTM that is <sup>embedded, exact</sup>  
 null-homotopic is isomorphic to  $C^* \otimes T_2^* M$   
↑ topological ↑ 2-chiral complex

Ex image of  $W(T^* S^n) \hookrightarrow \text{Tw } W(T^* S^n)$   
 $n \geq 5,$   
 $T_2^* S^n [n] \xrightarrow{\cong} T_2^* S^n \cong S^n$   
 $C^* \otimes T_2^* S^n \cong \mathbb{Z} D_u^n$


• not all twisted complexes geometric  
Ex.  $S^n [1] \xrightarrow{5} S^n$

Post-talk notes



$H_*(S^n, \{F \ll 0\})$   
 $\text{WH}(S^n, \mathcal{D}_{\text{def}}) \stackrel{\text{Flav}}{\cong} H_{\text{non}}^a(F)$   
 $\text{WH}(S^n, T_0^* S^n) \cong \mathbb{Z}$


$$\text{WH}(S^1, D_u) \cong H^1(U)$$



$\partial(TD^n) \supset \partial D^n = \text{Legendrian unknot}$


$\partial D_u^n = \text{Legendrian unknot}$

$\partial D^n, \partial D_u^n$  are Legendrian linked



$U = \text{neighborhood of Morse spec in } S^{n-1}$

$$\begin{aligned} TD^n &\longleftrightarrow TS^n \\ T^*D^n &\longrightarrow T^*_2 S^n \\ D^n &\hookrightarrow S^n \end{aligned}$$



remove this

