

# SMALL DOUBLING IN ORDERED GROUPS

GREGORY FREIMAN, MARCEL HERZOG, PATRIZIA LONGOBARDI,  
AND MERCEDE MAJ

ABSTRACT. We prove that if  $S$  is finite set which generates a non-abelian ordered group, then  $|S^2| \geq 3|S| - 2$ . This generalizes a classical result from the theory of set addition.

## 1. INTRODUCTION

The structure theory of set addition, or Freiman-type theory, is an area founded by the first named author some time ago, and which concerns the structure of subsets of groups having so-called small ‘doubling’ (see [F]). This area is very popular (see [B],[C],[GR],[GT],[HLS],[R],[S] and [T]) and this paper contributes to the current programme of trying to understand what happens when we move from the abelian to nonabelian setting.

First we mention the following theorem, which is a classical result in the theory of set addition.

**Theorem 1.1.** *Let  $S$  be a finite subset of an ordered group. Then*

$$|S^2| \geq 2|S| - 1.$$

*Proof.* Let  $S = \{x_1, x_2, \dots, x_k\}$ , with  $x_1 < x_2 < \dots < x_k$ . Then:

$$x_1^2 < x_1x_2 < x_2^2 < x_2x_3 < x_3^2 < \dots < x_{k-1}^2 < x_{k-1}x_k < x_k^2$$

and each of these elements belongs to  $S^2$ . Hence  $|S^2| \geq 2k - 1 = 2|S| - 1$ , as required.  $\square$

This result is best possible as can be seen by considering geometric progressions. However, the critical examples (geometric progressions)

---

2000 *Mathematics Subject Classification.* Primary: 20F60,20F18; Secondary: 20F99.

*Key words and phrases.* ordered groups, finite subsets, small doubling, nilpotent groups.

The second author is grateful to the Department of Mathematics and Informatics of the University of Salerno for its hospitality and support, while this investigation was carried out.

are abelian in character and the main result of this paper is the following theorem, which is a strengthening of Theorem 1.1 if the group generated by  $S$  is nonabelian.

**Theorem 1.2.** *Let  $S$  be a finite subset of an ordered group, which generates a nonabelian subgroup. Then*

$$|S^2| \geq 3|S| - 2.$$

Theorem 1.2 can be restated in the following Freiman-type equivalent form.

**Theorem 1.3.** *Let  $S$  be a finite subset of an ordered group and suppose that*

$$|S^2| \leq 3|S| - 3.$$

*Then  $S$  generates an abelian subgroup.*

This result is best possible, so that there is an ordered group  $G$  and a subset  $S$  generating a nonabelian group with  $|S^2| = 3|S| - 2$ .

We prove Theorem 1.3 in Section 3. Under a bit stronger assumption, we obtain the following extension of Freiman's Theorem 1.9 in [F].

**Corollary 1.4.** *Let  $S$  be a finite subset of an ordered group  $G$  and suppose that*

$$t = |S^2| \leq 3|S| - 4.$$

*Then there exist  $x_1, g \in G$ , such that  $g > 1$ ,  $gx_1 = x_1g$  and  $S$  is a subset of the geometric progression*

$$\{x_1, x_1g, x_1g^2, \dots, x_1g^{t-|S|}\}.$$

Finally we mention the following interesting result concerning ordered groups, which is proved in Section 2.

**Corollary 1.5.** *Let  $S$  be a finite subset of an ordered group  $G$ . Then*

$$N_G(S) = C_G(S).$$

Since the class of ordered groups contains the class of *torsion-free nilpotent groups*, our results hold in particular for finite subsets of torsion-free nilpotent groups.

We conclude this section with the following basic definition.

**Definition 1.6.** *If  $S, T$  are subsets of a group  $G$ , then we denote*

$$ST = \{st \mid s \in S, t \in T\} \quad \text{and} \quad S^2 = \{s_1s_2 \mid s_1, s_2 \in S\}.$$

*If  $S = \{s\}$ , then we denote  $ST$  by  $sT$  and if  $T = \{t\}$ , then we write  $St$  instead of  $S\{t\}$ . If  $G$  is an additive group, then we denote*

$$2S = \{s_1 + s_2 \mid s_1, s_2 \in S\}.$$

## 2. FINITE SUBSETS OF ORDERED GROUPS

We begin this section with the definitions of ordered groups and of orderable groups. We recall some properties of these groups that we shall use in this paper, and we mention some interesting examples of orderable groups.

In the second part of this section we investigate finite subsets in ordered groups.

**Definition 2.1.** *Let  $G$  be a group and suppose that a total order relation  $<$  is defined on the set  $G$ . We say that  $(G, <)$  is an ordered group if for all  $a, b, x, y \in G$ , the inequality  $a < b$  implies that  $xy < xby$ .*

*A group  $G$  is orderable if there exists a total order relation  $<$  on the set  $G$ , such that  $(G, <)$  is an ordered group.*

The following properties of ordered groups follow easily from the definition (we apply the notation of the definition and denote by 1 the unit element of  $G$ ).

- If  $a < b$  and  $n$  is a positive integer, then  $a^n < b^n$  and  $a^{-n} > b^{-n}$ .
- If  $a < 1$ , then  $x^{-1}ax < 1$ .
- $G$  is torsion-free.
- If  $a, x \in G$  and  $a = x^{-1}a^{-1}x$ , then  $a = 1$ .

The next lemma due to B.H. Neumann (see [N]) will be very useful in the sequel.

**Lemma 2.2.** *Let  $(G, <)$  be an ordered group and let  $a, b \in G$ . If  $[a^n, b] = 1$  for some integer  $n \neq 0$ , then  $[a, b] = 1$ .*

*Proof.* For each integer  $m > 0$  we have the following identities:

$$[a^m, b] \equiv (a^{-(m-1)}[a, b]a^{m-1})(a^{-(m-2)}[a, b]a^{m-2}) \cdots \\ (a^{-1}[a, b]a^1)(a^0[a, b]a^0)$$

and

$$[a^{-m}, b] \equiv \prod_{k=-m}^{-1} (a^{-k}[a, b]^{-1}a^k).$$

Suppose that  $[a, b] > 1$  ( $[a, b] < 1$ ). Since  $[a^m, b]$  is a product of conjugates of  $[a, b]$ , each of which is  $> 1$  ( $< 1$ ), it follows that  $[a^m, b] > 1$  ( $[a^m, b] < 1$ ). Similarly, it follows that  $[a^{-m}, b] < 1$  ( $[a^{-m}, b] > 1$ ). Hence if  $[a, b] \neq 1$ , then  $[a^n, b] \neq 1$  and the result follows.  $\square$

There are many examples of orderable groups. An **abelian group** is orderable if and only if it is torsion-free, by a theorem of F.W. Levi

(see [L]). K. Iwasawa (see [I]), A.I. Mal'cev (see [MA]) and B.H. Neumann (see [N]) proved, independently, that the class of ordered groups contains the class of **torsion-free nilpotent groups**.

Other examples of solvable orderable groups can be obtained using the following theorem of Kargapolov (see [K]).

**Theorem 2.3.** *A torsion-free group  $G$  has the property that every full order for any subgroup of  $G$  can be extended to some full order of  $G$  if and only if there exists a normal abelian subgroup  $A$  of  $G$  such that  $G/A$  is abelian and for any  $a \in A$  and  $b \in G \setminus A$ , there exist **positive** integers  $m, n$ ,  $m \neq n$ , such that  $(a^m)^b = a^n$ .*

More information concerning ordered groups may be found, for example, in [GL] and in [BMR].

We now prove an important proposition concerning finite subsets in ordered groups.

**Proposition 2.4.** *Let  $(G, <)$  be an ordered group and let  $S$  be a finite subset of  $G$  of size  $k$ . If  $y \in G \setminus C_G(S)$ , then*

$$|yS \cup Sy| \geq k + 1.$$

*In particular, there exist  $x_i, x_j \in S$  such that  $yx_i \notin Sy$  and  $x_jy \notin yS$ .*

*Proof.* Suppose, to the contrary, that  $yS = Sy$ . Since  $y \notin C_G(S)$ , there exists  $x_1 \in S$  such that

$$yx_1 \neq x_1y.$$

As  $yS = Sy$ , there exists  $x_2 \in S$  such that  $x_2 \neq x_1$  and  $yx_1 = x_2y$ . Suppose that there exist  $x_1, x_2, \dots, x_t \in S$  such that

$$(1) \quad \begin{aligned} yx_1 &= x_2y \\ yx_2 &= x_3y \\ &\vdots \\ yx_{t-1} &= x_ty \end{aligned}$$

where  $x_i = x_j$  if and only if  $i = j$ .

Since  $yS = Sy$ , there exists  $x_{t+1} \in S$  such that

$$yx_t = x_{t+1}y.$$

We claim that  $x_{t+1} \notin \{x_1, x_2, \dots, x_t\}$ . Indeed, if  $x_{t+1} = x_u$  for some integer  $u$ ,  $1 \leq u \leq t$ , then by (1)

$$x_t = y^{-1}x_{t+1}y = y^{-1}x_u y = y^{-2}x_{u+1}y^2 = \dots = y^{-(t-u+1)}x_t y^{t-u+1}$$

and hence  $[x_t, y^{t-u+1}] = 1$ . It follows by Lemma 2.2 and (1) that  $yx_t = x_t y = yx_{t-1}$ . But then  $x_t = x_{t-1}$ , in contradiction to (1). This

proves our claim. Since this procedure can be carried out indefinitely, we have reached a contradiction to the finiteness of  $S$ . Hence  $yS \neq Sy$  and the proposition follows.  $\square$

From Proposition 2.4 we derive Corollary 1.5.

**Corollary 1.5.** *Let  $S$  be a finite subset of an ordered group  $G$ . Then*

$$N_G(S) = C_G(S).$$

*Proof.* If  $y \in N_G(S)$ , then  $yS = Sy$  and it follows from Proposition 2.4 that  $y \in C_G(S)$ . The opposite containment is trivial.  $\square$

### 3. THE MAIN RESULTS

In this section we prove our main results and some corollaries. First we prove Theorem 1.3.

**Theorem 1.3.** *Let  $S$  be a finite subset of an ordered group  $G$  and suppose that*

$$(*) \quad |S^2| \leq 3k - 3.$$

*Then  $S$  generates an abelian subgroup.*

*Proof.* Let  $S = \{x_1, x_2, \dots, x_k\}$ , with  $x_1 < x_2 < \dots < x_k$ .

If  $k = 2$ , then  $|S^2| \leq 3$ . As  $x_1^2 < x_1x_2 < x_2^2$ , it follows that  $S^2 = \{x_1^2, x_1x_2, x_2^2\}$  and we must have  $x_2x_1 = x_1x_2$ , as required.

So assume that  $k > 2$  and that all subsets  $X$  of  $G$  satisfying  $2 \leq |X| < k$  and  $|X^2| \leq 3|X| - 3$  generate an abelian subgroup. Assume, moreover, that  $\langle S \rangle$  is **nonabelian**. Our aim is to reach a contradiction.

Let  $i$  be the **maximal** integer such that

$$A = \{x_1, x_2, \dots, x_i\}$$

generates an **abelian** subgroup. Then

$$(i) \quad 1 \leq i < k, \quad x_{i+1} \notin C_G(A), \quad x_{i+1} \notin \langle A \rangle$$

and there exists  $x_j \in A$  such that

$$(ii) \quad x_{i+1}x_j \neq x_jx_{i+1}.$$

Let  $x_j$  be the **maximal** such element of  $A$ . Then

$$(iii) \quad x_a \in C_G(x_{i+1}) \quad \text{for each } x_a \in A \text{ satisfying } x_a > x_j.$$

Moreover, it follows from (i) that

$$(iv) \quad A^2 \cap (x_{i+1}A \cup Ax_{i+1}) = \emptyset.$$

Write

$$D = \{x_{i+1}, x_{i+2}, \dots, x_k\}.$$

If  $|D| = 1$ , then  $i = k - 1$  and the order in  $S$  implies that  $x_k^2 \notin A^2 \cup (x_{i+1}A \cup Ax_{i+1})$ . Thus, by (iv), (ii), Theorem 1.1 and Proposition 2.4, we get that

$$\begin{aligned} |S^2| &\geq |A^2| + |x_{i+1}A \cup Ax_{i+1}| + |\{x_k^2\}| \geq (2i - 1) + (i + 1) + 1 \\ &= 3i + 1 = 3(k - 1) + 1 = 3k - 2 \end{aligned}$$

in contradiction to (\*).

So assume that  $|D| \geq 2$ . We claim that

$$|D^2| \leq 3|D| - 3.$$

First we notice, that the order in  $S$  implies that

$$(v) \quad D^2 \cap (A^2 \cup x_{i+1}A \cup Ax_{i+1}) = \emptyset.$$

This observation, together with (iv), (\*), (ii), Theorem 1.1 and Proposition 2.4, yields the following inequality:

$$\begin{aligned} |D^2| &\leq |S^2| - |A^2| - |x_{i+1}A \cup Ax_{i+1}| \leq (3k - 3) - (2i - 1) - (i + 1) \\ &= 3(k - i) - 3 = 3|D| - 3. \end{aligned}$$

This proves our claim.

Since  $2 \leq |D| < k$ , it follows by the inductive assumption that  $\langle D \rangle$  is **abelian**. In particular,

$$(vi) \quad \langle D \rangle \leq C_G(x_{i+1}).$$

This implies, in view of (ii), that

$$(vii) \quad D^2 \cap (x_j D \cup D x_j) = \emptyset.$$

We claim that

$$(viii) \quad Ax_{i+1} \cap x_j D = \{x_j x_{i+1}\}.$$

Indeed, suppose that

$$(ix) \quad x_a x_{i+1} = x_j x_d \quad \text{for some } x_a \in A \text{ and } x_d \in D.$$

If  $x_a > x_j$ , then it follows by (ix), (iii) and (vi) that  $x_j \in \langle x_a, x_{i+1}, x_d \rangle \leq C_G(x_{i+1})$ , in contradiction to (ii). On the other hand, if  $x_a < x_j$ , then it follows by (ix) that  $x_{i+1} > x_d$ , which is impossible, since  $x_{i+1}$  is the smallest element in  $D$ . Thus  $x_a = x_j$ ,  $x_d = x_{i+1}$  and our claim follows. Since  $|Ax_{i+1}| = |A| = i$  and  $|x_j D| = |D| = k - i$ , (viii) implies that

$$(x) \quad |Ax_{i+1} \cup x_j D| = k - 1.$$

We also claim that

$$(xi) \quad A^2 \cap (x_j D \cup D x_j) = \emptyset.$$

Indeed, suppose that there exist  $x_a, x_b \in A$  and  $x_d \in D$  satisfying

$$x_a x_b = x_j x_d.$$

Since  $x_b < x_d$ , it follows that  $x_a > x_j$ . But  $\langle A \rangle$  is abelian, so  $x_a x_b = x_b x_a$  and similarly we get  $x_b > x_j$ . Thus, by (iii) and (vi),  $x_j \in \langle x_a, x_b, x_d \rangle \leq C_G(x_{i+1})$ , in contradiction to (ii). Hence  $A^2 \cap x_j D = \emptyset$  and a similar proof yields  $A^2 \cap D x_j = \emptyset$ . Thus our claim holds.

It follows by (iv), (v), (vii) and (xi) that

$$|A^2 \cup D^2 \cup A x_{i+1} \cup x_j D| = |A^2| + |D^2| + |A x_{i+1} \cup x_j D|$$

and hence, by Theorem 1.1 and (x), we get

$$|A^2 \cup D^2 \cup A x_{i+1} \cup x_j D| \geq (2i - 1) + (2(k - i) - 1) + (k - 1) = 3k - 3.$$

Thus, by (\*),

$$(xii) \quad S^2 = A^2 \cup D^2 \cup A x_{i+1} \cup x_j D.$$

Consider now the element  $x_{i+1} x_j \in S^2$ . By (v),  $x_{i+1} x_j \notin D^2$  and by (iv),  $x_{i+1} x_j \notin A^2$ .

Suppose, first, that  $x_{i+1} x_j \in A x_{i+1}$ . Then

$$(xiii) \quad x_{i+1} x_j = x_a x_{i+1} \quad \text{for some } x_a \in A.$$

If  $x_a > x_j$ , then by (xiii) and (iii)  $x_j \in C_G(x_{i+1})$ , in contradiction to (ii). Again by (ii)  $x_a \neq x_j$ . Hence  $x_a < x_j$ .

By (ii) and Proposition 2.4, there exists  $x_b \in A$  such that  $x_{i+1} x_b \notin A x_{i+1}$ . Since  $x_{i+1} x_j \in A x_{i+1}$ , we know that  $x_b \neq x_j$  and if  $x_b > x_j$ , then (iii) implies that  $x_{i+1} x_b = x_b x_{i+1} \in A x_{i+1}$ , a contradiction. Hence  $x_b < x_j$ .

Since  $x_{i+1} x_b \notin A x_{i+1}$  and since by (iv) and (v), also  $x_{i+1} x_b \notin A^2 \cup D^2$ , it follows by (xii) that  $x_{i+1} x_b \in x_j D$  and there exists  $x_d \in D$  such that  $x_j x_d = x_{i+1} x_b$ . Since  $\langle A \rangle$  is abelian, it follows that  $x_j x_d x_j = x_{i+1} x_b x_j = x_{i+1} x_j x_b$ . As by (xiii)  $x_{i+1} x_j = x_a x_{i+1}$ , we get  $x_j x_d x_j = x_a x_{i+1} x_b$  and  $x_j > x_b$  implies that  $x_j x_d < x_a x_{i+1}$ . But  $x_j > x_a$  and  $x_d \geq x_{i+1}$ , so  $x_j x_d > x_a x_{i+1}$ , a contradiction.

Suppose, finally, that  $x_{i+1} x_j \in x_j D$ . It follows that

$$x_{i+1} x_j = x_j x_d \quad \text{for some } x_d \in D.$$

By (ii) and Proposition 2.4 there exists  $x_f \in D$  such that  $x_f x_j \notin x_j D$ . Since  $x_{i+1} x_j \in x_j D$ , we must have  $x_{i+1} < x_f$ .

Now,  $x_f x_j \notin x_j D$  and it follows from (vii) and (xi) that  $x_f x_j \notin D^2 \cup A^2$ . Hence by (xii) we must have  $x_f x_j \in A x_{i+1}$ . Thus

$$(xiv) \quad x_a x_{i+1} = x_f x_j \quad \text{for some } x_a \in A.$$

Since  $x_f x_j \notin x_j D$ , we must have  $x_a \neq x_j$ . If  $x_a > x_j$ , then it follows by (iii) and (vi) that  $x_j \in \langle x_f, x_a, x_{i+1} \rangle \leq C_G(x_{i+1})$ , in contradiction to (ii). Hence  $x_a < x_j$ .

Since  $\langle D \rangle$  is abelian, it follows from (xiv) that  $x_{i+1} x_a x_{i+1} = x_{i+1} x_f x_j = x_f x_{i+1} x_j$ . Now  $x_{i+1} x_j = x_j x_d$ , so  $x_{i+1} x_a x_{i+1} = x_f x_j x_d$ . But  $x_{i+1} < x_f$ , so  $x_a x_{i+1} > x_j x_d$ . However,  $x_a < x_j$  and  $x_{i+1} \leq x_d$ , so  $x_a x_{i+1} < x_j x_d$ , a contradiction.

We have shown that  $x_{i+1} x_j \in S^2$  does not belong to  $A^2 \cup D^2 \cup A x_{i+1} \cup x_j D$ , in contradiction to (xii). It follows from this contradiction that  $\langle S \rangle$  is abelian.  $\square$

The result of the previous theorem is best possible. In fact, we exhibit in the following example an ordered group  $G$  and a finite subset  $S$  of  $G$ , such that  $\langle S \rangle$  is **nonabelian**,  $|S| = k \geq 2$  and  $|S^2| = 3k - 2$ .

**Example.**

Let  $G = A \rtimes \langle b \rangle$  be a semidirect product of an abelian subgroup  $A$ , isomorphic to the additive rational group  $(\mathbb{Q}, +)$ , with an infinite cyclic group  $\langle b \rangle$ , such that

$$b^{-1} a b = a^2 \quad \text{for each } a \in A.$$

Then  $G$  is torsion-free and it is orderable by Theorem 2.3.

Let  $a \in A \setminus \{1\}$  and let  $S = \{b, ba, ba^2, \dots, ba^{k-1}\}$ . Since  $ab = ba^2$ , it is easy to see that  $S^2 = \{b^2, b^2 a, b^2 a^2, b^2 a^3, \dots, b^2 a^{3k-3}\}$ . Thus  $\langle S \rangle$  is nonabelian and  $|S^2| = 3k - 2$ .

Theorem 1.3 is clearly equivalent to Theorem 1.2.

**Theorem 1.2.** *Let  $S$  be a finite subset of an ordered group, which generates a nonabelian subgroup. Then*

$$|S^2| \geq 3|S| - 2.$$

In order to prove Corollary 1.4 we need the following proposition, which extends Freiman's Theorem 1.9 in [F] from finite subsets of integers to finite subsets in ordered groups, generating abelian subgroups. Although this result is mentioned in [HLS], for sake of completeness we decided to report it with its proof.

**Proposition 3.1.** *Let  $S$  be a finite subset of an ordered group  $G$  and suppose that*

$$t = |S^2| \leq 3|S| - 4$$

*and  $S$  generates an abelian group. Then there exist  $x_1, g \in G$ , such that  $g > 1$ ,  $g x_1 = x_1 g$  and  $S$  is a subset of the geometric progression*

$$\{x_1, x_1 g, x_1 g^2, \dots, x_1 g^{t-|S|}\}.$$



*Proof.* Let  $S = \{x_1, x_2, \dots, x_k\}$ , with  $x_1 < x_2 < \dots < x_k$ . Clearly we may assume that  $G = \langle S \rangle$ , an abelian group.

Write  $y_i = x_1^{-1}x_i$  for  $i \in \{1, \dots, k\}$  and let  $K = \{1, y_2, \dots, y_k\}$ . Then  $1 < y_2 < y_3 < \dots < y_k$ ,  $S = x_1K$ ,  $S^2 = x_1^2K^2$  and  $|S^2| = |K^2|$ , so it suffices to prove the theorem when  $x_1 = 1$ . So assume that  $x_1 = 1$ . We argue by induction on  $k$ .

Assume first that  $k = 3$  and  $S = \{1, x_2, x_3\}$ . Then the elements  $1, x_2, x_2^2, x_2x_3, x_3^2$  are all different, since  $1 < x_2 < x_3$ . But  $|S^2| \leq 3 \cdot 3 - 4 = 5$ , so  $S^2 = \{1, x_2, x_2^2, x_2x_3, x_3^2\}$ , and the only possibility for  $x_3 \in S^2$  is  $x_3 = x_2^2$ . Hence  $S = \{1, g, g^2\}$  with  $g = x_2 > 1$  and  $2 = t - k$ , as required.

Suppose now that  $k > 3$  and that the theorem holds for subsets  $X$  of  $G$  satisfying  $3 \leq |X| < k$  and  $|X^2| \leq 3|X| - 4$ . Let  $g = x_kx_{k-1}^{-1}$ . Then  $g > 1$ , since  $x_k > x_{k-1}$ .

Assume first that for each  $i$ ,  $1 \leq i \leq k - 1$ , we have  $x_{i+1} = x_i g^{s_{i+1}}$ , where  $s_{i+1}$  are positive integers. Then, as  $x_1 = 1$ , it follows that  $x_{i+1} = g^{q_{i+1}}$ , where  $q_{i+1}$  are integers and  $0 < q_2 < q_3 < \dots < q_k$ . Let  $D = \{0, q_2, \dots, q_k\}$ . Since  $S = \{1, g^{q_2}, \dots, g^{q_k}\}$ , it follows that  $|2D| = |S^2| \leq 3k - 4$ . As  $q_{i+1}$  are integers, Freiman's Theorem 1.9 in [F] implies that  $D$  is a subset of the arithmetic progression  $\{0, q, 2q, \dots, (t - k)q\}$  for some integer  $q > 0$ . Thus  $S$  is a subset of the set  $\{1, g^q, g^{2q}, \dots, g^{(t-k)q}\}$ , where  $g^q > 1$ , as required.

Now assume that there exists an integer  $i$ ,  $1 \leq i \leq k - 1$ , such that for all positive integers  $l$

$$x_{i+1} \neq x_i g^l,$$

and let  $i$  be the **maximal** such integer. It follows by the definition of  $g$  that  $i < k - 1$ . Moreover, the definition of  $i$  implies that for each integer  $s$ ,  $i < s \leq k - 1$ , there exists a positive integer  $t_s$  such that  $x_k = x_s g^{t_s}$ , but for  $s = i$  such integer does not exist.

Let  $S' = S \setminus \{x_k\}$ . Obviously  $x_k^2, x_k x_{k-1} \in S^2 \setminus (S')^2$  because of the order in  $S$ . We also claim that  $x_k x_i \in S^2 \setminus (S')^2$ . In fact, if  $x_k x_i \in (S')^2$ , then  $x_k x_i = x_u x_v = x_v x_u$  for some integers  $u, v$ ,  $1 \leq u, v \leq k - 1$ . Since  $x_k > x_u$ , we must have  $i < v$  and similarly  $i < u$ . Therefore there exist positive integers  $t_u, t_v$  such that  $x_k = x_u g^{t_u}$  and  $x_k = x_v g^{t_v}$ . Thus  $x_k x_i = x_u x_v = x_k^2 g^{-(t_u + t_v)}$ , yielding  $x_k = x_i g^{t_u + t_v}$  with  $t_u + t_v > 0$ , in contradiction to the definition of  $i$ . This contradiction proves that  $x_k x_i \in S^2 \setminus (S')^2$ . Since  $x_k x_i \notin \{x_k^2, x_k x_{k-1}\}$ , it follows that

$$|(S')^2| \leq |S^2| - 3 \leq 3k - 4 - 3 = 3(k - 1) - 4 = 3|S'| - 4.$$

By induction there exists  $g' > 1$  such that each  $x_j$ ,  $1 < j \leq k - 1$ , satisfies  $x_j = (g')^{q_j}$  for some positive integer  $q_j$ . In particular, if

$x_w, x_j \in S'$  and  $x_w x_j > 1$ , then  $x_w x_j = (g')^{q_{w,j}}$ , where  $q_{w,j}$  is a positive integer.

Recall that  $x_k > 1$  and  $x_k^2 \notin (S')^2$ . We claim that if  $x_k \neq (g')^h$  for all positive integers  $h$ , then each  $x_b \in S'$  satisfies  $x_k x_b \notin (S')^2$ . Indeed, assume that this is not the case and  $x_k x_b = (g')^z$  for some positive integer  $z$ . Then  $x_k = (g')^l$  for some integer  $l$  and since  $x_k, g' > 1$ ,  $l$  is positive. We have reached a contradiction to our assumption. This proves our claim, and it follows that  $|S^2| - |(S')^2| \geq k$ . Thus

$$|(S')^2| \leq |S^2| - k \leq 3k - 4 - k = 2(k - 1) - 2 = 2|S'| - 2,$$

in contradiction to Theorem 1.1. Hence also  $x_k = (g')^{q_k}$  for some positive integer  $q_k$ . It follows from the order in  $S$  and from  $g' > 1$ , that  $0 < q_2 < q_3 < \dots < q_k$ . Applying again Freiman's Theorem 1.9 in [F] to  $D = \{0, q_2, \dots, q_k\}$ , it follows as above that  $S$  is as required.  $\square$

Corollary 1.4 follows immediately from Theorem 1.3 and Proposition 3.1.

**Corollary 1.4.** *Let  $S$  be a finite subset of an ordered group  $G$  and suppose that*

$$t = |S^2| \leq 3|S| - 4.$$

*Then there exist  $x_1, g \in G$ , such that  $g > 1$ ,  $gx_1 = x_1g$  and  $S$  is a subset of the geometric progression*

$$\{x_1, x_1g, x_1g^2, \dots, x_1g^{t-|S|}\}.$$

*Proof.* By Theorem 1.3,  $\langle S \rangle$  is abelian, and hence, by Proposition 3.1, it is a subset of a geometric progression, as stated.  $\square$

Acknowledgement. The authors are grateful to the referees for their constructive remarks.

#### REFERENCES

- [B] Y. Bilu, Structure of sets with small sumsets, *Asterisque* **258** (1999), 77-108.
- [BMR] R. Botto Mura and A. Rhemtulla, Orderable groups, Lecture Notes in Pure and Applied Mathematics, Marcel Dekker, Inc., New York and Basel, 1977.
- [C] M. Chang, A polynomial bound in Freiman's theorem. *Duke Math. J.* **113(3)** (2002), 399-419.
- [F] G. A. Freiman, Foundations of a structural theory of set addition. Translations of mathematical monographs, v. 37. American Mathematical Society, Providence, Rhode Island, 1973.
- [GL] A. M. W. Glass, Partially ordered groups, World Scientific Publishing Co., Series in Algebra, v. 7, 1999.
- [GR] B. J. Green and I. Z. Ruzsa, Freiman's theorem in an arbitrary abelian group, *J. Lond. Math. Soc. (2)* **75(1)** (2007), 163-175.
- [GT] B. J. Green and T. C. Tao, Freiman's theorem in finite fields via extremal set theory, *Combin. Probab. Comput.* **18** (2009), 335-355.

- [HLS] Y. O. Hamidoune, A. S. Llado and O. Serra, On subsets with small product in torsion-free groups, *Combinatorica* **18(4)** (1998), 529-540.
- [I] K. Iwasawa, On linearly ordered groups, *J. Math. Soc. Japan* **1** (1948), 1-9.
- [K] M. I. Kargapolov, Completely orderable groups, *Algebra i Logica (2)* **1** (1962), 16-21.
- [L] F. W. Levi, Arithmetische Gesetze im Gebiete diskreter Gruppen, *Rend. Circ. Mat. Palermo* **35** (1913), 225-236.
- [MA] A. I. Mal'cev, On ordered groups, *Izv. Akad. Nauk. SSSR Ser. Mat.* **13** (1948), 473-482.
- [N] B. H. Neumann, On ordered groups, *Amer. J. Math.* **71** (1949), 1-18.
- [R] I. Z. Ruzsa, Generalized arithmetic progressions and sumsets, *Acta Math. Hungar.* **65(4)** (1994), 379-388.
- [S] T. Sanders, A note on Freiman's theorem in vector spaces, *Combin. Probab. Comput.* **17(2)** (2008), 297-305.
- [T] T. C. Tao, Product set estimates for non-commutative groups, *Combinatorica* **28(5)** (2008), 547-594.

G. A. FREIMAN, DEPARTMENT OF MATHEMATICS  
 RAYMOND AND BEVERLY SACKLER FACULTY OF EXACT SCIENCES  
 TEL AVIV UNIVERSITY, TEL AVIV, ISRAEL  
*E-mail address:* `grisha@post.tau.ac.il`

M. HERZOG, DEPARTMENT OF MATHEMATICS  
 RAYMOND AND BEVERLY SACKLER FACULTY OF EXACT SCIENCES  
 TEL AVIV UNIVERSITY, TEL AVIV, ISRAEL  
*E-mail address:* `herzogm@post.tau.ac.il`

P. LONGOBARDI, DIPARTIMENTO DI MATEMATICA E INFORMATICA  
 UNIVERSITÀ DI SALERNO  
 VIA PONTE DON MELILLO, 84084 FISCIANO (SALERNO), ITALY  
*E-mail address:* `plongobardi@unisa.it`

M. MAJ, DIPARTIMENTO DI MATEMATICA E INFORMATICA  
 UNIVERSITÀ DI SALERNO  
 VIA PONTE DON MELILLO, 84084 FISCIANO (SALERNO), ITALY  
*E-mail address:* `mmaj@unisa.it`